

025250A

FUSED TETRAHYDROPYRIDINE DERIVATIVES AS MATRIX
METALLOPROTEINASE INHIBITORS



* 2 8 8 8 0 *

28880

PATENT TRADEMARK OFFICE

FUSED TETRAHYDROPYRIDINE DERIVATIVES AS MATRIX
METALLOPROTEINASE INHIBITORS

CROSS-REFERENCE TO RELATED APPLICATIONS

5 This application claims benefit of priority from United States Provisional
Patent Application Number 60/403,172, filed August 13, 2002.

FIELD OF THE INVENTION

10 This invention relates to fused tetrahydropyridine derivatives which inhibit
matrix metalloproteinase enzymes and thus are useful for treating diseases
resulting from MMP-mediated tissue breakdown such as heart disease, cardiac
insufficiency, inflammatory bowel disease, multiple sclerosis, osteo- and
rheumatoid arthritis, arthritis other than osteo- or rheumatoid arthritis, heart
failure, age-related macular degeneration, chronic obstructive pulmonary disease,
asthma, periodontal diseases, psoriasis, atherosclerosis, and osteoporosis.

BACKGROUND OF THE INVENTION

15 Matrix metalloproteinases (sometimes referred to as MMPs) are naturally
occurring enzymes found in most mammals. Over-expression and activation of
MMPs, or an imbalance between MMPs and inhibitors of MMPs, have been
suggested as factors in the pathogenesis of diseases characterized by the
breakdown of extracellular matrix or connective tissues.

20 Stromelysin-1 and gelatinase A are members of the MMP family. Other
members include fibroblast collagenase (MMP-1), neutrophil collagenase
(MMP-8), gelatinase B (92 kDa gelatinase) (MMP-9), stromelysin-2 (MMP-10),
stromelysin-3 (MMP-11), matrilysin (MMP-7), collagenase 3 (MMP-13),
TNF-alpha converting enzyme (TACE), and other newly discovered membrane-
25 associated matrix metalloproteinases (Sato H., Takino T., Okada Y., Cao J.,
Shinagawa A., Yamamoto E., and Seiki M., *Nature*, 1994;370:61-65). These

enzymes have been implicated with a number of diseases which result from breakdown of connective tissue, including such diseases as rheumatoid arthritis, osteoarthritis, osteoporosis, periodontitis, multiple sclerosis, gingivitis, corneal epidermal and gastric ulceration, atherosclerosis, neointimal proliferation which
5 leads to restenosis and ischemic heart failure, and tumor metastasis. A method for preventing and treating these and other diseases is now recognized to be by inhibiting matrix metalloproteinase enzymes, thereby curtailing and/or eliminating the breakdown of connective tissues that results in the disease states.

There is a catalytic zinc domain in matrix metalloproteinases that is
10 typically the focal point for inhibitor design. The modification of substrates by introducing zinc-chelating groups has generated potent inhibitors such as peptide hydroxamates and thiol-containing peptides. Peptide hydroxamates and the natural endogenous inhibitors of MMPs (TIMPs) have been used successfully to treat animal models of cancer and inflammation. MMP inhibitors have also been used
15 to prevent and treat congestive heart failure and other cardiovascular diseases, United States Patent No. 5,948,780.

A major limitation on the use of currently known MMP inhibitors is their lack of specificity for any particular enzyme. Recent data has established that specific MMP enzymes are associated with some diseases, with no effect on
20 others. The MMPs are generally categorized based on their substrate specificity, and indeed the collagenase subfamily of MMP-1, MMP-8, and MMP-13 selectively cleave native interstitial collagens, and thus are associated only with diseases linked to such interstitial collagen tissue. This is evidenced by the recent discovery that MMP-13 alone is over expressed in breast carcinoma, while
25 MMP-1 alone is over expressed in papillary carcinoma (see Chen et al., *J. Am. Chem. Soc.*, 2000;122:9648-9654).

Selective inhibitors of MMP-13 include a compound named WAY-170523, which has been reported by Chen et al., supra., 2000, and other compounds are reported in PCT International Patent Application Publication
30 numbers WO 01/63244; WO 00/09485; WO 01/12611; WO 02/34726; and WO 02/34753, and European Patent Application numbers EP 935,963 and EP 1,138,680. Further, United States Patent number 6,008,243 discloses inhibitors of MMP-13. However, no selective or nonselective inhibitor of MMP-13 has been


approved and marketed for the treatment of any disease in any mammal. Accordingly, the need continues to find new low molecular weight compounds that are potent and selective MMP inhibitors, and that have an acceptable therapeutic index of toxicity/potency to make them amenable for use clinically in the prevention and treatment of the associated disease states. An object of this invention is to provide a group of selective MMP-13 inhibitor compounds characterized as being fused tetrahydropyridine derivatives.

SUMMARY OF THE INVENTION

This invention provides a fused tetrahydropyridine derived compound defined by Formula I.

Accordingly, embodiments of the invention include:

1. A compound of Formula I



or a pharmaceutically acceptable salt thereof, or a pyrido-N-oxide thereof, wherein:

R¹ is independently selected from:

- C₅ or C₆ cycloalkyl-(C₁-C₈ alkylenyl);
- Substituted C₅ or C₆ cycloalkyl-(C₁-C₈ alkylenyl);
- C₈-C₁₀ bicycloalkyl-(C₁-C₈ alkylenyl);
- Substituted C₈-C₁₀ bicycloalkyl-(C₁-C₈ alkylenyl);
- 5- or 6-membered heterocycloalkyl-(C₁-C₈ alkylenyl);
- Substituted 5- or 6-membered heterocycloalkyl-(C₁-C₈ alkylenyl);
- 8- to 10-membered heterobicycloalkyl-(C₁-C₈ alkylenyl);
- Substituted 8- to 10-membered heterobicycloalkyl-(C₁-C₈ alkylenyl);
- Phenyl-(C₁-C₈ alkylenyl);
- Substituted phenyl-(C₁-C₈ alkylenyl);


- Naphthyl-(C₁-C₈ alkylenyl);
Substituted naphthyl-(C₁-C₈ alkylenyl);
5- or 6-membered heteroaryl-(C₁-C₈ alkylenyl);
Substituted 5- or 6-membered heteroaryl-(C₁-C₈ alkylenyl);
5 8- to 10-membered heterobiaryl-(C₁-C₈ alkylenyl);
Substituted 8- to 10-membered heterobiaryl-(C₁-C₈ alkylenyl);
Phenyl;
Substituted phenyl;
Naphthyl;
10 Substituted naphthyl;
5- or 6-membered heteroaryl;
Substituted 5- or 6-membered heteroaryl;
8- to 10-membered heterobiaryl; and
Substituted 8- to 10-membered heterobiaryl;
15 R² is independently selected from:
H;
C₁-C₆ alkyl;
Phenyl-(C₁-C₈ alkylenyl);
Substituted phenyl-(C₁-C₈ alkylenyl);
20 Naphthyl-(C₁-C₈ alkylenyl);
Substituted naphthyl-(C₁-C₈ alkylenyl);
5- or 6-membered heteroaryl-(C₁-C₈ alkylenyl);
Substituted 5- or 6-membered heteroaryl-(C₁-C₈ alkylenyl);
8- to 10-membered heterobiaryl-(C₁-C₈ alkylenyl);
25 Substituted 8- to 10-membered heterobiaryl-(C₁-C₈ alkylenyl);
Phenyl-O-(C₁-C₈ alkylenyl);
Substituted phenyl-O-(C₁-C₈ alkylenyl);
Phenyl-S-(C₁-C₈ alkylenyl);
Substituted phenyl-S-(C₁-C₈ alkylenyl);
30 Phenyl-S(O)-(C₁-C₈ alkylenyl);
Substituted phenyl-S(O)-(C₁-C₈ alkylenyl);
Phenyl-S(O)₂-(C₁-C₈ alkylenyl); and
Substituted phenyl-S(O)₂-(C₁-C₈ alkylenyl);

Each substituted R¹ and R² group contains from 1 to 4 substituents, each independently on a carbon or nitrogen atom, independently selected from:

- C₁-C₆ alkyl;
- CN;
- 5 CF₃;
- HO;
- (C₁-C₆ alkyl)-O;
- (C₁-C₆ alkyl)-S(O)₂;
- H₂N;
- 10 (C₁-C₆ alkyl)-N(H);
- (C₁-C₆ alkyl)₂-N;
- (C₁-C₆ alkyl)-C(O)O-(C₁-C₈ alkylenyl)_m;
- (C₁-C₆ alkyl)-C(O)O-(1- to 8-membered heteroalkylenyl)_m;
- (C₁-C₆ alkyl)-C(O)N(H)-(C₁-C₈ alkylenyl)_m;
- 15 (C₁-C₆ alkyl)-C(O)N(H)-(1- to 8-membered heteroalkylenyl)_m;
- H₂NS(O)₂-(C₁-C₈ alkylenyl);
- (C₁-C₆ alkyl)-N(H)S(O)₂-(C₁-C₈ alkylenyl)_m;
- (C₁-C₆ alkyl)₂-NS(O)₂-(C₁-C₈ alkylenyl)_m;
- 3- to 6-membered heterocycloalkyl-(G)_m;
- 20 Substituted 3- to 6-membered heterocycloalkyl-(G)_m;
- 5- or 6-membered heteroaryl-(G)_m;
- Substituted 5- or 6-membered heteroaryl-(G)_m;
- (C₁-C₆ alkyl)-S(O)₂-N(H)-C(O)-(C₁-C₈ alkylenyl)_m; and
- (C₁-C₆ alkyl)-C(O)-N(H)-S(O)₂-(C₁-C₈ alkylenyl)_m;
- 25 wherein each substituent on a carbon atom may further be independently selected from:

Halo; and
HO₂C;

- wherein 2 substituents may be taken together with a carbon atom to which they
- 30 are both bonded to form the group C=O;
- wherein two adjacent, substantially sp² carbon atoms may be taken together with a diradical substituent to form a cyclic diradical selected from:



5

R is H or C₁-C₆ alkyl;

G is CH₂; O, S, S(O); or S(O)₂;

m is an integer of 0 or 1;


Y¹ is CH₂, C(O), or S(O)₂;

10

Y² is C(O);

Y³ is N(R⁴); or

Y² and Y³ may be taken together to form a diradical group selected from:



R³ is independently selected from the groups:

15

H;

CH₃;

CH₃O;
CH=CH₂;
HO;
CF₃;
CN;
F; and
Cl;

5

R⁴ is independently selected from the groups:

H;
CH₃;
CH₃O;
HO;
CF₃; and
CN; and

10

15 wherein R⁴ is bonded to a carbon atom, R⁴ may further independently be
halo or CO₂H;

Q is selected from:

OC(O);
CH(R⁵)C(O);
OC(NR⁵);
CH(R⁵)C(NR⁵);
N(R⁵)C(O);
N(R⁵)C(S);
N(R⁵)C(NR⁵);
CH₂N(R⁵);
SC(O);
CH(R⁵)C(S);
SC(NR⁵);
trans-(H)C=C(H);
cis-(H)C=C(H);
C≡C;
CH₂C≡C;
C≡CCH₂;

20







25

30


$\text{CF}_2\text{C}\equiv\text{C}$; and

$\text{C}\equiv\text{CCF}_2$;

$\text{C}\equiv\text{CC}(\text{O})$;



; and



5

Each R^5 is independently selected from: H, $\text{C}_1\text{-C}_6$ alkyl, $\text{C}_3\text{-C}_6$ cycloalkyl; 3- to 6-membered heterocycloalkyl; phenyl; benzyl; and 5- or 6-membered heteroaryl;

X is O, S, N(H), or N($\text{C}_1\text{-C}_6$ alkyl);

10

Each V is independently C(H) or N;

wherein each $\text{C}_8\text{-C}_{10}$ bicycloalkyl is a bicyclic carbocyclic ring that contains 8-, 9-, or 10-member carbon atoms which are 5,5-fused, 6,5-fused, or 6,6-fused bicyclic rings, respectively, and wherein the ring is saturated or optionally contains one carbon-carbon double bond;

15

wherein each 8- to 10-membered heterobicycloalkyl is a bicyclic ring that contains carbon atoms and from 1 to 4 heteroatoms independently selected from 2 O, 1 S, 1 S(O), 1 S(O)₂, 1 N, 4 N(H), and 4 N($\text{C}_1\text{-C}_6$ alkyl), and wherein when two O atoms or one O atom and one S atom are present, the two O atoms or one O atom and one S atom are not bonded to each other, and wherein the ring is

20

saturated or optionally contains one carbon-carbon or carbon-nitrogen double bond, and wherein the heterobicycloalkyl is a 5,5-fused, 6,5-fused, or 6,6-fused bicyclic ring, respectively,

wherein each heterocycloalkyl is a ring that contains carbon atoms and from 1 to 4 heteroatoms independently selected from 2 O, 1 S, 1 S(O), 1 S(O)₂, 1 N, 4

N(H), and 4 N(C₁-C₆ alkyl), and wherein when two O atoms or one O atom and one S atom are present, the two O atoms or one O atom and one S atom are not bonded to each other, and wherein the ring is saturated or optionally contains one carbon-carbon or carbon-nitrogen double bond;

5 wherein each 5-membered heteroaryl contains carbon atoms and from 1 to 4 heteroatoms independently selected from 1 O, 1 S, 1 N(H), 1 N(C₁-C₆ alkyl), and 4 N, and each 6-membered heteroaryl contains carbon atoms and 1 or 2 heteroatoms independently selected from N, N(H), and N(C₁-C₆ alkyl), and 5- and 6-membered heteroaryl are monocyclic rings;

10 wherein each heterobiaryl contains carbon atoms and from 1 to 4 heteroatoms independently selected from 1 O, 1 S, 1 N(H), 1 N(C₁-C₆ alkyl), and 4 N, and where the 8-, 9-, and 10-membered heterobiaryl are 5,5-fused, 6,5-fused, and 6,6-fused bicyclic rings, respectively, and wherein at least 1 of the 2 fused rings of a bicyclic ring is aromatic, and wherein when the O

15 and S atoms both are present, the O and S atoms are not bonded to each other;


wherein with any (C₁-C₆ alkyl)₂-N group, the C₁-C₆ alkyl groups may be optionally taken together with the nitrogen atom to which they are attached to form a 5- or 6-membered heterocycloalkyl; and

20 wherein each group and each substituent recited above is independently selected.

2. The compound according to Embodiment 1, or a pharmaceutically acceptable salt thereof, wherein Y¹ is CH₂.
- 25 3. The compound according to Embodiment 1, or a pharmaceutically acceptable salt thereof, wherein Y¹ is C(=O).
4. The compound according to Embodiment 1, or a pharmaceutically acceptable salt thereof, wherein Y¹ is S(O)₂.
- 30 5. The compound according to Embodiment 1, or a pharmaceutically acceptable salt thereof, wherein Y² is C(=O).


6. The compound according to Embodiment 1, or a pharmaceutically acceptable salt thereof, wherein Y^3 is $N(R^4)$, wherein R^4 is as defined above.

5 7. The compound according to Embodiment 1, or a pharmaceutically acceptable salt thereof, wherein Y^2 and Y^3 are taken together to form the diradical group




, wherein R^3 and R^4 are as defined above.

10 8. The compound according to Embodiment 1, or a pharmaceutically acceptable salt thereof, wherein Y^2 and Y^3 are taken together to form the diradical group



, wherein R^3 is as defined above.

15 9. The compound according to Embodiment 1, or a pharmaceutically acceptable salt thereof, wherein Y^2 and Y^3 are taken together to form the diradical group



, wherein R^3 is H.

20 10. The compound according to any one of Embodiments 1 to 9, or a pharmaceutically acceptable salt thereof, wherein Q is $OC(O)$.

25 11. The compound according to any one of Embodiments 1 to 9, or a pharmaceutically acceptable salt thereof, wherein Q is $CH(R^5)C(O)$, wherein R^5 is as defined above.

12. The compound according to any one of Embodiments 1 to 9, or a pharmaceutically acceptable salt thereof, wherein Q is $\text{OC}(\text{NR}^5)$, $\text{N}(\text{R}^5)\text{C}(\text{NR}^5)$, or $\text{SC}(\text{NR}^5)$, wherein each R^5 is as defined above.
- 5 13. The compound according to any one of Embodiments 1 to 9, or a pharmaceutically acceptable salt thereof, wherein Q is $\text{CH}(\text{R}^5)\text{C}(\text{NR}^5)$, wherein each R^5 is as defined above.
- 10 14. The compound according to any one of Embodiments 1 to 9, or a pharmaceutically acceptable salt thereof, wherein Q is $\text{N}(\text{R}^5)\text{C}(\text{O})$, wherein R^5 is as defined above.
- 15 15. The compound according to any one of Embodiments 1 to 9, or a pharmaceutically acceptable salt thereof, wherein Q is $\text{CH}_2\text{N}(\text{R}^5)$, wherein R^5 is as defined above.
16. The compound according to any one of Embodiments 1 to 9, or a pharmaceutically acceptable salt thereof, wherein Q is $\text{SC}(\text{O})$.
- 20 17. The compound according to any one of Embodiments 1 to 9, or a pharmaceutically acceptable salt thereof, wherein Q is $\text{CH}(\text{R}^5)\text{C}(\text{S})$, wherein R^5 is as defined above.
- 25 18. The compound according to any one of Embodiments 1 to 9, or a pharmaceutically acceptable salt thereof, wherein Q is $\text{C}\equiv\text{C}$, $\text{CH}_2\text{C}\equiv\text{C}$, $\text{C}\equiv\text{CCH}_2$, $\text{CF}_2\text{C}\equiv\text{C}$, or $\text{C}\equiv\text{CCF}_2$.
- 30 19. The compound according to any one of Embodiments 1 to 9, or a pharmaceutically acceptable salt thereof, wherein Q is $\text{C}\equiv\text{CCH}_2$ or $\text{C}\equiv\text{CCF}_2$.
20. The compound according to any one of Embodiments 1 to 9, or a pharmaceutically acceptable salt thereof, wherein Q is $\text{C}\equiv\text{CC}(\text{O})$.

21. The compound according to any one of Embodiments 1 to 20, or a pharmaceutically acceptable salt thereof, wherein at least one of R^3 and R^4 is H.
- 5 23. The compound according to any one of Embodiments 1 to 20, or a pharmaceutically acceptable salt thereof, wherein at least one of R^3 and R^4 is F.
24. The compound according to any one of Embodiments 1 to 20, or a pharmaceutically acceptable salt thereof, wherein at least one of R^3 and R^4 is HO.
- 10 25. The compound according to any one of Embodiments 1 to 20, or a pharmaceutically acceptable salt thereof, wherein at least one of R^3 and R^4 is CH_3O .
- 15 26. The compound according to any one of Embodiments 1 to 25, or a pharmaceutically acceptable salt thereof, wherein at least one of R^1 is independently selected from:
- Phenyl-(C_1 - C_8 alkylenyl);
 - Substituted phenyl-(C_1 - C_8 alkylenyl);
 - 5- or 6-membered heteroaryl-(C_1 - C_8 alkylenyl);
 - 20 Substituted 5- or 6-membered heteroaryl-(C_1 - C_8 alkylenyl);
 - 8- to 10-membered heterobiaryl-(C_1 - C_8 alkylenyl); and
 - Substituted 8- to 10-membered heterobiaryl-(C_1 - C_8 alkylenyl); or
- at least one of R^2 is independently selected from:
- Phenyl-(C_1 - C_8 alkylenyl) $_m$;
 - 25 Substituted phenyl-(C_1 - C_8 alkylenyl) $_m$;
 - 5- or 6-membered heteroaryl-(C_1 - C_8 alkylenyl) $_m$;
 - Substituted 5- or 6-membered heteroaryl-(C_1 - C_8 alkylenyl) $_m$;
 - 8- to 10-membered heterobiaryl-(C_1 - C_8 alkylenyl) $_m$; and
 - Substituted 8- to 10-membered heterobiaryl-(C_1 - C_8 alkylenyl) $_m$;
- 30 wherein m is an integer of 0 or 1; and
- wherein each group and each substituent is independently selected.

27. The compound according to any one of Embodiments 1 to 26, or a pharmaceutically acceptable salt thereof, wherein R^1 is independently selected from:

- Phenyl-(C_1 - C_8 alkylene);
- 5 Substituted phenyl-(C_1 - C_8 alkylene);
- 5- or 6-membered heteroaryl-(C_1 - C_8 alkylene);
- Substituted 5- or 6-membered heteroaryl-(C_1 - C_8 alkylene);
- 8- to 10-membered heterobiaryl-(C_1 - C_8 alkylene); and
- Substituted 8- to 10-membered heterobiaryl-(C_1 - C_8 alkylene); and

10 R^2 is independently selected from:

- Phenyl-(C_1 - C_8 alkylene)_m;
- Substituted phenyl-(C_1 - C_8 alkylene)_m;
- 5- or 6-membered heteroaryl-(C_1 - C_8 alkylene)_m;
- Substituted 5- or 6-membered heteroaryl-(C_1 - C_8 alkylene)_m;
- 15 8- to 10-membered heterobiaryl-(C_1 - C_8 alkylene)_m; and
- Substituted 8- to 10-membered heterobiaryl-(C_1 - C_8 alkylene)_m;

wherein m is an integer of 0 or 1; and

wherein each group and each substituent is independently selected.

20 28. The compound according to any one of Embodiments 1 to 27, or a pharmaceutically acceptable salt thereof, wherein R^1 is independently selected from:

- Phenyl-(C_1 - C_8 alkylene);
- Substituted phenyl-(C_1 - C_8 alkylene);
- 25 5- or 6-membered heteroaryl-(C_1 - C_8 alkylene); and
- Substituted 5- or 6-membered heteroaryl-(C_1 - C_8 alkylene);

and R^2 is independently selected from:

- Phenyl-(C_1 - C_8 alkylene)_m;
- Substituted phenyl-(C_1 - C_8 alkylene)_m;
- 30 5- or 6-membered heteroaryl-(C_1 - C_8 alkylene)_m; and
- Substituted 5- or 6-membered heteroaryl-(C_1 - C_8 alkylene)_m;

wherein m is an integer of 0 or 1; and


wherein each group and each substituent is independently selected.

29. The compound according to any one of Embodiments 1 to 25, or a pharmaceutically acceptable salt thereof, wherein R^1 is independently selected from:

- 5 C_3 - C_6 cycloalkyl- $(C_1$ - C_8 alkylenyl);
 Substituted C_3 - C_6 cycloalkyl- $(C_1$ - C_8 alkylenyl);
 3- to 6-membered heterocycloalkyl- $(C_1$ - C_8 alkylenyl); and
 Substituted 3- to 6-membered heterocycloalkyl- $(C_1$ - C_8 alkylenyl) and
 10 wherein each group and each substituent recited above is independently selected.

30. The compound according to any one of Embodiments 1 to 25, or a pharmaceutically acceptable salt thereof, wherein R^1 is substituted phenyl- $(C_1$ - C_8 alkylenyl).

15 31. The compound according to any one of Embodiments 1 to 9 and 21 to 30, or a pharmaceutically acceptable salt thereof, wherein Q is selected from:



20 wherein V, X, and R^5 are as defined above.

32. The compound according to any one of Embodiments 1 to 31, or a pharmaceutically acceptable salt thereof, wherein each C_1 - C_8 alkylenyl is CH_2 , $C(CH_3)_2$, $C(=O)$, or CF_2 .

25

33. The compound according to any one of Embodiments 1 to 32, or a pharmaceutically acceptable salt thereof, wherein each C_1 - C_8 alkylenyl is CH_2 .


34. The compound according to any one of Embodiments 1 to 33, or a pharmaceutically acceptable salt thereof, wherein at least one substituent is selected from the groups:

- 5 CO₂H;
 CO₂CH₃;
 F;
 Cl;
 CN;
10 CF₃;
 HO;
 CH₃O; and
 CH₃; or

wherein at least two substituents are each F, F and CH₃O, or F and HO; or

15 wherein at least three substituents are selected from 2 F, 1 CH₃O, and 1 HO.

35. A compound of Formula II



II

or a pharmaceutically acceptable salt thereof, or a pyrido-N-oxide thereof,

20 wherein:

R¹ is independently selected from:

- C₅ or C₆ cycloalkyl-(C₁-C₈ alkylenyl);
 Substituted C₅ or C₆ cycloalkyl-(C₁-C₈ alkylenyl);
 C₈-C₁₀ bicycloalkyl-(C₁-C₈ alkylenyl);
25 Substituted C₈-C₁₀ bicycloalkyl-(C₁-C₈ alkylenyl);
 5- or 6-membered heterocycloalkyl-(C₁-C₈ alkylenyl);
 Substituted 5- or 6-membered heterocycloalkyl-(C₁-C₈ alkylenyl);
 8- to 10-membered heterobicycloalkyl-(C₁-C₈ alkylenyl);
 Substituted 8- to 10-membered heterobicycloalkyl-(C₁-C₈ alkylenyl);

- Phenyl-(C₁-C₈ alkylenyl);
 Substituted phenyl-(C₁-C₈ alkylenyl);
 Naphthyl-(C₁-C₈ alkylenyl);
 Substituted naphthyl-(C₁-C₈ alkylenyl);
 5 5- or 6-membered heteroaryl-(C₁-C₈ alkylenyl);
 Substituted 5- or 6-membered heteroaryl-(C₁-C₈ alkylenyl);
 8- to 10-membered heterobiaryl-(C₁-C₈ alkylenyl);
 Substituted 8- to 10-membered heterobiaryl-(C₁-C₈ alkylenyl);
 Phenyl;
 10 Substituted phenyl;
 Naphthyl;
 Substituted naphthyl;
 5- or 6-membered heteroaryl;
 Substituted 5- or 6-membered heteroaryl;
 15 8- to 10-membered heterobiaryl; and
 Substituted 8- to 10-membered heterobiaryl;
- R² is independently selected from:
- H;
 C₁-C₆ alkyl;
 20 Phenyl-(C₁-C₈ alkylenyl)_m;
 Substituted phenyl-(C₁-C₈ alkylenyl)_m;
 Naphthyl-(C₁-C₈ alkylenyl)_m;
 Substituted naphthyl-(C₁-C₈ alkylenyl)_m;
 5- or 6-membered heteroaryl-(C₁-C₈ alkylenyl)_m;
 25 Substituted 5- or 6-membered heteroaryl-(C₁-C₈ alkylenyl)_m;
 8- to 10-membered heterobiaryl-(C₁-C₈ alkylenyl)_m;
 Substituted 8- to 10-membered heterobiaryl-(C₁-C₈ alkylenyl)_m;
 Phenyl-O-(C₁-C₈ alkylenyl);
 Substituted phenyl-O-(C₁-C₈ alkylenyl);
 30 Phenyl-S-(C₁-C₈ alkylenyl);
 Substituted phenyl-S-(C₁-C₈ alkylenyl);
 Phenyl-S(O)-(C₁-C₈ alkylenyl);
 Substituted phenyl-S(O)-(C₁-C₈ alkylenyl);


Phenyl-S(O)₂-(C₁-C₈ alkylenyl); and

Substituted phenyl-S(O)₂-(C₁-C₈ alkylenyl);

Each substituted R¹ and R² group contains from 1 to 4 substituents, each independently on a carbon or nitrogen atom, independently selected from:

- 5 C₁-C₆ alkyl;
CN;
CF₃;
HO;
(C₁-C₆ alkyl)-O;
10 (C₁-C₆ alkyl)-S(O)₂;
H₂N;
(C₁-C₆ alkyl)-N(H);
(C₁-C₆ alkyl)₂-N;
(C₁-C₆ alkyl)-C(O)O-(C₁-C₈ alkylenyl)_m;
15 (C₁-C₆ alkyl)-C(O)O-(1- to 8-membered heteroalkylenyl)_m;
(C₁-C₆ alkyl)-C(O)N(H)-(C₁-C₈ alkylenyl)_m;
(C₁-C₆ alkyl)-C(O)N(H)-(1- to 8-membered heteroalkylenyl)_m;
H₂NS(O)₂-(C₁-C₈ alkylenyl);
(C₁-C₆ alkyl)-N(H)S(O)₂-(C₁-C₈ alkylenyl)_m;
20 (C₁-C₆ alkyl)₂-NS(O)₂-(C₁-C₈ alkylenyl)_m;
3- to 6-membered heterocycloalkyl-(G)_m;
Substituted 3- to 6-membered heterocycloalkyl-(G)_m;
5- or 6-membered heteroaryl-(G)_m;
Substituted 5- or 6-membered heteroaryl-(G)_m;
25 (C₁-C₆ alkyl)-S(O)₂-N(H)-C(O)-(C₁-C₈ alkylenyl)_m;
(C₁-C₆ alkyl)-C(O)-N(H)-S(O)₂-(C₁-C₈ alkylenyl)_m;
Halo; and
HO₂C;

- wherein 2 substituents may be taken together with a carbon atom to which they
30 are both bonded to form the group C=O;
wherein two adjacent, substantially sp² carbon atoms may be taken together with a
diradical substituent to form a cyclic diradical selected from:



5

R is H or C₁-C₆ alkyl;

G is CH₂; O, S, S(O); or S(O)₂;

m is an integer of 0 or 1;

10 wherein each C₈-C₁₀ bicycloalkyl is a bicyclic carbocyclic ring that contains 8-, 9-, or 10-member carbon atoms which are 5,5-fused, 6,5-fused, or 6,6-fused bicyclic rings, respectively, and wherein the ring is saturated or optionally contains one carbon-carbon double bond;

15 wherein each 8- to 10-membered heterobicycloalkyl is a bicyclic ring that contains carbon atoms and from 1 to 4 heteroatoms independently selected from 2 O, 1 S, 1 S(O), 1 S(O)₂, 1 N, 4 N(H), and 4 N(C₁-C₆ alkyl), and wherein when two O atoms or one O atom and one S atom are present, the two O atoms or one O atom and one S atom are not bonded to each other, and wherein the ring is saturated or optionally contains one carbon-carbon or carbon-nitrogen double

bond, and wherein the heterobicycloalkyl is a 5,5-fused, 6,5-fused, or 6,6-fused bicyclic ring, respectively,

wherein each heterocycloalkyl is a ring that contains carbon atoms and from 1 to 4 heteroatoms independently selected from 2 O, 1 S, 1 S(O), 1 S(O)₂, 1 N, 4 N(H), and 4 N(C₁-C₆ alkyl), and wherein when two O atoms or one O atom and one S atom are present, the two O atoms or one O atom and one S atom are not bonded to each other, and wherein the ring is saturated or optionally contains one carbon-carbon or carbon-nitrogen double bond;

wherein each 5-membered heteroaryl contains carbon atoms and from 1 to 4 heteroatoms independently selected from 1 O, 1 S, 1 N(H), 1 N(C₁-C₆ alkyl), and 4 N, and each 6-membered heteroaryl contains carbon atoms and 1 or 2 heteroatoms independently selected from N, N(H), and N(C₁-C₆ alkyl), and 5- and 6-membered heteroaryl are monocyclic rings;

wherein each heterobiaryl contains carbon atoms and from 1 to 4 heteroatoms independently selected from 1 O, 1 S, 1 N(H), 1 N(C₁-C₆ alkyl), and 4 N, and where the 8-, 9-, and 10-membered heterobiaryl are 5,5-fused, 6,5-fused, and 6,6-fused bicyclic rings, respectively, and wherein at least 1 of the 2 fused rings of a bicyclic ring is aromatic, and wherein when the O and S atoms both are present, the O and S atoms are not bonded to each other;

wherein with any (C₁-C₆ alkyl)₂-N group, the C₁-C₆ alkyl groups may be optionally taken together with the nitrogen atom to which they are attached to form a 5- or 6-membered heterocycloalkyl; and

wherein each group and each substituent recited above is independently selected.

36. The compound according to Embodiment 35, selected from:

3-Benzyl-6-(3-phenyl-prop-2-ynyl)-5,6,7,8-tetrahydro-3H-pyrido[4,3-d]pyrimidin-4-one;

6-(3-Phenyl-prop-2-ynyl)-3-pyridin-4-ylmethyl-5,6,7,8-tetrahydro-3H-pyrido[4,3-d]pyrimidin-4-one;

4-[4-Oxo-6-(3-phenyl-prop-2-ynyl)-5,6,7,8-tetrahydro-4H-pyrido[4,3-d]pyrimidin-3-ylmethyl]-benzoic acid; and

4-[4-Oxo-6-(3-phenyl-prop-2-ynyl)-5,6,7,8-tetrahydro-4H-pyrido[4,3-d]pyrimidin-3-ylmethyl]-benzonitrile; or
a pharmaceutically acceptable salt thereof.

- 5 37. The compound according to Embodiment 35, selected from:
6-(3-Phenyl-prop-2-ynyl)-3-[4-(1H-tetrazol-5-yl)-benzyl]-5,6,7,8-
 tetrahydro-3H-pyrido[4,3-d]pyrimidin-4-one;
3-Biphenyl-4-ylmethyl-6-(3-phenyl-prop-2-ynyl)-5,6,7,8-tetrahydro-3H-
 pyrido[4,3-d]pyrimidin-4-one;
10 6-(3-Phenyl-prop-2-ynyl)-3-(4-pyridin-4-yl-benzyl)-5,6,7,8-tetrahydro-
 3H-pyrido[4,3-d]pyrimidin-4-one;
3-(4-Furan-3-yl-benzyl)-6-(3-phenyl-prop-2-ynyl)-5,6,7,8-tetrahydro-3H-
 pyrido[4,3-d]pyrimidin-4-one;
15 3-(4-Furan-2-yl-benzyl)-6-(3-phenyl-prop-2-ynyl)-5,6,7,8-tetrahydro-3H-
 pyrido[4,3-d]pyrimidin-4-one;
6-(3-Phenyl-prop-2-ynyl)-3-(4-thiophen-3-yl-benzyl)-5,6,7,8-tetrahydro-
 3H-pyrido[4,3-d]pyrimidin-4-one;
6-(3-Phenyl-prop-2-ynyl)-3-(4-thiophen-2-yl-benzyl)-5,6,7,8-tetrahydro-
 3H-pyrido[4,3-d]pyrimidin-4-one;
20 6-(3-Phenyl-prop-2-ynyl)-3-(4-thiophen-3-yl-benzyl)-5,6,7,8-tetrahydro-
 3H-pyrido[4,3-d]pyrimidin-4-one;
6-(3-Phenyl-prop-2-ynyl)-3-[4-(1H-pyrrol-2-yl)-benzyl]-5,6,7,8-
 tetrahydro-3H-pyrido[4,3-d]pyrimidin-4-one;
3-[4-(1-Methyl-1H-pyrrol-3-yl)-benzyl]-6-(3-phenyl-prop-2-ynyl)-5,6,7,8-
25 tetrahydro-3H-pyrido[4,3-d]pyrimidin-4-one; and
3-[4-(1-Methyl-1H-pyrrol-2-yl)-benzyl]-6-(3-phenyl-prop-2-ynyl)-5,6,7,8-
 tetrahydro-3H-pyrido[4,3-d]pyrimidin-4-one; or
a pharmaceutically acceptable salt thereof.
- 30 38. The compound according to Embodiment 35, selected from:
3-(4-Morpholin-4-yl-benzyl)-6-(3-phenyl-prop-2-ynyl)-5,6,7,8-tetrahydro-
 3H-pyrido[4,3-d]pyrimidin-4-one;

- 3-(4-Dimethylamino-benzyl)-6-(3-phenyl-prop-2-ynyl)-5,6,7,8-tetrahydro-
3H-pyrido[4,3-d]pyrimidin-4-one;
- 3-(4-Aziridin-1-yl-benzyl)-6-(3-phenyl-prop-2-ynyl)-5,6,7,8-tetrahydro-
3H-pyrido[4,3-d]pyrimidin-4-one;
- 5 3-(4-Methyl-benzyl)-6-(3-phenyl-prop-2-ynyl)-5,6,7,8-tetrahydro-3H-
pyrido[4,3-d]pyrimidin-4-one;
- 6-(3-Phenyl-prop-2-ynyl)-3-(4-trifluoromethyl-benzyl)-5,6,7,8-tetrahydro-
3H-pyrido[4,3-d]pyrimidin-4-one;
- 6-(3-Phenyl-prop-2-ynyl)-3-(4-trichloromethyl-benzyl)-5,6,7,8-tetrahydro-
10 3H-pyrido[4,3-d]pyrimidin-4-one;
- 3-(4-Methylsulfanyl-benzyl)-6-(3-phenyl-prop-2-ynyl)-5,6,7,8-tetrahydro-
3H-pyrido[4,3-d]pyrimidin-4-one;
- 3-(4-Methoxy-benzyl)-6-(3-phenyl-prop-2-ynyl)-5,6,7,8-tetrahydro-3H-
pyrido[4,3-d]pyrimidin-4-one;
- 15 3-Pentafluorophenylmethyl-6-(3-phenyl-prop-2-ynyl)-5,6,7,8-tetrahydro-
3H-pyrido[4,3-d]pyrimidin-4-one;
- 6-(3-Phenyl-prop-2-ynyl)-3-(3,4,5-trifluoro-benzyl)-5,6,7,8-tetrahydro-
3H-pyrido[4,3-d]pyrimidin-4-one;
- 3-(4-Fluoro-benzyl)-6-(3-phenyl-prop-2-ynyl)-5,6,7,8-tetrahydro-3H-
20 pyrido[4,3-d]pyrimidin-4-one;
- 3-(4-Methanesulfonyl-benzyl)-6-(3-phenyl-prop-2-ynyl)-5,6,7,8-
tetrahydro-3H-pyrido[4,3-d]pyrimidin-4-one;
- 4-[4-Oxo-6-(3-phenyl-prop-2-ynyl)-5,6,7,8-tetrahydro-4H-pyrido[4,3-
d]pyrimidin-3-ylmethyl]-benzenesulfonamide;
- 25 4-[4-Oxo-6-(3-phenyl-prop-2-ynyl)-5,6,7,8-tetrahydro-4H-pyrido[4,3-
d]pyrimidin-3-ylmethyl]-benzenesulfonamide; and
- 4-[4-Oxo-6-(3-phenyl-prop-2-ynyl)-5,6,7,8-tetrahydro-4H-pyrido[4,3-
d]pyrimidin-3-ylmethyl]-benzenesulfonic acid; or
- a pharmaceutically acceptable salt thereof.
- 30
39. The compound according to Embodiment 35, selected from:
3-Benzyl-6-[3-(4-methoxy-phenyl)-prop-2-ynyl]-5,6,7,8-tetrahydro-3H-
pyrido[4,3-d]pyrimidin-4-one;

- 6-[3-(4-Methoxy-phenyl)-prop-2-ynyl]-3-pyridin-4-ylmethyl-5,6,7,8-tetrahydro-3H-pyrido[4,3-d]pyrimidin-4-one;
- 4-{6-[3-(4-Methoxy-phenyl)-prop-2-ynyl]-4-oxo-5,6,7,8-tetrahydro-4H-pyrido[4,3-d]pyrimidin-3-ylmethyl}-benzoic acid; and
- 5 4-{6-[3-(4-Methoxy-phenyl)-prop-2-ynyl]-4-oxo-5,6,7,8-tetrahydro-4H-pyrido[4,3-d]pyrimidin-3-ylmethyl}-benzonitrile; or
a pharmaceutically acceptable salt thereof.
40. The compound according to Embodiment 35, selected from:
- 10 6-[3-(4-Methoxy-phenyl)-prop-2-ynyl]-3-[4-(1H-tetrazol-5-yl)-benzyl]-5,6,7,8-tetrahydro-3H-pyrido[4,3-d]pyrimidin-4-one;
- 3-Biphenyl-4-ylmethyl-6-[3-(4-methoxy-phenyl)-prop-2-ynyl]-5,6,7,8-tetrahydro-3H-pyrido[4,3-d]pyrimidin-4-one;
- 15 6-(3-Phenyl-prop-2-ynyl)-3-(4-pyridin-4-yl-benzyl)-5,6,7,8-tetrahydro-3H-pyrido[4,3-d]pyrimidin-4-one;
- 3-(4-Furan-3-yl-benzyl)-6-[3-(4-methoxy-phenyl)-prop-2-ynyl]-5,6,7,8-tetrahydro-3H-pyrido[4,3-d]pyrimidin-4-one;
- 3-(4-Furan-2-yl-benzyl)-6-[3-(4-methoxy-phenyl)-prop-2-ynyl]-5,6,7,8-tetrahydro-3H-pyrido[4,3-d]pyrimidin-4-one;
- 20 6-[3-(4-Methoxy-phenyl)-prop-2-ynyl]-3-(4-thiophen-3-yl-benzyl)-5,6,7,8-tetrahydro-3H-pyrido[4,3-d]pyrimidin-4-one;
- 6-[3-(4-Methoxy-phenyl)-prop-2-ynyl]-3-(4-thiophen-2-yl-benzyl)-5,6,7,8-tetrahydro-3H-pyrido[4,3-d]pyrimidin-4-one;
- 25 6-[3-(4-Methoxy-phenyl)-prop-2-ynyl]-3-[4-(1H-pyrrol-3-yl)-benzyl]-5,6,7,8-tetrahydro-3H-pyrido[4,3-d]pyrimidin-4-one;
- 6-[3-(4-Methoxy-phenyl)-prop-2-ynyl]-3-[4-(1H-pyrrol-2-yl)-benzyl]-5,6,7,8-tetrahydro-3H-pyrido[4,3-d]pyrimidin-4-one;
- 6-[3-(4-Methoxy-phenyl)-prop-2-ynyl]-3-[4-(1-methyl-1H-pyrrol-3-yl)-benzyl]-5,6,7,8-tetrahydro-3H-pyrido[4,3-d]pyrimidin-4-one; and
- 30 6-[3-(4-Methoxy-phenyl)-prop-2-ynyl]-3-[4-(1-methyl-1H-pyrrol-2-yl)-benzyl]-5,6,7,8-tetrahydro-3H-pyrido[4,3-d]pyrimidin-4-one; or
a pharmaceutically acceptable salt thereof.

41. The compound according to Embodiment 35, selected from:
- 6-[3-(4-Methoxy-phenyl)-prop-2-ynyl]-3-(4-morpholin-4-yl-benzyl)-
5,6,7,8-tetrahydro-3H-pyrido[4,3-d]pyrimidin-4-one;
- 3-(4-Dimethylamino-benzyl)-6-[3-(4-methoxy-phenyl)-prop-2-ynyl]-
5,6,7,8-tetrahydro-3H-pyrido[4,3-d]pyrimidin-4-one;
- 3-(4-Aziridin-1-yl-benzyl)-6-[3-(4-methoxy-phenyl)-prop-2-ynyl]-5,6,7,8-
tetrahydro-3H-pyrido[4,3-d]pyrimidin-4-one;
- 6-[3-(4-Methoxy-phenyl)-prop-2-ynyl]-3-(4-methyl-benzyl)-5,6,7,8-
tetrahydro-3H-pyrido[4,3-d]pyrimidin-4-one;
- 6-[3-(4-Methoxy-phenyl)-prop-2-ynyl]-3-(4-trifluoromethyl-benzyl)-
5,6,7,8-tetrahydro-3H-pyrido[4,3-d]pyrimidin-4-one;
- 6-[3-(4-Methoxy-phenyl)-prop-2-ynyl]-3-(4-trichloromethyl-benzyl)-
5,6,7,8-tetrahydro-3H-pyrido[4,3-d]pyrimidin-4-one;
- 6-[3-(4-Methoxy-phenyl)-prop-2-ynyl]-3-(4-methylsulfanyl-benzyl)-
5,6,7,8-tetrahydro-3H-pyrido[4,3-d]pyrimidin-4-one;
- 3-(4-Methoxy-benzyl)-6-[3-(4-methoxy-phenyl)-prop-2-ynyl]-5,6,7,8-
tetrahydro-3H-pyrido[4,3-d]pyrimidin-4-one;
- 6-[3-(4-Methoxy-phenyl)-prop-2-ynyl]-3-pentafluorophenylmethyl-
5,6,7,8-tetrahydro-3H-pyrido[4,3-d]pyrimidin-4-one;
- 6-[3-(4-Methoxy-phenyl)-prop-2-ynyl]-3-(3,4,5-trifluoro-benzyl)-5,6,7,8-
tetrahydro-3H-pyrido[4,3-d]pyrimidin-4-one;
- 3-(4-Fluoro-benzyl)-6-[3-(4-methoxy-phenyl)-prop-2-ynyl]-5,6,7,8-
tetrahydro-3H-pyrido[4,3-d]pyrimidin-4-one;
- 3-(4-Methanesulfonyl-benzyl)-6-[3-(4-methoxy-phenyl)-prop-2-ynyl]-
5,6,7,8-tetrahydro-3H-pyrido[4,3-d]pyrimidin-4-one;
- 4-{6-[3-(4-Methoxy-phenyl)-prop-2-ynyl]-4-oxo-5,6,7,8-tetrahydro-4H-
pyrido[4,3-d]pyrimidin-3-ylmethyl}-benzenesulfonamide;
- 4-{6-[3-(4-Methoxy-phenyl)-prop-2-ynyl]-4-oxo-5,6,7,8-tetrahydro-4H-
pyrido[4,3-d]pyrimidin-3-ylmethyl}-N,N-dimethyl-
benzenesulfonamide; and
- 4-{6-[3-(4-Methoxy-phenyl)-prop-2-ynyl]-4-oxo-5,6,7,8-tetrahydro-4H-
pyrido[4,3-d]pyrimidin-3-ylmethyl}-benzenesulfonic acid; or


a pharmaceutically acceptable salt thereof.

42. The compound according to Embodiment 35, selected from:
3-Benzyl-6-(3-pyridin-4-yl-prop-2-ynyl)-5,6,7,8-tetrahydro-3H-
5 pyrido[4,3-d]pyrimidin-4-one; and
3-Benzyl-6-[3-(2-methoxy-pyridin-4-yl)-prop-2-ynyl]-5,6,7,8-tetrahydro-
3H-pyrido[4,3-d]pyrimidin-4-one; or
a pharmaceutically acceptable salt thereof.
- 10 43. The compound according to Embodiment 35, selected from:
3-Benzyl-6-[3-(3-methoxy-phenyl)-prop-2-ynyl]-5,6,7,8-tetrahydro-3H-
pyrido[4,3-d]pyrimidin-4-one;
3-Benzyl-6-[3-(4-dimethylamino-phenyl)-prop-2-ynyl]-5,6,7,8-tetrahydro-
3H-pyrido[4,3-d]pyrimidin-4-one;
15 3-Benzyl-6-[3-(4-methylsulfanyl-phenyl)-prop-2-ynyl]-5,6,7,8-tetrahydro-
3H-pyrido[4,3-d]pyrimidin-4-one;
3-Benzyl-6-[3-(4-fluoro-phenyl)-prop-2-ynyl]-5,6,7,8-tetrahydro-3H-
pyrido[4,3-d]pyrimidin-4-one;
3-Benzyl-6-[3-(4-chloro-phenyl)-prop-2-ynyl]-5,6,7,8-tetrahydro-3H-
20 pyrido[4,3-d]pyrimidin-4-one;
3-Benzyl-6-[3-(4-bromo-phenyl)-prop-2-ynyl]-5,6,7,8-tetrahydro-3H-
pyrido[4,3-d]pyrimidin-4-one; and
3-Benzyl-6-[3-(4-iodo-phenyl)-prop-2-ynyl]-5,6,7,8-tetrahydro-3H-
pyrido[4,3-d]pyrimidin-4-one; or
25 a pharmaceutically acceptable salt thereof.
44. The compound according to Embodiment 35, selected from:
3-Benzyl-6-(3-furan-2-yl-prop-2-ynyl)-5,6,7,8-tetrahydro-3H-pyrido[4,3-
d]pyrimidin-4-one;
30 3-Benzyl-6-(3-thiophen-2-yl-prop-2-ynyl)-5,6,7,8-tetrahydro-3H-
pyrido[4,3-d]pyrimidin-4-one;

- 3-Benzyl-6-[3-(1H-pyrrol-2-yl)-prop-2-ynyl]-5,6,7,8-tetrahydro-3H-pyrido[4,3-d]pyrimidin-4-one;
3-Benzyl-6-[3-(1-methyl-1H-pyrrol-2-yl)-prop-2-ynyl]-5,6,7,8-tetrahydro-3H-pyrido[4,3-d]pyrimidin-4-one;
5 3-Benzyl-6-[3-(1-methyl-1H-imidazol-2-yl)-prop-2-ynyl]-5,6,7,8-tetrahydro-3H-pyrido[4,3-d]pyrimidin-4-one;
3-Benzyl-6-[3-(3-methyl-3H-imidazol-4-yl)-prop-2-ynyl]-5,6,7,8-tetrahydro-3H-pyrido[4,3-d]pyrimidin-4-one;
3-Benzyl-6-[3-(4-methyl-4H-[1,2,4]triazol-3-yl)-prop-2-ynyl]-5,6,7,8-tetrahydro-3H-pyrido[4,3-d]pyrimidin-4-one;
10 3-Benzyl-6-[3-(1-methyl-1H-pyrrol-3-yl)-prop-2-ynyl]-5,6,7,8-tetrahydro-3H-pyrido[4,3-d]pyrimidin-4-one;
3-Benzyl-6-(3-oxazol-2-yl-prop-2-ynyl)-5,6,7,8-tetrahydro-3H-pyrido[4,3-d]pyrimidin-4-one;
15 3-Benzyl-6-(3-oxazol-5-yl-prop-2-ynyl)-5,6,7,8-tetrahydro-3H-pyrido[4,3-d]pyrimidin-4-one; and
3-Benzyl-6-(3-isoxazol-5-yl-prop-2-ynyl)-5,6,7,8-tetrahydro-3H-pyrido[4,3-d]pyrimidin-4-one; or
a pharmaceutically acceptable salt thereof.

20

45. A compound of Formula III



III

or a pharmaceutically acceptable salt thereof, or a pyrido-N-oxide thereof,
wherein:

25

R¹ is independently selected from:

- C₅ or C₆ cycloalkyl-(C₁-C₈ alkylenyl);
- Substituted C₅ or C₆ cycloalkyl-(C₁-C₈ alkylenyl);
- C₈-C₁₀ bicycloalkyl-(C₁-C₈ alkylenyl);
- Substituted C₈-C₁₀ bicycloalkyl-(C₁-C₈ alkylenyl);

- 5- or 6-membered heterocycloalkyl-(C₁-C₈ alkylenyl);
Substituted 5- or 6-membered heterocycloalkyl-(C₁-C₈ alkylenyl);
8- to 10-membered heterobicycloalkyl-(C₁-C₈ alkylenyl);
Substituted 8- to 10-membered heterobicycloalkyl-(C₁-C₈ alkylenyl);
5 Phenyl-(C₁-C₈ alkylenyl);
Substituted phenyl-(C₁-C₈ alkylenyl);
Naphthyl-(C₁-C₈ alkylenyl);
Substituted naphthyl-(C₁-C₈ alkylenyl);
5- or 6-membered heteroaryl-(C₁-C₈ alkylenyl);
10 Substituted 5- or 6-membered heteroaryl-(C₁-C₈ alkylenyl);
8- to 10-membered heterobiaryl-(C₁-C₈ alkylenyl);
Substituted 8- to 10-membered heterobiaryl-(C₁-C₈ alkylenyl);
Phenyl;
Substituted phenyl;
15 Naphthyl;
Substituted naphthyl;
5- or 6-membered heteroaryl;
Substituted 5- or 6-membered heteroaryl;
8- to 10-membered heterobiaryl; and
20 Substituted 8- to 10-membered heterobiaryl;
- R² is independently selected from:
- H;
C₁-C₆ alkyl;
Phenyl-(C₁-C₈ alkylenyl)_m;
25 Substituted phenyl-(C₁-C₈ alkylenyl)_m;
Naphthyl-(C₁-C₈ alkylenyl)_m;
Substituted naphthyl-(C₁-C₈ alkylenyl)_m;
5- or 6-membered heteroaryl-(C₁-C₈ alkylenyl)_m;
Substituted 5- or 6-membered heteroaryl-(C₁-C₈ alkylenyl)_m;
30 8- to 10-membered heterobiaryl-(C₁-C₈ alkylenyl)_m;
Substituted 8- to 10-membered heterobiaryl-(C₁-C₈ alkylenyl)_m;
Phenyl-O-(C₁-C₈ alkylenyl);
Substituted phenyl-O-(C₁-C₈ alkylenyl);


Phenyl-S-(C₁-C₈ alkylenyl);
Substituted phenyl-S-(C₁-C₈ alkylenyl);
Phenyl-S(O)-(C₁-C₈ alkylenyl);
Substituted phenyl-S(O)-(C₁-C₈ alkylenyl);
5 Phenyl-S(O)₂-(C₁-C₈ alkylenyl); and
Substituted phenyl-S(O)₂-(C₁-C₈ alkylenyl)

Each substituted R¹ and R² group contains from 1 to 4 substituents, each independently on a carbon or nitrogen atom, independently selected from:

- 10 C₁-C₆ alkyl;
CN;
CF₃;
HO;
(C₁-C₆ alkyl)-O;
(C₁-C₆ alkyl)-S(O)₂;
15 H₂N;
(C₁-C₆ alkyl)-N(H);
(C₁-C₆ alkyl)₂-N;
(C₁-C₆ alkyl)-C(O)O-(C₁-C₈ alkylenyl)_m;
(C₁-C₆ alkyl)-C(O)O-(1- to 8-membered heteroalkylenyl)_m;
20 (C₁-C₆ alkyl)-C(O)N(H)-(C₁-C₈ alkylenyl)_m;
(C₁-C₆ alkyl)-C(O)N(H)-(1- to 8-membered heteroalkylenyl)_m;
H₂NS(O)₂-(C₁-C₈ alkylenyl);
(C₁-C₆ alkyl)-N(H)S(O)₂-(C₁-C₈ alkylenyl)_m;
(C₁-C₆ alkyl)₂-NS(O)₂-(C₁-C₈ alkylenyl)_m;
25 3- to 6-membered heterocycloalkyl-(G)_m;
Substituted 3- to 6-membered heterocycloalkyl-(G)_m;
5- or 6-membered heteroaryl-(G)_m;
Substituted 5- or 6-membered heteroaryl-(G)_m;
(C₁-C₆ alkyl)-S(O)₂-N(H)-C(O)-(C₁-C₈ alkylenyl)_m;
30 (C₁-C₆ alkyl)-C(O)-N(H)-S(O)₂-(C₁-C₈ alkylenyl)_m;
Halo; and
HO₂C;

wherein 2 substituents may be taken together with a carbon atom to which they are both bonded to form the group C=O;

wherein two adjacent, substantially sp^2 carbon atoms may be taken together with a diradical substituent to form a cyclic diradical selected from:



R is H or C₁-C₆ alkyl;

G is CH₂; O, S, S(O); or S(O)₂;

m is an integer of 0 or 1;

wherein each C₈-C₁₀ bicycloalkyl is a bicyclic carbocyclic ring that contains 8-, 9-, or 10-member carbon atoms which are 5,5-fused, 6,5-fused, or 6,6-fused bicyclic rings, respectively, and wherein the ring is saturated or optionally contains one carbon-carbon double bond;

wherein each 8- to 10-membered heterobicycloalkyl is a bicyclic ring that contains carbon atoms and from 1 to 4 heteroatoms independently selected from 2 O, 1 S, 1 S(O), 1 S(O)₂, 1 N, 4 N(H), and 4 N(C₁-C₆ alkyl), and wherein when two

O atoms or one O atom and one S atom are present, the two O atoms or one O atom and one S atom are not bonded to each other, and wherein the ring is saturated or optionally contains one carbon-carbon or carbon-nitrogen double bond, and wherein the heterobicycloalkyl is a 5,5-fused, 6,5-fused, or 6,6-fused bicyclic ring, respectively,

wherein each heterocycloalkyl is a ring that contains carbon atoms and from 1 to 4 heteroatoms independently selected from 2 O, 1 S, 1 S(O), 1 S(O)₂, 1 N, 4 N(H), and 4 N(C₁-C₆ alkyl), and wherein when two O atoms or one O atom and one S atom are present, the two O atoms or one O atom and one S atom are not bonded to each other, and wherein the ring is saturated or optionally contains one carbon-carbon or carbon-nitrogen double bond;

wherein each 5-membered heteroaryl contains carbon atoms and from 1 to 4 heteroatoms independently selected from 1 O, 1 S, 1 N(H), 1 N(C₁-C₆ alkyl), and 4 N, and each 6-membered heteroaryl contains carbon atoms and 1 or 2 heteroatoms independently selected from N, N(H), and N(C₁-C₆ alkyl), and 5- and 6-membered heteroaryl are monocyclic rings;

wherein each heterobiaryl contains carbon atoms and from 1 to 4 heteroatoms independently selected from 1 O, 1 S, 1 N(H), 1 N(C₁-C₆ alkyl), and 4 N, and where the 8-, 9-, and 10-membered heterobiaryl are 5,5-fused, 6,5-fused, and 6,6-fused bicyclic rings, respectively, and wherein at least 1 of the 2 fused rings of a bicyclic ring is aromatic, and wherein when the O and S atoms both are present, the O and S atoms are not bonded to each other;

wherein with any (C₁-C₆ alkyl)₂-N group, the C₁-C₆ alkyl groups may be optionally taken together with the nitrogen atom to which they are attached to form a 5- or 6-membered heterocycloalkyl; and

wherein each group and each substituent recited above is independently selected.

46. The compound according to Embodiment 45, selected from:
- 3-Benzyl-4-oxo-3,5,7,8-tetrahydro-4H-pyrido[4,3-d]pyrimidine-6-carboxylic acid benzyl ester; and
- 3-Benzyl-4-oxo-3,5,7,8-tetrahydro-4H-pyrido[4,3-d]pyrimidine-6-carboxylic acid pyridin-4-ylmethyl ester; or

a pharmaceutically acceptable salt thereof.

47. The compound according to Embodiment 45, selected from:
- 5 3-Benzyl-4-oxo-3,5,7,8-tetrahydro-4H-pyrido[4,3-d]pyrimidine-6-carboxylic acid 2-methoxy-pyridin-4-ylmethyl ester;
- 3-Benzyl-4-oxo-3,5,7,8-tetrahydro-4H-pyrido[4,3-d]pyrimidine-6-carboxylic acid 3-methoxy-benzyl ester;
- 3-Benzyl-4-oxo-3,5,7,8-tetrahydro-4H-pyrido[4,3-d]pyrimidine-6-carboxylic acid 4-methoxy-benzyl ester;
- 10 3-Benzyl-4-oxo-3,5,7,8-tetrahydro-4H-pyrido[4,3-d]pyrimidine-6-carboxylic acid 4-fluoro-benzyl ester;
- 3-Benzyl-4-oxo-3,5,7,8-tetrahydro-4H-pyrido[4,3-d]pyrimidine-6-carboxylic acid 4-chloro-benzyl ester;
- 3-Benzyl-4-oxo-3,5,7,8-tetrahydro-4H-pyrido[4,3-d]pyrimidine-6-carboxylic acid 4-bromo-benzyl ester;
- 15 3-Benzyl-4-oxo-3,5,7,8-tetrahydro-4H-pyrido[4,3-d]pyrimidine-6-carboxylic acid 4-iodo-benzyl ester;
- 3-Benzyl-4-oxo-3,5,7,8-tetrahydro-4H-pyrido[4,3-d]pyrimidine-6-carboxylic acid 4-dimethylamino-benzyl ester; and
- 20 3-Benzyl-4-oxo-3,5,7,8-tetrahydro-4H-pyrido[4,3-d]pyrimidine-6-carboxylic acid 4-methylsulfanyl-benzyl ester; or
- a pharmaceutically acceptable salt thereof.
48. The compound according to Embodiment 45, selected from:
- 25 3-Benzyl-4-oxo-3,5,7,8-tetrahydro-4H-pyrido[4,3-d]pyrimidine-6-carboxylic acid furan-2-ylmethyl ester;
- 3-Benzyl-4-oxo-3,5,7,8-tetrahydro-4H-pyrido[4,3-d]pyrimidine-6-carboxylic acid thiophen-2-ylmethyl ester;
- 3-Benzyl-4-oxo-3,5,7,8-tetrahydro-4H-pyrido[4,3-d]pyrimidine-6-carboxylic acid 1H-pyrrol-2-ylmethyl ester;
- 30 3-Benzyl-4-oxo-3,5,7,8-tetrahydro-4H-pyrido[4,3-d]pyrimidine-6-carboxylic acid 1-methyl-1H-pyrrol-2-ylmethyl ester;


3-Benzyl-4-oxo-3,5,7,8-tetrahydro-4H-pyrido[4,3-d]pyrimidine-6-carboxylic acid 1-methyl-1H-imidazol-2-ylmethyl ester;
3-Benzyl-4-oxo-3,5,7,8-tetrahydro-4H-pyrido[4,3-d]pyrimidine-6-carboxylic acid 3-methyl-3H-imidazol-4-ylmethyl ester;
5 3-Benzyl-4-oxo-3,5,7,8-tetrahydro-4H-pyrido[4,3-d]pyrimidine-6-carboxylic acid 4-methyl-4H-[1,2,4]triazol-3-ylmethyl ester;
3-Benzyl-4-oxo-3,5,7,8-tetrahydro-4H-pyrido[4,3-d]pyrimidine-6-carboxylic acid oxazol-2-ylmethyl ester;
3-Benzyl-4-oxo-3,5,7,8-tetrahydro-4H-pyrido[4,3-d]pyrimidine-6-carboxylic acid oxazol-5-ylmethyl ester;
10 3-Benzyl-4-oxo-3,5,7,8-tetrahydro-4H-pyrido[4,3-d]pyrimidine-6-carboxylic acid isoxazol-5-ylmethyl ester; and
3-Benzyl-4-oxo-3,5,7,8-tetrahydro-4H-pyrido[4,3-d]pyrimidine-6-carboxylic acid 1-methyl-1H-pyrrol-3-ylmethyl ester; or
15 a pharmaceutically acceptable salt thereof.

49. The compound according to Embodiment 45, selected from:
3-Benzyl-4-oxo-3,5,7,8-tetrahydro-4H-pyrido[4,3-d]pyrimidine-6-carboxylic acid pyrrol-1-ylmethyl ester;
20 3-Benzyl-4-oxo-3,5,7,8-tetrahydro-4H-pyrido[4,3-d]pyrimidine-6-carboxylic acid 3-methyl-pyrrol-1-ylmethyl ester; and
3-Benzyl-4-oxo-3,5,7,8-tetrahydro-4H-pyrido[4,3-d]pyrimidine-6-carboxylic acid 3,4-dimethyl-pyrrol-1-ylmethyl ester; or
a pharmaceutically acceptable salt thereof.

25 50. The compound according to Embodiment 45, selected from:
3-Benzyl-4-oxo-3,5,7,8-tetrahydro-4H-pyrido[4,3-d]pyrimidine-6-carboxylic acid morpholin-4-ylmethyl ester; and
3-Benzyl-4-oxo-3,5,7,8-tetrahydro-4H-pyrido[4,3-d]pyrimidine-6-carboxylic acid pyrrolidin-1-ylmethyl ester; or
30 a pharmaceutically acceptable salt thereof.

51. The compound according to Embodiment 45, selected from:
3-Benzyl-4-oxo-3,5,7,8-tetrahydro-4H-pyrido[4,3-d]pyrimidine-6-carboxylic acid 1,3-dihydro-isoindol-2-ylmethyl ester;
3-Benzyl-4-oxo-3,5,7,8-tetrahydro-4H-pyrido[4,3-d]pyrimidine-6-carboxylic acid 1H-indol-2-ylmethyl ester;
3-Benzyl-4-oxo-3,5,7,8-tetrahydro-4H-pyrido[4,3-d]pyrimidine-6-carboxylic acid 1-methyl-1H-indol-2-ylmethyl ester;
3-Benzyl-4-oxo-3,5,7,8-tetrahydro-4H-pyrido[4,3-d]pyrimidine-6-carboxylic acid benzo[b]thiophen-2-ylmethyl ester; and
3-Benzyl-4-oxo-3,5,7,8-tetrahydro-4H-pyrido[4,3-d]pyrimidine-6-carboxylic acid benzofuran-2-ylmethyl ester; or
a pharmaceutically acceptable salt thereof.

52. A compound of Formula IV



IV

or a pharmaceutically acceptable salt thereof, or a pyrido-N-oxide thereof, wherein:


R¹ is independently selected from:

- C₅ or C₆ cycloalkyl-(C₁-C₈ alkylenyl);
- Substituted C₅ or C₆ cycloalkyl-(C₁-C₈ alkylenyl);
- C₈-C₁₀ bicycloalkyl-(C₁-C₈ alkylenyl);
- Substituted C₈-C₁₀ bicycloalkyl-(C₁-C₈ alkylenyl);
- 5- or 6-membered heterocycloalkyl-(C₁-C₈ alkylenyl);
- Substituted 5- or 6-membered heterocycloalkyl-(C₁-C₈ alkylenyl);
- 8- to 10-membered heterobicycloalkyl-(C₁-C₈ alkylenyl);
- Substituted 8- to 10-membered heterobicycloalkyl-(C₁-C₈ alkylenyl);
- Phenyl-(C₁-C₈ alkylenyl);
- Substituted phenyl-(C₁-C₈ alkylenyl);

- Naphthyl-(C₁-C₈ alkylenyl);
Substituted naphthyl-(C₁-C₈ alkylenyl);
5- or 6-membered heteroaryl-(C₁-C₈ alkylenyl);
Substituted 5- or 6-membered heteroaryl-(C₁-C₈ alkylenyl);
5 8- to 10-membered heterobiaryl-(C₁-C₈ alkylenyl);
Substituted 8- to 10-membered heterobiaryl-(C₁-C₈ alkylenyl);
Phenyl;
Substituted phenyl;
Naphthyl;
10 Substituted naphthyl;
5- or 6-membered heteroaryl;
Substituted 5- or 6-membered heteroaryl;
8- to 10-membered heterobiaryl; and
Substituted 8- to 10-membered heterobiaryl;
15 R² is independently selected from:
H;
C₁-C₆ alkyl;
Phenyl-(C₁-C₈ alkylenyl);
Substituted phenyl-(C₁-C₈ alkylenyl);
20 Naphthyl-(C₁-C₈ alkylenyl);
Substituted naphthyl-(C₁-C₈ alkylenyl);
5- or 6-membered heteroaryl-(C₁-C₈ alkylenyl);
Substituted 5- or 6-membered heteroaryl-(C₁-C₈ alkylenyl);
8- to 10-membered heterobiaryl-(C₁-C₈ alkylenyl);
25 Substituted 8- to 10-membered heterobiaryl-(C₁-C₈ alkylenyl);
Phenyl-O-(C₁-C₈ alkylenyl);
Substituted phenyl-O-(C₁-C₈ alkylenyl);
Phenyl-S-(C₁-C₈ alkylenyl);
Substituted phenyl-S-(C₁-C₈ alkylenyl);
30 Phenyl-S(O)-(C₁-C₈ alkylenyl);
Substituted phenyl-S(O)-(C₁-C₈ alkylenyl);
Phenyl-S(O)₂-(C₁-C₈ alkylenyl); and
Substituted phenyl-S(O)₂-(C₁-C₈ alkylenyl);

Each substituted R^1 and R^2 group contains from 1 to 4 substituents, each independently on a carbon or nitrogen atom, independently selected from:

- C₁-C₆ alkyl;
- CN;
- 5 CF₃;
- HO;
- (C₁-C₆ alkyl)-O;
- (C₁-C₆ alkyl)-S(O)₂;
- H₂N;
- 10 (C₁-C₆ alkyl)-N(H);
- (C₁-C₆ alkyl)₂-N;
- (C₁-C₆ alkyl)-C(O)O-(C₁-C₈ alkylenyl)_m;
- (C₁-C₆ alkyl)-C(O)O-(1- to 8-membered heteroalkylenyl)_m;
- (C₁-C₆ alkyl)-C(O)N(H)-(C₁-C₈ alkylenyl)_m;
- 15 (C₁-C₆ alkyl)-C(O)N(H)-(1- to 8-membered heteroalkylenyl)_m;
- H₂NS(O)₂-(C₁-C₈ alkylenyl);
- (C₁-C₆ alkyl)-N(H)S(O)₂-(C₁-C₈ alkylenyl)_m;
- (C₁-C₆ alkyl)₂-NS(O)₂-(C₁-C₈ alkylenyl)_m;
- 3- to 6-membered heterocycloalkyl-(G)_m;
- 20 Substituted 3- to 6-membered heterocycloalkyl-(G)_m;
- 5- or 6-membered heteroaryl-(G)_m;
- Substituted 5- or 6-membered heteroaryl-(G)_m;
- (C₁-C₆ alkyl)-S(O)₂-N(H)-C(O)-(C₁-C₈ alkylenyl)_m; and
- (C₁-C₆ alkyl)-C(O)-N(H)-S(O)₂-(C₁-C₈ alkylenyl)_m;
- 25 wherein each substituent on a carbon atom may further be independently selected from:
- Halo; and
- HO₂C;
- wherein 2 substituents may be taken together with a carbon atom to which they
- 30 are both bonded to form the group C=O;
- wherein two adjacent, substantially sp² carbon atoms may be taken together with a diradical substituent to form a cyclic diradical selected from:



5

R is H or C₁-C₆ alkyl;

G is CH₂; O, S, S(O); or S(O)₂;

m is an integer of 0 or 1;

R⁴ is independently selected from the groups:

10

H;

CH₃;

CH₃O;

HO;

CF₃; and

15

CN;

wherein each C₈-C₁₀ bicycloalkyl is a bicyclic carbocyclic ring that contains 8-, 9-, or 10-member carbon atoms which are 5,5-fused, 6,5-fused, or 6,6-fused bicyclic rings, respectively, and wherein the ring is saturated or optionally contains one carbon-carbon double bond;

wherein each 8- to 10-membered heterobicycloalkyl is a bicyclic ring that contains carbon atoms and from 1 to 4 heteroatoms independently selected from 2 O, 1 S, 1 S(O), 1 S(O)₂, 1 N, 4 N(H), and 4 N(C₁-C₆ alkyl), and wherein when two O atoms or one O atom and one S atom are present, the two O atoms or one O atom and one S atom are not bonded to each other, and wherein the ring is saturated or optionally contains one carbon-carbon or carbon-nitrogen double bond, and wherein the heterobicycloalkyl is a 5,5-fused, 6,5-fused, or 6,6-fused bicyclic ring, respectively,

wherein each heterocycloalkyl is a ring that contains carbon atoms and from 1 to 4 heteroatoms independently selected from 2 O, 1 S, 1 S(O), 1 S(O)₂, 1 N, 4 N(H), and 4 N(C₁-C₆ alkyl), and wherein when two O atoms or one O atom and one S atom are present, the two O atoms or one O atom and one S atom are not bonded to each other, and wherein the ring is saturated or optionally contains one carbon-carbon or carbon-nitrogen double bond;

wherein each 5-membered heteroaryl contains carbon atoms and from 1 to 4 heteroatoms independently selected from 1 O, 1 S, 1 N(H), 1 N(C₁-C₆ alkyl), and 4 N, and each 6-membered heteroaryl contains carbon atoms and 1 or 2 heteroatoms independently selected from N, N(H), and N(C₁-C₆ alkyl), and 5- and 6-membered heteroaryl are monocyclic rings;

wherein each heterobiaryl contains carbon atoms and from 1 to 4 heteroatoms independently selected from 1 O, 1 S, 1 N(H), 1 N(C₁-C₆ alkyl), and 4 N, and where the 8-, 9-, and 10-membered heterobiaryl are 5,5-fused, 6,5-fused, and 6,6-fused bicyclic rings, respectively, and wherein at least 1 of the 2 fused rings of a bicyclic ring is aromatic, and wherein when the O and S atoms both are present, the O and S atoms are not bonded to each other;


wherein with any (C₁-C₆ alkyl)₂-N group, the C₁-C₆ alkyl groups may be optionally taken together with the nitrogen atom to which they are attached to form a 5- or 6-membered heterocycloalkyl; and

wherein each group and each substituent recited above is independently selected.

53. The compound according to Embodiment 52, selected from:

- 3-Benzyl-1-methyl-2,4-dioxo-1,3,4,5,7,8-hexahydro-2H-pyrido[4,3-d]pyrimidine-6-carboxylic acid benzyl ester; and
3-Benzyl-1-methyl-2,4-dioxo-1,3,4,5,7,8-hexahydro-2H-pyrido[4,3-d]pyrimidine-6-carboxylic acid pyridin-4-ylmethyl ester; or
5 a pharmaceutically acceptable salt thereof.
54. The compound according to Embodiment 52, selected from:
3-Benzyl-1-methyl-2,4-dioxo-1,3,4,5,7,8-hexahydro-2H-pyrido[4,3-d]pyrimidine-6-carboxylic acid 4-methoxy-benzyl ester;
10 3-Benzyl-1-methyl-2,4-dioxo-1,3,4,5,7,8-hexahydro-2H-pyrido[4,3-d]pyrimidine-6-carboxylic acid 3-methoxy-benzyl ester; and
3-Benzyl-1-methyl-2,4-dioxo-1,3,4,5,7,8-hexahydro-2H-pyrido[4,3-d]pyrimidine-6-carboxylic acid 2-methoxy-pyridin-4-ylmethyl ester; or
15 a pharmaceutically acceptable salt thereof.
55. The compound according to Embodiment 52, selected from:
3-Benzyl-1-methyl-2,4-dioxo-1,3,4,5,7,8-hexahydro-2H-pyrido[4,3-d]pyrimidine-6-carboxylic acid pyridin-4-ylmethyl ester;
20 3-Benzyl-1-methyl-2,4-dioxo-1,3,4,5,7,8-hexahydro-2H-pyrido[4,3-d]pyrimidine-6-carboxylic acid 4-chloro-benzyl ester;
3-Benzyl-1-methyl-2,4-dioxo-1,3,4,5,7,8-hexahydro-2H-pyrido[4,3-d]pyrimidine-6-carboxylic acid 4-bromo-benzyl ester;
3-Benzyl-1-methyl-2,4-dioxo-1,3,4,5,7,8-hexahydro-2H-pyrido[4,3-d]pyrimidine-6-carboxylic acid 4-iodo-benzyl ester;
25 3-Benzyl-1-methyl-2,4-dioxo-1,3,4,5,7,8-hexahydro-2H-pyrido[4,3-d]pyrimidine-6-carboxylic acid 4-methyl-benzyl ester;
3-Benzyl-1-methyl-2,4-dioxo-1,3,4,5,7,8-hexahydro-2H-pyrido[4,3-d]pyrimidine-6-carboxylic acid 4-trifluoromethyl-benzyl ester; and
30 3-Benzyl-1-methyl-2,4-dioxo-1,3,4,5,7,8-hexahydro-2H-pyrido[4,3-d]pyrimidine-6-carboxylic acid 4-trichloromethyl-benzyl ester; or
a pharmaceutically acceptable salt thereof.

56. The compound according to Embodiment 52, selected from:
- 3-Benzyl-1-methyl-2,4-dioxo-1,3,4,5,7,8-hexahydro-2H-pyrido[4,3-
d]pyrimidine-6-carboxylic acid benzofuran-2-ylmethyl ester;
- 5 3-Benzyl-1-methyl-2,4-dioxo-1,3,4,5,7,8-hexahydro-2H-pyrido[4,3-
d]pyrimidine-6-carboxylic acid benzo[b]thiophen-2-ylmethyl ester;
- 3-Benzyl-1-methyl-2,4-dioxo-1,3,4,5,7,8-hexahydro-2H-pyrido[4,3-
d]pyrimidine-6-carboxylic acid 1-methyl-1H-indol-2-ylmethyl
ester;
- 10 3-Benzyl-1-methyl-2,4-dioxo-1,3,4,5,7,8-hexahydro-2H-pyrido[4,3-
d]pyrimidine-6-carboxylic acid 1H-indol-2-ylmethyl ester; and
- 3-Benzyl-1-methyl-2,4-dioxo-1,3,4,5,7,8-hexahydro-2H-pyrido[4,3-
d]pyrimidine-6-carboxylic acid 1,3-dihydro-isindol-2-ylmethyl
ester; or
- 15 a pharmaceutically acceptable salt thereof.
57. The compound according to Embodiment 52, selected from:
- 3-Benzyl-1-methyl-2,4-dioxo-1,3,4,5,7,8-hexahydro-2H-pyrido[4,3-
d]pyrimidine-6-carboxylic acid pyrrolidin-1-ylmethyl ester;
- 20 3-Benzyl-1-methyl-2,4-dioxo-1,3,4,5,7,8-hexahydro-2H-pyrido[4,3-
d]pyrimidine-6-carboxylic acid pyrrol-1-ylmethyl ester;
- 3-Benzyl-1-methyl-2,4-dioxo-1,3,4,5,7,8-hexahydro-2H-pyrido[4,3-
d]pyrimidine-6-carboxylic acid piperidin-1-ylmethyl ester; and
- 25 3-Benzyl-1-methyl-2,4-dioxo-1,3,4,5,7,8-hexahydro-2H-pyrido[4,3-
d]pyrimidine-6-carboxylic acid 4-methyl-piperazin-1-ylmethyl
ester; or
- a pharmaceutically acceptable salt thereof.
58. A compound of Formula V



V

or a pharmaceutically acceptable salt thereof, or a pyrido-N-oxide thereof,
wherein:

R¹ is independently selected from:

- 5 C₅ or C₆ cycloalkyl-(C₁-C₈ alkylenyl);
 Substituted C₅ or C₆ cycloalkyl-(C₁-C₈ alkylenyl);
 C₈-C₁₀ bicycloalkyl-(C₁-C₈ alkylenyl);
 Substituted C₈-C₁₀ bicycloalkyl-(C₁-C₈ alkylenyl);
 5- or 6-membered heterocycloalkyl-(C₁-C₈ alkylenyl);
10 Substituted 5- or 6-membered heterocycloalkyl-(C₁-C₈ alkylenyl);
 8- to 10-membered heterobicycloalkyl-(C₁-C₈ alkylenyl);
 Substituted 8- to 10-membered heterobicycloalkyl-(C₁-C₈ alkylenyl);
 Phenyl-(C₁-C₈ alkylenyl);
 Substituted phenyl-(C₁-C₈ alkylenyl);
15 Naphthyl-(C₁-C₈ alkylenyl);
 Substituted naphthyl-(C₁-C₈ alkylenyl);
 5- or 6-membered heteroaryl-(C₁-C₈ alkylenyl);
 Substituted 5- or 6-membered heteroaryl-(C₁-C₈ alkylenyl);
 8- to 10-membered heterobiaryl-(C₁-C₈ alkylenyl);
20 Substituted 8- to 10-membered heterobiaryl-(C₁-C₈ alkylenyl);
 Phenyl;
 Substituted phenyl;
 Naphthyl;
 Substituted naphthyl;
25 5- or 6-membered heteroaryl;
 Substituted 5- or 6-membered heteroaryl;
 8- to 10-membered heterobiaryl; and
 Substituted 8- to 10-membered heterobiaryl;

R² is independently selected from:

- H;
- C₁-C₆ alkyl;
- Phenyl-(C₁-C₈ alkylenyl);
- 5 Substituted phenyl-(C₁-C₈ alkylenyl);
- Naphthyl-(C₁-C₈ alkylenyl);
- Substituted naphthyl-(C₁-C₈ alkylenyl);
- 5- or 6-membered heteroaryl-(C₁-C₈ alkylenyl);
- Substituted 5- or 6-membered heteroaryl-(C₁-C₈ alkylenyl);
- 10 8- to 10-membered heterobiaryl-(C₁-C₈ alkylenyl);
- Substituted 8- to 10-membered heterobiaryl-(C₁-C₈ alkylenyl);
- Phenyl-O-(C₁-C₈ alkylenyl);
- Substituted phenyl-O-(C₁-C₈ alkylenyl);
- Phenyl-S-(C₁-C₈ alkylenyl);
- 15 Substituted phenyl-S-(C₁-C₈ alkylenyl);
- Phenyl-S(O)-(C₁-C₈ alkylenyl);
- Substituted phenyl-S(O)-(C₁-C₈ alkylenyl);
- Phenyl-S(O)₂-(C₁-C₈ alkylenyl); and
- Substituted phenyl-S(O)₂-(C₁-C₈ alkylenyl);
- 20 Each substituted R¹ and R² group contains from 1 to 4 substituents, each independently on a carbon or nitrogen atom, independently selected from:
 - C₁-C₆ alkyl;
 - CN;
 - CF₃;
 - 25 HO;
 - (C₁-C₆ alkyl)-O;
 - (C₁-C₆ alkyl)-S(O)₂;
 - H₂N;
 - (C₁-C₆ alkyl)-N(H);
 - 30 (C₁-C₆ alkyl)₂-N;
 - (C₁-C₆ alkyl)-C(O)O-(C₁-C₈ alkylenyl)_m;
 - (C₁-C₆ alkyl)-C(O)O-(1- to 8-membered heteroalkylenyl)_m;
 - (C₁-C₆ alkyl)-C(O)N(H)-(C₁-C₈ alkylenyl)_m;

(C₁-C₆ alkyl)-C(O)N(H)-(1- to 8-membered heteroalkylenyl)_m;

H₂NS(O)₂-(C₁-C₈ alkylenyl);

(C₁-C₆ alkyl)-N(H)S(O)₂-(C₁-C₈ alkylenyl)_m;

(C₁-C₆ alkyl)₂-NS(O)₂-(C₁-C₈ alkylenyl)_m;

5 3- to 6-membered heterocycloalkyl-(G)_m;

Substituted 3- to 6-membered heterocycloalkyl-(G)_m;

5- or 6-membered heteroaryl-(G)_m;

Substituted 5- or 6-membered heteroaryl-(G)_m;

(C₁-C₆ alkyl)-S(O)₂-N(H)-C(O)-(C₁-C₈ alkylenyl)_m; and

10 (C₁-C₆ alkyl)-C(O)-N(H)-S(O)₂-(C₁-C₈ alkylenyl)_m;


wherein each substituent on a carbon atom may further be independently selected from:


Halo; and

HO₂C;

15 wherein 2 substituents may be taken together with a carbon atom to which they are both bonded to form the group C=O;

wherein two adjacent, substantially sp² carbon atoms may be taken together with a diradical substituent to form a cyclic diradical selected from:





R is H or C₁-C₆ alkyl;

G is CH₂; O, S, S(O); or S(O)₂;

m is an integer of 0 or 1;

5 R⁴ is independently selected from the groups:

H;

CH₃;

CH₃O;

HO;

10 CF₃; and

CN;

wherein each C₈-C₁₀ bicycloalkyl is a bicyclic carbocyclic ring that contains 8-, 9-, or 10-member carbon atoms which are 5,5-fused, 6,5-fused, or 6,6-fused bicyclic rings, respectively, and wherein the ring is saturated or optionally contains one carbon-carbon double bond;

15 wherein each 8- to 10-membered heterobicycloalkyl is a bicyclic ring that contains carbon atoms and from 1 to 4 heteroatoms independently selected from 2 O, 1 S, 1 S(O), 1 S(O)₂, 1 N, 4 N(H), and 4 N(C₁-C₆ alkyl), and wherein when two O atoms or one O atom and one S atom are present, the two O atoms or one O atom and one S atom are not bonded to each other, and wherein the ring is saturated or optionally contains one carbon-carbon or carbon-nitrogen double bond, and wherein the heterobicycloalkyl is a 5,5-fused, 6,5-fused, or 6,6-fused bicyclic ring, respectively,

25 wherein each heterocycloalkyl is a ring that contains carbon atoms and from 1 to 4 heteroatoms independently selected from 2 O, 1 S, 1 S(O), 1 S(O)₂, 1 N, 4 N(H), and 4 N(C₁-C₆ alkyl), and wherein when two O atoms or one O atom and one S atom are present, the two O atoms or one O atom and one

S atom are not bonded to each other, and wherein the ring is saturated or optionally contains one carbon-carbon or carbon-nitrogen double bond; wherein each 5-membered heteroaryl contains carbon atoms and from 1 to 4 heteroatoms independently selected from 1 O, 1 S, 1 N(H), 1 N(C₁-C₆ alkyl), and 4 N, and each 6-membered heteroaryl contains carbon atoms and 1 or 2 heteroatoms independently selected from N, N(H), and N(C₁-C₆ alkyl), and 5- and 6-membered heteroaryl are monocyclic rings; wherein each heterobiaryl contains carbon atoms and from 1 to 4 heteroatoms independently selected from 1 O, 1 S, 1 N(H), 1 N(C₁-C₆ alkyl), and 4 N, and where the 8-, 9-, and 10-membered heterobiaryl are 5,5-fused, 6,5-fused, and 6,6-fused bicyclic rings, respectively, and wherein at least 1 of the 2 fused rings of a bicyclic ring is aromatic, and wherein when the O and S atoms both are present, the O and S atoms are not bonded to each other; wherein with any (C₁-C₆ alkyl)₂-N group, the C₁-C₆ alkyl groups may be optionally taken together with the nitrogen atom to which they are attached to form a 5- or 6-membered heterocycloalkyl; and wherein each group and each substituent recited above is independently selected.

59. The compound according to Embodiment 58, selected from:
3-Benzyl-1-methyl-6-(3-phenyl-prop-2-ynyl)-5,6,7,8-tetrahydro-1H-pyrido[4,3-d]pyrimidine-2,4-dione; and
3-Benzyl-1-methyl-6-(3-pyridin-4-yl-prop-2-ynyl)-5,6,7,8-tetrahydro-1H-pyrido[4,3-d]pyrimidine-2,4-dione; or
a pharmaceutically acceptable salt thereof.

60. The compound according to Embodiment 58, selected from:
3-Benzyl-6-[3-(4-methoxy-phenyl)-prop-2-ynyl]-1-methyl-5,6,7,8-tetrahydro-1H-pyrido[4,3-d]pyrimidine-2,4-dione;
3-Benzyl-6-[3-(3-methoxy-phenyl)-prop-2-ynyl]-1-methyl-5,6,7,8-tetrahydro-1H-pyrido[4,3-d]pyrimidine-2,4-dione; and
3-Benzyl-6-[3-(2-methoxy-pyridin-4-yl)-prop-2-ynyl]-1-methyl-5,6,7,8-tetrahydro-1H-pyrido[4,3-d]pyrimidine-2,4-dione; or

a pharmaceutically acceptable salt thereof.

61. The compound according to Embodiment 58, selected from:
- 5 3-Benzyl-1-methyl-6-(3-pyrrolidin-1-yl-prop-2-ynyl)-5,6,7,8-tetrahydro-1H-pyrido[4,3-d]pyrimidine-2,4-dione;
- 3-Benzyl-1-methyl-6-(3-pyrrol-1-yl-prop-2-ynyl)-5,6,7,8-tetrahydro-1H-pyrido[4,3-d]pyrimidine-2,4-dione;
- 3-Benzyl-1-methyl-6-(3-piperidin-1-yl-prop-2-ynyl)-5,6,7,8-tetrahydro-1H-pyrido[4,3-d]pyrimidine-2,4-dione; and
- 10 3-Benzyl-1-methyl-6-[3-(4-methyl-piperazin-1-yl)-prop-2-ynyl]-5,6,7,8-tetrahydro-1H-pyrido[4,3-d]pyrimidine-2,4-dione; or
- a pharmaceutically acceptable salt thereof.
62. The compound according to Embodiment 58, selected from:
- 15 3-Benzyl-6-(3-furan-2-yl-prop-2-ynyl)-1-methyl-5,6,7,8-tetrahydro-1H-pyrido[4,3-d]pyrimidine-2,4-dione;
- 3-Benzyl-1-methyl-6-(3-thiophen-2-yl-prop-2-ynyl)-5,6,7,8-tetrahydro-1H-pyrido[4,3-d]pyrimidine-2,4-dione;
- 3-Benzyl-1-methyl-6-(3-oxazol-2-yl-prop-2-ynyl)-5,6,7,8-tetrahydro-1H-pyrido[4,3-d]pyrimidine-2,4-dione; and
- 20 3-Benzyl-1-methyl-6-(3-thiazol-2-yl-prop-2-ynyl)-5,6,7,8-tetrahydro-1H-pyrido[4,3-d]pyrimidine-2,4-dione; or
- a pharmaceutically acceptable salt thereof.
63. The compound according to Embodiment 58, selected from:
- 25 3-Benzyl-6-[3-(1H-indol-2-yl)-prop-2-ynyl]-1-methyl-5,6,7,8-tetrahydro-1H-pyrido[4,3-d]pyrimidine-2,4-dione;
- 3-Benzyl-1-methyl-6-[3-(1-methyl-1H-indol-2-yl)-prop-2-ynyl]-5,6,7,8-tetrahydro-1H-pyrido[4,3-d]pyrimidine-2,4-dione;
- 30 6-(3-Benzo[b]thiophen-2-yl-prop-2-ynyl)-3-benzyl-1-methyl-5,6,7,8-tetrahydro-1H-pyrido[4,3-d]pyrimidine-2,4-dione;

6-(3-Benzofuran-2-yl-prop-2-ynyl)-3-benzyl-1-methyl-5,6,7,8-tetrahydro-
1H-pyrido[4,3-d]pyrimidine-2,4-dione; and
6-(3-Benzoxazol-2-yl-prop-2-ynyl)-3-benzyl-1-methyl-5,6,7,8-
tetrahydro-1H-pyrido[4,3-d]pyrimidine-2,4-dione; or
5 a pharmaceutically acceptable salt thereof.

64. The compound according to Embodiment 58, selected from:

3-Benzyl-6-[3-(4-fluoro-phenyl)-prop-2-ynyl]-1-methyl-5,6,7,8-
tetrahydro-1H-pyrido[4,3-d]pyrimidine-2,4-dione;

10 3-Benzyl-1-methyl-6-(3-pentafluorophenyl-prop-2-ynyl)-5,6,7,8-
tetrahydro-1H-pyrido[4,3-d]pyrimidine-2,4-dione;

3-Benzyl-6-[3-(4-chloro-phenyl)-prop-2-ynyl]-1-methyl-5,6,7,8-
tetrahydro-1H-pyrido[4,3-d]pyrimidine-2,4-dione;

15 3-Benzyl-6-[3-(4-bromo-phenyl)-prop-2-ynyl]-1-methyl-5,6,7,8-
tetrahydro-1H-pyrido[4,3-d]pyrimidine-2,4-dione;

3-Benzyl-6-[3-(4-iodo-phenyl)-prop-2-ynyl]-1-methyl-5,6,7,8-tetrahydro-
1H-pyrido[4,3-d]pyrimidine-2,4-dione;

3-Benzyl-1-methyl-6-[3-(4-trifluoromethyl-phenyl)-prop-2-ynyl]-5,6,7,8-
tetrahydro-1H-pyrido[4,3-d]pyrimidine-2,4-dione;

20 3-Benzyl-1-methyl-6-[3-(4-trichloromethyl-phenyl)-prop-2-ynyl]-5,6,7,8-
tetrahydro-1H-pyrido[4,3-d]pyrimidine-2,4-dione; and

3-Benzyl-1-methyl-6-[3-(4-trifluoromethoxy-phenyl)-prop-2-ynyl]-
5,6,7,8-tetrahydro-1H-pyrido[4,3-d]pyrimidine-2,4-dione; or
a pharmaceutically acceptable salt thereof.

25

65. The compound according to Embodiment 58, selected from:

3-Benzyl-1-methyl-6-(3-[1,2,4]triazol-4-yl-prop-2-ynyl)-5,6,7,8-
tetrahydro-1H-pyrido[4,3-d]pyrimidine-2,4-dione;

3-Benzyl-6-(3-imidazol-1-yl-prop-2-ynyl)-1-methyl-5,6,7,8-tetrahydro-
1H-pyrido[4,3-d]pyrimidine-2,4-dione;

30

3-Benzyl-1-methyl-6-[3-(4-methyl-imidazol-1-yl)-prop-2-ynyl]-5,6,7,8-
tetrahydro-1H-pyrido[4,3-d]pyrimidine-2,4-dione;


- 3-Benzyl-1-methyl-6-[3-(4-methyl-4H-[1,2,4]triazol-3-yl)-prop-2-ynyl]-
5,6,7,8-tetrahydro-1H-pyrido[4,3-d]pyrimidine-2,4-dione;
3-Benzyl-1-methyl-6-[3-(1-methyl-1H-imidazol-2-yl)-prop-2-ynyl]-
5,6,7,8-tetrahydro-1H-pyrido[4,3-d]pyrimidine-2,4-dione;
5 3-Benzyl-1-methyl-6-[3-(3-methyl-3H-imidazol-4-yl)-prop-2-ynyl]-
5,6,7,8-tetrahydro-1H-pyrido[4,3-d]pyrimidine-2,4-dione; and
3-Benzyl-1-methyl-6-[3-(1-methyl-1H-tetrazol-5-yl)-prop-2-ynyl]-5,6,7,8-
tetrahydro-1H-pyrido[4,3-d]pyrimidine-2,4-dione; or
a pharmaceutically acceptable salt thereof.
- 10
66. The compound according to Embodiment 58, selected from:
- 4-[1-Methyl-2,4-dioxo-6-(3-phenyl-prop-2-ynyl)-1,4,5,6,7,8-hexahydro-
2H-pyrido[4,3-d]pyrimidin-3-ylmethyl]-benzoic acid;
4-[1-Methyl-2,4-dioxo-6-(3-phenyl-prop-2-ynyl)-1,4,5,6,7,8-hexahydro-
15 2H-pyrido[4,3-d]pyrimidin-3-ylmethyl]-benzoic acid methyl ester;
3-(4-Methoxy-benzyl)-1-methyl-6-(3-phenyl-prop-2-ynyl)-5,6,7,8-
tetrahydro-1H-pyrido[4,3-d]pyrimidine-2,4-dione;
1-Methyl-3-(4-methylsulfanyl-benzyl)-6-(3-phenyl-prop-2-ynyl)-5,6,7,8-
tetrahydro-1H-pyrido[4,3-d]pyrimidine-2,4-dione;
20 3-(4-Methanesulfonyl-benzyl)-1-methyl-6-(3-phenyl-prop-2-ynyl)-5,6,7,8-
tetrahydro-1H-pyrido[4,3-d]pyrimidine-2,4-dione;
4-[1-Methyl-2,4-dioxo-6-(3-phenyl-prop-2-ynyl)-1,4,5,6,7,8-hexahydro-
2H-pyrido[4,3-d]pyrimidin-3-ylmethyl]-benzenesulfonic acid;
4-[1-Methyl-2,4-dioxo-6-(3-phenyl-prop-2-ynyl)-1,4,5,6,7,8-hexahydro-
25 2H-pyrido[4,3-d]pyrimidin-3-ylmethyl]-benzenesulfonamide;
N,N-Dimethyl-4-[1-methyl-2,4-dioxo-6-(3-phenyl-prop-2-ynyl)-
1,4,5,6,7,8-hexahydro-2H-pyrido[4,3-d]pyrimidin-3-ylmethyl]-
benzenesulfonamide;
3-(4-Dimethylamino-benzyl)-1-methyl-6-(3-phenyl-prop-2-ynyl)-5,6,7,8-
30 tetrahydro-1H-pyrido[4,3-d]pyrimidine-2,4-dione;
4-[1-Methyl-2,4-dioxo-6-(3-phenyl-prop-2-ynyl)-1,4,5,6,7,8-hexahydro-
2H-pyrido[4,3-d]pyrimidin-3-ylmethyl]-benzonitrile;

- 3-(4-Fluoro-benzyl)-1-methyl-6-(3-phenyl-prop-2-ynyl)-5,6,7,8-tetrahydro-1H-pyrido[4,3-d]pyrimidine-2,4-dione;
3-(4-Fluoro-benzyl)-1-methyl-6-(3-phenyl-prop-2-ynyl)-5,6,7,8-tetrahydro-1H-pyrido[4,3-d]pyrimidine-2,4-dione;
5 3-(4-Fluoro-benzyl)-1-methyl-6-(3-phenyl-prop-2-ynyl)-5,6,7,8-tetrahydro-1H-pyrido[4,3-d]pyrimidine-2,4-dione;
1-Methyl-6-(3-phenyl-prop-2-ynyl)-3-(4-trichloromethyl-benzyl)-5,6,7,8-tetrahydro-1H-pyrido[4,3-d]pyrimidine-2,4-dione;
1-Methyl-6-(3-phenyl-prop-2-ynyl)-3-(4-trifluoromethyl-benzyl)-5,6,7,8-tetrahydro-1H-pyrido[4,3-d]pyrimidine-2,4-dione; and
10 1-Methyl-3-(4-methyl-benzyl)-6-(3-phenyl-prop-2-ynyl)-5,6,7,8-tetrahydro-1H-pyrido[4,3-d]pyrimidine-2,4-dione; or
a pharmaceutically acceptable salt thereof.
- 15 67. The compound according to Embodiment 58, selected from:
1-Methyl-6-(3-phenyl-prop-2-ynyl)-3-(4-pyrrolidin-1-yl-benzyl)-5,6,7,8-tetrahydro-1H-pyrido[4,3-d]pyrimidine-2,4-dione;
1-Methyl-6-(3-phenyl-prop-2-ynyl)-3-(4-pyrrol-1-yl-benzyl)-5,6,7,8-tetrahydro-1H-pyrido[4,3-d]pyrimidine-2,4-dione;
20 3-(4-Imidazol-1-yl-benzyl)-1-methyl-6-(3-phenyl-prop-2-ynyl)-5,6,7,8-tetrahydro-1H-pyrido[4,3-d]pyrimidine-2,4-dione;
1-Methyl-6-(3-phenyl-prop-2-ynyl)-3-(4-[1,2,4]triazol-4-yl-benzyl)-5,6,7,8-tetrahydro-1H-pyrido[4,3-d]pyrimidine-2,4-dione;
1-Methyl-6-(3-phenyl-prop-2-ynyl)-3-(4-tetrazol-1-yl-benzyl)-5,6,7,8-tetrahydro-1H-pyrido[4,3-d]pyrimidine-2,4-dione;
25 1-Methyl-3-[4-(1-methyl-1H-tetrazol-5-yl)-benzyl]-6-(3-phenyl-prop-2-ynyl)-5,6,7,8-tetrahydro-1H-pyrido[4,3-d]pyrimidine-2,4-dione;
1-Methyl-6-(3-phenyl-prop-2-ynyl)-3-[4-(1H-tetrazol-5-yl)-benzyl]-5,6,7,8-tetrahydro-1H-pyrido[4,3-d]pyrimidine-2,4-dione;
30 1-Methyl-6-(3-phenyl-prop-2-ynyl)-3-[4-(1H-pyrrol-2-yl)-benzyl]-5,6,7,8-tetrahydro-1H-pyrido[4,3-d]pyrimidine-2,4-dione;
1-Methyl-6-(3-phenyl-prop-2-ynyl)-3-(4-thiophen-2-yl-benzyl)-5,6,7,8-tetrahydro-1H-pyrido[4,3-d]pyrimidine-2,4-dione;

3-(4-Furan-2-yl-benzyl)-1-methyl-6-(3-phenyl-prop-2-ynyl)-5,6,7,8-tetrahydro-1H-pyrido[4,3-d]pyrimidine-2,4-dione;
1-Methyl-3-(4-oxazol-2-yl-benzyl)-6-(3-phenyl-prop-2-ynyl)-5,6,7,8-tetrahydro-1H-pyrido[4,3-d]pyrimidine-2,4-dione;
5 1-Methyl-6-(3-phenyl-prop-2-ynyl)-3-(4-thiazol-2-yl-benzyl)-5,6,7,8-tetrahydro-1H-pyrido[4,3-d]pyrimidine-2,4-dione; and
1-Methyl-3-[4-(1-methyl-1H-imidazol-2-yl)-benzyl]-6-(3-phenyl-prop-2-ynyl)-5,6,7,8-tetrahydro-1H-pyrido[4,3-d]pyrimidine-2,4-dione; or
a pharmaceutically acceptable salt thereof.

10

68. A compound of Formula VI



VI

or a pharmaceutically acceptable salt thereof, or a pyrido-N-oxide thereof,
wherein:

15

R¹ is independently selected from:

- C₅ or C₆ cycloalkyl-(C₁-C₈ alkylenyl);
- Substituted C₅ or C₆ cycloalkyl-(C₁-C₈ alkylenyl);
- C₈-C₁₀ bicycloalkyl-(C₁-C₈ alkylenyl);
- Substituted C₈-C₁₀ bicycloalkyl-(C₁-C₈ alkylenyl);
- 20 5- or 6-membered heterocycloalkyl-(C₁-C₈ alkylenyl);
- Substituted 5- or 6-membered heterocycloalkyl-(C₁-C₈ alkylenyl);
- 8- to 10-membered heterobicycloalkyl-(C₁-C₈ alkylenyl);
- Substituted 8- to 10-membered heterobicycloalkyl-(C₁-C₈ alkylenyl);
- Phenyl-(C₁-C₈ alkylenyl);
- 25 Substituted phenyl-(C₁-C₈ alkylenyl);
- Naphthyl-(C₁-C₈ alkylenyl);
- Substituted naphthyl-(C₁-C₈ alkylenyl);
- 5- or 6-membered heteroaryl-(C₁-C₈ alkylenyl);

- Substituted 5- or 6-membered heteroaryl-(C₁-C₈ alkylene);
8- to 10-membered heterobiaryl-(C₁-C₈ alkylene);
Substituted 8- to 10-membered heterobiaryl-(C₁-C₈ alkylene);
Phenyl;
5 Substituted phenyl;
Naphthyl;
Substituted naphthyl;
5- or 6-membered heteroaryl;
Substituted 5- or 6-membered heteroaryl;
10 8- to 10-membered heterobiaryl; and
Substituted 8- to 10-membered heterobiaryl;

R² is independently selected from:

- H;
C₁-C₆ alkyl;
15 Phenyl-(C₁-C₈ alkylene);
Substituted phenyl-(C₁-C₈ alkylene);
Naphthyl-(C₁-C₈ alkylene);
Substituted naphthyl-(C₁-C₈ alkylene);
5- or 6-membered heteroaryl-(C₁-C₈ alkylene);
20 Substituted 5- or 6-membered heteroaryl-(C₁-C₈ alkylene);
8- to 10-membered heterobiaryl-(C₁-C₈ alkylene);
Substituted 8- to 10-membered heterobiaryl-(C₁-C₈ alkylene);
Phenyl-O-(C₁-C₈ alkylene);
Substituted phenyl-O-(C₁-C₈ alkylene);
25 Phenyl-S-(C₁-C₈ alkylene);
Substituted phenyl-S-(C₁-C₈ alkylene);
Phenyl-S(O)-(C₁-C₈ alkylene);
Substituted phenyl-S(O)-(C₁-C₈ alkylene);
Phenyl-S(O)₂-(C₁-C₈ alkylene); and
30 Substituted phenyl-S(O)₂-(C₁-C₈ alkylene);

Each substituted R¹ and R² group contains from 1 to 4 substituents, each independently on a carbon or nitrogen atom, independently selected from:

C₁-C₆ alkyl;


- CN;
 CF₃;
 HO;
 (C₁-C₆ alkyl)-O;
 5 (C₁-C₆ alkyl)-S(O)₂;
 H₂N;
 (C₁-C₆ alkyl)-N(H);
 (C₁-C₆ alkyl)₂-N;
 (C₁-C₆ alkyl)-C(O)O-(C₁-C₈ alkylenyl)_m;
 10 (C₁-C₆ alkyl)-C(O)O-(1- to 8-membered heteroalkylenyl)_m;
 (C₁-C₆ alkyl)-C(O)N(H)-(C₁-C₈ alkylenyl)_m;
 (C₁-C₆ alkyl)-C(O)N(H)-(1- to 8-membered heteroalkylenyl)_m;
 H₂NS(O)₂-(C₁-C₈ alkylenyl);
 (C₁-C₆ alkyl)-N(H)S(O)₂-(C₁-C₈ alkylenyl)_m;
 15 (C₁-C₆ alkyl)₂-NS(O)₂-(C₁-C₈ alkylenyl)_m;
 3- to 6-membered heterocycloalkyl-(G)_m;
 Substituted 3- to 6-membered heterocycloalkyl-(G)_m;
 5- or 6-membered heteroaryl-(G)_m;
 Substituted 5- or 6-membered heteroaryl-(G)_m;
 20 (C₁-C₆ alkyl)-S(O)₂-N(H)-C(O)-(C₁-C₈ alkylenyl)_m; and
 (C₁-C₆ alkyl)-C(O)-N(H)-S(O)₂-(C₁-C₈ alkylenyl)_m;


wherein each substituent on a carbon atom may further be independently selected from:

- Halo; and
 25 HO₂C;

wherein 2 substituents may be taken together with a carbon atom to which they are both bonded to form the group C=O;

wherein two adjacent, substantially sp² carbon atoms may be taken together with a diradical substituent to form a cyclic diradical selected from:





5 R is H or C₁-C₆ alkyl;

G is CH₂; O, S, S(O); or S(O)₂;

m is an integer of 0 or 1;

R⁴ is independently selected from the groups:

H;

10 CH₃;

CH₃O;

HO;

CF₃; and

CN;

15 wherein each C₈-C₁₀ bicycloalkyl is a bicyclic carbocyclic ring that contains 8-, 9-, or 10-member carbon atoms which are 5,5-fused, 6,5-fused, or 6,6-fused bicyclic rings, respectively, and wherein the ring is saturated or optionally contains one carbon-carbon double bond;

20 wherein each 8- to 10-membered heterobicycloalkyl is a bicyclic ring that contains carbon atoms and from 1 to 4 heteroatoms independently selected from 2 O, 1 S, 1 S(O), 1 S(O)₂, 1 N, 4 N(H), and 4 N(C₁-C₆ alkyl), and wherein when two

O atoms or one O atom and one S atom are present, the two O atoms or one O atom and one S atom are not bonded to each other, and wherein the ring is saturated or optionally contains one carbon-carbon or carbon-nitrogen double bond, and wherein the heterobicycloalkyl is a 5,5-fused, 6,5-fused, or 6,6-fused bicyclic ring, respectively,

wherein each heterocycloalkyl is a ring that contains carbon atoms and from 1 to 4 heteroatoms independently selected from 2 O, 1 S, 1 S(O), 1 S(O)₂, 1 N, 4 N(H), and 4 N(C₁-C₆ alkyl), and wherein when two O atoms or one O atom and one S atom are present, the two O atoms or one O atom and one S atom are not bonded to each other, and wherein the ring is saturated or optionally contains one carbon-carbon or carbon-nitrogen double bond;

wherein each 5-membered heteroaryl contains carbon atoms and from 1 to 4 heteroatoms independently selected from 1 O, 1 S, 1 N(H), 1 N(C₁-C₆ alkyl), and 4 N, and each 6-membered heteroaryl contains carbon atoms and 1 or 2 heteroatoms independently selected from N, N(H), and N(C₁-C₆ alkyl), and 5- and 6-membered heteroaryl are monocyclic rings;

wherein each heterobiaryl contains carbon atoms and from 1 to 4 heteroatoms independently selected from 1 O, 1 S, 1 N(H), 1 N(C₁-C₆ alkyl), and 4 N, and where the 8-, 9-, and 10-membered heterobiaryl are 5,5-fused, 6,5-fused, and 6,6-fused bicyclic rings, respectively, and wherein at least 1 of the 2 fused rings of a bicyclic ring is aromatic, and wherein when the O and S atoms both are present, the O and S atoms are not bonded to each other;

wherein with any (C₁-C₆ alkyl)₂-N group, the C₁-C₆ alkyl groups may be optionally taken together with the nitrogen atom to which they are attached to form a 5- or 6-membered heterocycloalkyl; and

wherein each group and each substituent recited above is independently selected.

69. The compound according to Embodiment 68, selected from:

3-Benzyl-1-methyl-4-oxo-1,3,4,5,7,8-hexahydro-2H-pyrido[4,3-d]pyrimidine-6-carboxylic acid benzyl ester; and

3-Benzyl-1-methyl-4-oxo-1,3,4,5,7,8-hexahydro-2H-pyrido[4,3-d]pyrimidine-6-carboxylic acid pyridin-4-ylmethyl ester; or

a pharmaceutically acceptable salt thereof.

70. The compound according to Embodiment 68, selected from:

3-Benzyl-1-methyl-4-oxo-1,3,4,5,7,8-hexahydro-2H-pyrido[4,3-

5 d]pyrimidine-6-carboxylic acid 4-methoxy-benzyl ester;

3-Benzyl-1-methyl-4-oxo-1,3,4,5,7,8-hexahydro-2H-pyrido[4,3-

d]pyrimidine-6-carboxylic acid 3-methoxy-benzyl ester; and

3-Benzyl-1-methyl-4-oxo-1,3,4,5,7,8-hexahydro-2H-pyrido[4,3-

d]pyrimidine-6-carboxylic acid 2-methoxy-pyridin-4-ylmethyl

10 ester; or

a pharmaceutically acceptable salt thereof.

71. The compound according to Embodiment 68, selected from:

3-Benzyl-1-methyl-4-oxo-1,3,4,5,7,8-hexahydro-2H-pyrido[4,3-

15 d]pyrimidine-6-carboxylic acid pyrrolidin-1-ylmethyl ester;

3-Benzyl-1-methyl-4-oxo-1,3,4,5,7,8-hexahydro-2H-pyrido[4,3-

d]pyrimidine-6-carboxylic acid pyrrol-1-ylmethyl ester;

3-Benzyl-1-methyl-4-oxo-1,3,4,5,7,8-hexahydro-2H-pyrido[4,3-

d]pyrimidine-6-carboxylic acid imidazol-1-ylmethyl ester;

20 3-Benzyl-1-methyl-4-oxo-1,3,4,5,7,8-hexahydro-2H-pyrido[4,3-

d]pyrimidine-6-carboxylic acid [1,2,4]triazol-4-ylmethyl ester; and

3-Benzyl-1-methyl-4-oxo-1,3,4,5,7,8-hexahydro-2H-pyrido[4,3-

d]pyrimidine-6-carboxylic acid tetrazol-1-ylmethyl ester; or

a pharmaceutically acceptable salt thereof.

25

72. The compound according to Embodiment 68, selected from:

3-Benzyl-1-methyl-4-oxo-1,3,4,5,7,8-hexahydro-2H-pyrido[4,3-

d]pyrimidine-6-carboxylic acid 4-methyl-piperazin-1-ylmethyl

ester; and

30 3-Benzyl-1-methyl-4-oxo-1,3,4,5,7,8-hexahydro-2H-pyrido[4,3-


d]pyrimidine-6-carboxylic acid piperidin-1-ylmethyl ester; or

a pharmaceutically acceptable salt thereof.

73. The compound according to Embodiment 68, selected from:
- 3-Benzyl-1-methyl-4-oxo-1,3,4,5,7,8-hexahydro-2H-pyrido[4,3-
d]pyrimidine-6-carboxylic acid 1,3-dihydro-indol-2-ylmethyl
5 ester;
- 3-Benzyl-1-methyl-4-oxo-1,3,4,5,7,8-hexahydro-2H-pyrido[4,3-
d]pyrimidine-6-carboxylic acid 1H-indol-2-ylmethyl ester;
- 3-Benzyl-1-methyl-4-oxo-1,3,4,5,7,8-hexahydro-2H-pyrido[4,3-
d]pyrimidine-6-carboxylic acid 1-methyl-1H-indol-2-ylmethyl
10 ester;
- 3-Benzyl-1-methyl-4-oxo-1,3,4,5,7,8-hexahydro-2H-pyrido[4,3-
d]pyrimidine-6-carboxylic acid benzo[b]thiophen-2-ylmethyl ester;
and
- 3-Benzyl-1-methyl-4-oxo-1,3,4,5,7,8-hexahydro-2H-pyrido[4,3-
d]pyrimidine-6-carboxylic acid benzofuran-2-ylmethyl ester; or
15 a pharmaceutically acceptable salt thereof.
74. The compound according to Embodiment 68, selected from:
- 3-Benzyl-1-methyl-4-oxo-1,3,4,5,7,8-hexahydro-2H-pyrido[4,3-
d]pyrimidine-6-carboxylic acid 4-fluoro-benzyl ester;
20
- 3-Benzyl-1-methyl-4-oxo-1,3,4,5,7,8-hexahydro-2H-pyrido[4,3-
d]pyrimidine-6-carboxylic acid 4-chloro-benzyl ester;
- 3-Benzyl-1-methyl-4-oxo-1,3,4,5,7,8-hexahydro-2H-pyrido[4,3-
d]pyrimidine-6-carboxylic acid 4-bromo-benzyl ester;
- 25 3-Benzyl-1-methyl-4-oxo-1,3,4,5,7,8-hexahydro-2H-pyrido[4,3-
d]pyrimidine-6-carboxylic acid 4-iodo-benzyl ester;
- 3-Benzyl-1-methyl-4-oxo-1,3,4,5,7,8-hexahydro-2H-pyrido[4,3-
d]pyrimidine-6-carboxylic acid 4-trichloromethyl-benzyl ester;
- 3-Benzyl-1-methyl-4-oxo-1,3,4,5,7,8-hexahydro-2H-pyrido[4,3-
d]pyrimidine-6-carboxylic acid 4-trifluoromethyl-benzyl ester; and
30
- 3-Benzyl-1-methyl-4-oxo-1,3,4,5,7,8-hexahydro-2H-pyrido[4,3-
d]pyrimidine-6-carboxylic acid 4-trifluoromethoxy-benzyl ester; or

a pharmaceutically acceptable salt thereof.

75. A compound of Formula VII



VII

5 or a pharmaceutically acceptable salt thereof, or a pyrido-N-oxide thereof, wherein:

R¹ is independently selected from:

- C₅ or C₆ cycloalkyl-(C₁-C₈ alkylenyl);
- Substituted C₅ or C₆ cycloalkyl-(C₁-C₈ alkylenyl);
- 10 C₈-C₁₀ bicycloalkyl-(C₁-C₈ alkylenyl);
- Substituted C₈-C₁₀ bicycloalkyl-(C₁-C₈ alkylenyl);
- 5- or 6-membered heterocycloalkyl-(C₁-C₈ alkylenyl);
- Substituted 5- or 6-membered heterocycloalkyl-(C₁-C₈ alkylenyl);
- 8- to 10-membered heterobicycloalkyl-(C₁-C₈ alkylenyl);
- 15 Substituted 8- to 10-membered heterobicycloalkyl-(C₁-C₈ alkylenyl);
- Phenyl-(C₁-C₈ alkylenyl);
- Substituted phenyl-(C₁-C₈ alkylenyl);
- Naphthyl-(C₁-C₈ alkylenyl);
- Substituted naphthyl-(C₁-C₈ alkylenyl);
- 20 5- or 6-membered heteroaryl-(C₁-C₈ alkylenyl);
- Substituted 5- or 6-membered heteroaryl-(C₁-C₈ alkylenyl);
- 8- to 10-membered heterobiaryl-(C₁-C₈ alkylenyl);
- Substituted 8- to 10-membered heterobiaryl-(C₁-C₈ alkylenyl);
- Phenyl;
- 25 Substituted phenyl;
- Naphthyl;
- Substituted naphthyl;
- 5- or 6-membered heteroaryl;

Substituted 5- or 6-membered heteroaryl;
8- to 10-membered heterobiaryl; and
Substituted 8- to 10-membered heterobiaryl;

R² is independently selected from:

- 5 H;
 C₁-C₆ alkyl;
 Phenyl-(C₁-C₈ alkylene);
 Substituted phenyl-(C₁-C₈ alkylene);
 Naphthyl-(C₁-C₈ alkylene);
10 Substituted naphthyl-(C₁-C₈ alkylene);
 5- or 6-membered heteroaryl-(C₁-C₈ alkylene);
 Substituted 5- or 6-membered heteroaryl-(C₁-C₈ alkylene);
 8- to 10-membered heterobiaryl-(C₁-C₈ alkylene);
 Substituted 8- to 10-membered heterobiaryl-(C₁-C₈ alkylene);
15 Phenyl-O-(C₁-C₈ alkylene);
 Substituted phenyl-O-(C₁-C₈ alkylene);
 Phenyl-S-(C₁-C₈ alkylene);
 Substituted phenyl-S-(C₁-C₈ alkylene);
 Phenyl-S(O)-(C₁-C₈ alkylene);
20 Substituted phenyl-S(O)-(C₁-C₈ alkylene);
 Phenyl-S(O)₂-(C₁-C₈ alkylene); and
 Substituted phenyl-S(O)₂-(C₁-C₈ alkylene);

Each substituted R¹ and R² group contains from 1 to 4 substituents, each independently on a carbon or nitrogen atom, independently selected from:

- 25 C₁-C₆ alkyl;
 CN;
 CF₃;
 HO;
 (C₁-C₆ alkyl)-O;
30 (C₁-C₆ alkyl)-S(O)₂;
 H₂N;
 (C₁-C₆ alkyl)-N(H);
 (C₁-C₆ alkyl)₂-N;


$(C_1-C_6 \text{ alkyl})-C(O)O-(C_1-C_8 \text{ alkylenyl})_m$;
 $(C_1-C_6 \text{ alkyl})-C(O)O-(1\text{- to }8\text{-membered heteroalkylenyl})_m$;
 $(C_1-C_6 \text{ alkyl})-C(O)N(H)-(C_1-C_8 \text{ alkylenyl})_m$;
 $(C_1-C_6 \text{ alkyl})-C(O)N(H)-(1\text{- to }8\text{-membered heteroalkylenyl})_m$;
5 $H_2NS(O)_2-(C_1-C_8 \text{ alkylenyl})$;
 $(C_1-C_6 \text{ alkyl})-N(H)S(O)_2-(C_1-C_8 \text{ alkylenyl})_m$;
 $(C_1-C_6 \text{ alkyl})_2-NS(O)_2-(C_1-C_8 \text{ alkylenyl})_m$;
3- to 6-membered heterocycloalkyl-(G)_m;
Substituted 3- to 6-membered heterocycloalkyl-(G)_m;
10 5- or 6-membered heteroaryl-(G)_m;
Substituted 5- or 6-membered heteroaryl-(G)_m;
 $(C_1-C_6 \text{ alkyl})-S(O)_2-N(H)-C(O)-(C_1-C_8 \text{ alkylenyl})_m$; and
 $(C_1-C_6 \text{ alkyl})-C(O)-N(H)-S(O)_2-(C_1-C_8 \text{ alkylenyl})_m$;


wherein each substituent on a carbon atom may further be independently selected
 from:

Halo; and
 HO_2C ;

wherein 2 substituents may be taken together with a carbon atom to which they
 are both bonded to form the group $C=O$;

20 wherein two adjacent, substantially sp^2 carbon atoms may be taken together with a
 diradical substituent to form a cyclic diradical selected from:





R is H or C₁-C₆ alkyl;

G is CH₂, O, S, S(O); or S(O)₂;

5 m is an integer of 0 or 1;

R⁴ is independently selected from the groups:

H;

CH₃;

CH₃O;

10 HO;

CF₃; and

CN;

wherein each C₈-C₁₀ bicycloalkyl is a bicyclic carbocyclic ring that contains 8-, 9-, or 10-member carbon atoms which are 5,5-fused, 6,5-fused, or 6,6-fused bicyclic rings, respectively, and wherein the ring is saturated or optionally contains one carbon-carbon double bond;

wherein each 8- to 10-membered heterobicycloalkyl is a bicyclic ring that contains carbon atoms and from 1 to 4 heteroatoms independently selected from 2 O, 1 S, 1 S(O), 1 S(O)₂, 1 N, 4 N(H), and 4 N(C₁-C₆ alkyl), and wherein when two O atoms or one O atom and one S atom are present, the two O atoms or one O atom and one S atom are not bonded to each other, and wherein the ring is saturated or optionally contains one carbon-carbon or carbon-nitrogen double bond, and wherein the heterobicycloalkyl is a 5,5-fused, 6,5-fused, or 6,6-fused bicyclic ring, respectively,

25 wherein each heterocycloalkyl is a ring that contains carbon atoms and from 1 to 4 heteroatoms independently selected from 2 O, 1 S, 1 S(O), 1 S(O)₂, 1 N, 4

N(H), and 4 N(C₁-C₆ alkyl), and wherein when two O atoms or one O atom and one S atom are present, the two O atoms or one O atom and one S atom are not bonded to each other, and wherein the ring is saturated or optionally contains one carbon-carbon or carbon-nitrogen double bond;
5 wherein each 5-membered heteroaryl contains carbon atoms and from 1 to 4 heteroatoms independently selected from 1 O, 1 S, 1 N(H), 1 N(C₁-C₆ alkyl), and 4 N, and each 6-membered heteroaryl contains carbon atoms and 1 or 2 heteroatoms independently selected from N, N(H), and N(C₁-C₆ alkyl), and 5- and 6-membered heteroaryl are monocyclic rings;
10 wherein each heterobiaryl contains carbon atoms and from 1 to 4 heteroatoms independently selected from 1 O, 1 S, 1 N(H), 1 N(C₁-C₆ alkyl), and 4 N, and where the 8-, 9-, and 10-membered heterobiaryl are 5,5-fused, 6,5-fused, and 6,6-fused bicyclic rings, respectively, and wherein at least 1 of the 2 fused rings of a bicyclic ring is aromatic, and wherein when the O
15 and S atoms both are present, the O and S atoms are not bonded to each other;
wherein with any (C₁-C₆ alkyl)₂-N group, the C₁-C₆ alkyl groups may be optionally taken together with the nitrogen atom to which they are attached to form a 5- or 6-membered heterocycloalkyl; and
20 wherein each group and each substituent recited above is independently selected.

76. The compound according to Embodiment 75, selected from:
3-Benzyl-1-methyl-6-(3-phenyl-prop-2-ynyl)-2,3,5,6,7,8-hexahydro-1H-pyrido[4,3-d]pyrimidin-4-one; and
25 3-Benzyl-1-methyl-6-(3-pyridin-4-yl-prop-2-ynyl)-2,3,5,6,7,8-hexahydro-1H-pyrido[4,3-d]pyrimidin-4-one; or
a pharmaceutically acceptable salt thereof.
77. The compound according to Embodiment 75, selected from:
30 3-Benzyl-6-[3-(4-methoxy-phenyl)-prop-2-ynyl]-1-methyl-2,3,5,6,7,8-hexahydro-1H-pyrido[4,3-d]pyrimidin-4-one;

3-Benzyl-6-[3-(3-methoxy-phenyl)-prop-2-ynyl]-1-methyl-2,3,5,6,7,8-hexahydro-1H-pyrido[4,3-d]pyrimidin-4-one; and
3-Benzyl-6-[3-(2-methoxy-pyridin-4-yl)-prop-2-ynyl]-1-methyl-2,3,5,6,7,8-hexahydro-1H-pyrido[4,3-d]pyrimidin-4-one; or
a pharmaceutically acceptable salt thereof.

5

78. The compound according to Embodiment 75, selected from:

3-Benzyl-6-[3-(4-fluoro-phenyl)-prop-2-ynyl]-1-methyl-2,3,5,6,7,8-hexahydro-1H-pyrido[4,3-d]pyrimidin-4-one;

10

3-Benzyl-6-[3-(4-chloro-phenyl)-prop-2-ynyl]-1-methyl-2,3,5,6,7,8-hexahydro-1H-pyrido[4,3-d]pyrimidin-4-one;

3-Benzyl-6-[3-(4-bromo-phenyl)-prop-2-ynyl]-1-methyl-2,3,5,6,7,8-hexahydro-1H-pyrido[4,3-d]pyrimidin-4-one; and

3-Benzyl-6-[3-(4-iodo-phenyl)-prop-2-ynyl]-1-methyl-2,3,5,6,7,8-hexahydro-1H-pyrido[4,3-d]pyrimidin-4-one; or

15

a pharmaceutically acceptable salt thereof.

79. The compound according to Embodiment 75, selected from:

3-Benzyl-1-methyl-6-(3-pyrrolidin-1-yl-prop-2-ynyl)-2,3,5,6,7,8-hexahydro-1H-pyrido[4,3-d]pyrimidin-4-one;

20

3-Benzyl-1-methyl-6-(3-pyrrol-1-yl-prop-2-ynyl)-2,3,5,6,7,8-hexahydro-1H-pyrido[4,3-d]pyrimidin-4-one;

3-Benzyl-6-(3-imidazol-1-yl-prop-2-ynyl)-1-methyl-2,3,5,6,7,8-hexahydro-1H-pyrido[4,3-d]pyrimidin-4-one;

25

3-Benzyl-1-methyl-6-(3-[1,2,4]triazol-4-yl-prop-2-ynyl)-2,3,5,6,7,8-hexahydro-1H-pyrido[4,3-d]pyrimidin-4-one;

3-Benzyl-1-methyl-6-(3-tetrazol-1-yl-prop-2-ynyl)-2,3,5,6,7,8-hexahydro-1H-pyrido[4,3-d]pyrimidin-4-one;

3-Benzyl-1-methyl-6-[3-(4-methyl-imidazol-1-yl)-prop-2-ynyl]-

30

2,3,5,6,7,8-hexahydro-1H-pyrido[4,3-d]pyrimidin-4-one;

3-Benzyl-1-methyl-6-(3-piperidin-1-yl-prop-2-ynyl)-2,3,5,6,7,8-hexahydro-1H-pyrido[4,3-d]pyrimidin-4-one; and

3-Benzyl-1-methyl-6-[3-(4-methyl-piperazin-1-yl)-prop-2-ynyl]-
2,3,5,6,7,8-hexahydro-1H-pyrido[4,3-d]pyrimidin-4-one; or
a pharmaceutically acceptable salt thereof.

- 5 80. The compound according to Embodiment 75, selected from:
3-Benzyl-6-[3-(1H-indol-2-yl)-prop-2-ynyl]-1-methyl-2,3,5,6,7,8-
hexahydro-1H-pyrido[4,3-d]pyrimidin-4-one;
3-Benzyl-1-methyl-6-[3-(1-methyl-1H-indol-2-yl)-prop-2-ynyl]-
2,3,5,6,7,8-hexahydro-1H-pyrido[4,3-d]pyrimidin-4-one;
10 6-(3-Benzo[b]thiophen-2-yl-prop-2-ynyl)-3-benzyl-1-methyl-2,3,5,6,7,8-
hexahydro-1H-pyrido[4,3-d]pyrimidin-4-one; and
6-(3-Benzofuran-2-yl-prop-2-ynyl)-3-benzyl-1-methyl-2,3,5,6,7,8-
hexahydro-1H-pyrido[4,3-d]pyrimidin-4-one; or
a pharmaceutically acceptable salt thereof.
- 15
81. The compound according to Embodiment 75, selected from:
3-Benzyl-1-methyl-6-(3-oxazol-2-yl-prop-2-ynyl)-2,3,5,6,7,8-hexahydro-
1H-pyrido[4,3-d]pyrimidin-4-one;
3-Benzyl-1-methyl-6-(3-[1,3,4]oxadiazol-2-yl-prop-2-ynyl)-2,3,5,6,7,8-
20 hexahydro-1H-pyrido[4,3-d]pyrimidin-4-one;
3-Benzyl-1-methyl-6-(3-oxazol-5-yl-prop-2-ynyl)-2,3,5,6,7,8-hexahydro-
1H-pyrido[4,3-d]pyrimidin-4-one;
3-Benzyl-1-methyl-6-(3-thiazol-2-yl-prop-2-ynyl)-2,3,5,6,7,8-hexahydro-
1H-pyrido[4,3-d]pyrimidin-4-one;
25 3-Benzyl-1-methyl-6-[3-(5-methyl-thiazol-2-yl)-prop-2-ynyl]-2,3,5,6,7,8-
hexahydro-1H-pyrido[4,3-d]pyrimidin-4-one; and
3-Benzyl-1-methyl-6-[3-(1-methyl-1H-imidazol-2-yl)-prop-2-ynyl]-
2,3,5,6,7,8-hexahydro-1H-pyrido[4,3-d]pyrimidin-4-one; or
a pharmaceutically acceptable salt thereof.
- 30
82. The compound according to Embodiment 75, selected from:
4-[1-Methyl-4-oxo-6-(3-phenyl-prop-2-ynyl)-1,4,5,6,7,8-hexahydro-2H-
pyrido[4,3-d]pyrimidin-3-ylmethyl]-benzoic acid;

- 4-[1-Methyl-4-oxo-6-(3-phenyl-prop-2-ynyl)-1,4,5,6,7,8-hexahydro-2H-pyrido[4,3-d]pyrimidin-3-ylmethyl]-benzoic acid methyl ester;
3-(4-Methoxy-benzyl)-1-methyl-6-(3-phenyl-prop-2-ynyl)-2,3,5,6,7,8-hexahydro-1H-pyrido[4,3-d]pyrimidin-4-one;
- 5 1-Methyl-3-(4-methylsulfanyl-benzyl)-6-(3-phenyl-prop-2-ynyl)-2,3,5,6,7,8-hexahydro-1H-pyrido[4,3-d]pyrimidin-4-one;
- 4-[1-Methyl-4-oxo-6-(3-phenyl-prop-2-ynyl)-1,4,5,6,7,8-hexahydro-2H-pyrido[4,3-d]pyrimidin-3-ylmethyl]-benzonitrile;
3-(4-Dimethylamino-benzyl)-1-methyl-6-(3-phenyl-prop-2-ynyl)-2,3,5,6,7,8-hexahydro-1H-pyrido[4,3-d]pyrimidin-4-one;
- 10 3-(4-Aziridin-1-yl-benzyl)-1-methyl-6-(3-phenyl-prop-2-ynyl)-2,3,5,6,7,8-hexahydro-1H-pyrido[4,3-d]pyrimidin-4-one;
- 3-(4-Methanesulfonyl-benzyl)-1-methyl-6-(3-phenyl-prop-2-ynyl)-2,3,5,6,7,8-hexahydro-1H-pyrido[4,3-d]pyrimidin-4-one;
- 15 4-[1-Methyl-4-oxo-6-(3-phenyl-prop-2-ynyl)-1,4,5,6,7,8-hexahydro-2H-pyrido[4,3-d]pyrimidin-3-ylmethyl]-benzenesulfonic acid;
- 4-[1-Methyl-4-oxo-6-(3-phenyl-prop-2-ynyl)-1,4,5,6,7,8-hexahydro-2H-pyrido[4,3-d]pyrimidin-3-ylmethyl]-benzenesulfonamide;
- N,N-Dimethyl-4-[1-methyl-4-oxo-6-(3-phenyl-prop-2-ynyl)-1,4,5,6,7,8-hexahydro-2H-pyrido[4,3-d]pyrimidin-3-ylmethyl]-benzenesulfonamide;
- 20 3-(4-Fluoro-benzyl)-1-methyl-6-(3-phenyl-prop-2-ynyl)-2,3,5,6,7,8-hexahydro-1H-pyrido[4,3-d]pyrimidin-4-one;
- 3-(4-Chloro-benzyl)-1-methyl-6-(3-phenyl-prop-2-ynyl)-2,3,5,6,7,8-hexahydro-1H-pyrido[4,3-d]pyrimidin-4-one;
- 25 3-(4-Bromo-benzyl)-1-methyl-6-(3-phenyl-prop-2-ynyl)-2,3,5,6,7,8-hexahydro-1H-pyrido[4,3-d]pyrimidin-4-one;
- 3-(4-Iodo-benzyl)-1-methyl-6-(3-phenyl-prop-2-ynyl)-2,3,5,6,7,8-hexahydro-1H-pyrido[4,3-d]pyrimidin-4-one;
- 30 1-Methyl-3-pentafluorophenylmethyl-6-(3-phenyl-prop-2-ynyl)-2,3,5,6,7,8-hexahydro-1H-pyrido[4,3-d]pyrimidin-4-one; and
1-Methyl-6-(3-phenyl-prop-2-ynyl)-3-(3,4,5-trifluoro-benzyl)-2,3,5,6,7,8-hexahydro-1H-pyrido[4,3-d]pyrimidin-4-one; or

a pharmaceutically acceptable salt thereof.

83. The compound according to Embodiment 75, selected from:
1-Methyl-6-(3-phenyl-prop-2-ynyl)-3-pyridin-4-ylmethyl-2,3,5,6,7,8-
5 hexahydro-1H-pyrido[4,3-d]pyrimidin-4-one;
1-Methyl-6-(3-phenyl-prop-2-ynyl)-3-pyridin-3-ylmethyl-2,3,5,6,7,8-
hexahydro-1H-pyrido[4,3-d]pyrimidin-4-one; and
1-Methyl-6-(3-phenyl-prop-2-ynyl)-3-pyridin-2-ylmethyl-2,3,5,6,7,8-
hexahydro-1H-pyrido[4,3-d]pyrimidin-4-one; or
10 a pharmaceutically acceptable salt thereof.
84. The compound according to Embodiment 75, selected from:
1-Methyl-3-morpholin-4-ylmethyl-6-(3-phenyl-prop-2-ynyl)-2,3,5,6,7,8-
hexahydro-1H-pyrido[4,3-d]pyrimidin-4-one; and
15 1-Methyl-3-morpholin-4-ylmethyl-6-(3-phenyl-prop-2-ynyl)-2,3,5,6,7,8-
hexahydro-1H-pyrido[4,3-d]pyrimidin-4-one; or
a pharmaceutically acceptable salt thereof.
85. The compound according to Embodiment 75, selected from:
20 1-Methyl-6-(3-phenyl-prop-2-ynyl)-3-pyrrol-1-ylmethyl-2,3,5,6,7,8-
hexahydro-1H-pyrido[4,3-d]pyrimidin-4-one;
3-Furan-2-ylmethyl-1-methyl-6-(3-phenyl-prop-2-ynyl)-2,3,5,6,7,8-
hexahydro-1H-pyrido[4,3-d]pyrimidin-4-one;
1-Methyl-6-(3-phenyl-prop-2-ynyl)-3-thiophen-2-ylmethyl-2,3,5,6,7,8-
25 hexahydro-1H-pyrido[4,3-d]pyrimidin-4-one;
1-Methyl-6-(3-phenyl-prop-2-ynyl)-3-thiazol-2-ylmethyl-2,3,5,6,7,8-
hexahydro-1H-pyrido[4,3-d]pyrimidin-4-one;
1-Methyl-3-oxazol-2-ylmethyl-6-(3-phenyl-prop-2-ynyl)-2,3,5,6,7,8-
hexahydro-1H-pyrido[4,3-d]pyrimidin-4-one;
30 3-(1H-Imidazol-2-ylmethyl)-1-methyl-6-(3-phenyl-prop-2-ynyl)-
2,3,5,6,7,8-hexahydro-1H-pyrido[4,3-d]pyrimidin-4-one;

1-Methyl-3-(1-methyl-1H-imidazol-2-ylmethyl)-6-(3-phenyl-prop-2-ynyl)-
2,3,5,6,7,8-hexahydro-1H-pyrido[4,3-d]pyrimidin-4-one;

1-Methyl-3-(4-methyl-4H-[1,2,4]triazol-3-ylmethyl)-6-(3-phenyl-prop-2-
ynyl)-2,3,5,6,7,8-hexahydro-1H-pyrido[4,3-d]pyrimidin-4-one; and

5 1-Methyl-6-(3-phenyl-prop-2-ynyl)-3-(1H-tetrazol-5-ylmethyl)-
2,3,5,6,7,8-hexahydro-1H-pyrido[4,3-d]pyrimidin-4-one; or
a pharmaceutically acceptable salt thereof.

86. The compound according to Embodiment 75, selected from:


10 1-Methyl-6-(3-phenyl-prop-2-ynyl)-3-(4-tetrazol-1-yl-benzyl)-2,3,5,6,7,8-
hexahydro-1H-pyrido[4,3-d]pyrimidin-4-one; and

6-[3-(4-Methoxy-phenyl)-prop-2-ynyl]-1-methyl-3-(4-tetrazol-1-yl-
benzyl)-2,3,5,6,7,8-hexahydro-1H-pyrido[4,3-d]pyrimidin-4-one;

or

15 a pharmaceutically acceptable salt thereof.

87. A compound of Formula VIII



VIII

or a pharmaceutically acceptable salt thereof, or a pyrido-N-oxide thereof,

20 wherein:

R¹ is independently selected from:

C₅ or C₆ cycloalkyl-(C₁-C₈ alkylenyl);

Substituted C₅ or C₆ cycloalkyl-(C₁-C₈ alkylenyl);

C₈-C₁₀ bicycloalkyl-(C₁-C₈ alkylenyl);

25 Substituted C₈-C₁₀ bicycloalkyl-(C₁-C₈ alkylenyl);

5- or 6-membered heterocycloalkyl-(C₁-C₈ alkylenyl);

Substituted 5- or 6-membered heterocycloalkyl-(C₁-C₈ alkylenyl);

8- to 10-membered heterobicycloalkyl-(C₁-C₈ alkylenyl);

- Substituted 8- to 10-membered heterobicycloalkyl-(C₁-C₈ alkylenyl);
Phenyl-(C₁-C₈ alkylenyl);
Substituted phenyl-(C₁-C₈ alkylenyl);
Naphthyl-(C₁-C₈ alkylenyl);
5 Substituted naphthyl-(C₁-C₈ alkylenyl);
5- or 6-membered heteroaryl-(C₁-C₈ alkylenyl);
Substituted 5- or 6-membered heteroaryl-(C₁-C₈ alkylenyl);
8- to 10-membered heterobiaryl-(C₁-C₈ alkylenyl);
Substituted 8- to 10-membered heterobiaryl-(C₁-C₈ alkylenyl);
10 Phenyl;
Substituted phenyl;
Naphthyl;
Substituted naphthyl;
5- or 6-membered heteroaryl;
15 Substituted 5- or 6-membered heteroaryl;
8- to 10-membered heterobiaryl; and
Substituted 8- to 10-membered heterobiaryl;
- R² is independently selected from:
- H;
20 C₁-C₆ alkyl;
Phenyl-(C₁-C₈ alkylenyl);
Substituted phenyl-(C₁-C₈ alkylenyl);
Naphthyl-(C₁-C₈ alkylenyl);
Substituted naphthyl-(C₁-C₈ alkylenyl);
25 5- or 6-membered heteroaryl-(C₁-C₈ alkylenyl);
Substituted 5- or 6-membered heteroaryl-(C₁-C₈ alkylenyl);
8- to 10-membered heterobiaryl-(C₁-C₈ alkylenyl);
Substituted 8- to 10-membered heterobiaryl-(C₁-C₈ alkylenyl);
Phenyl-O-(C₁-C₈ alkylenyl);
30 Substituted phenyl-O-(C₁-C₈ alkylenyl);
Phenyl-S-(C₁-C₈ alkylenyl);
Substituted phenyl-S-(C₁-C₈ alkylenyl);
Phenyl-S(O)-(C₁-C₈ alkylenyl);

Substituted phenyl-S(O)-(C₁-C₈ alkylenyl);

Phenyl-S(O)₂-(C₁-C₈ alkylenyl); and

Substituted phenyl-S(O)₂-(C₁-C₈ alkylenyl);

Each substituted R¹ and R² group contains from 1 to 4 substituents, each

5 independently on a carbon or nitrogen atom, independently selected from:

C₁-C₆ alkyl;

CN;

CF₃;

HO;

10 (C₁-C₆ alkyl)-O;

(C₁-C₆ alkyl)-S(O)₂;

H₂N;

(C₁-C₆ alkyl)-N(H);

(C₁-C₆ alkyl)₂-N;

15 (C₁-C₆ alkyl)-C(O)O-(C₁-C₈ alkylenyl)_m;

(C₁-C₆ alkyl)-C(O)O-(1- to 8-membered heteroalkylenyl)_m;

(C₁-C₆ alkyl)-C(O)N(H)-(C₁-C₈ alkylenyl)_m;

(C₁-C₆ alkyl)-C(O)N(H)-(1- to 8-membered heteroalkylenyl)_m;

H₂NS(O)₂-(C₁-C₈ alkylenyl);

20 (C₁-C₆ alkyl)-N(H)S(O)₂-(C₁-C₈ alkylenyl)_m;

(C₁-C₆ alkyl)₂-NS(O)₂-(C₁-C₈ alkylenyl)_m;

3- to 6-membered heterocycloalkyl-(G)_m;

Substituted 3- to 6-membered heterocycloalkyl-(G)_m;

5- or 6-membered heteroaryl-(G)_m;

25 Substituted 5- or 6-membered heteroaryl-(G)_m;

(C₁-C₆ alkyl)-S(O)₂-N(H)-C(O)-(C₁-C₈ alkylenyl)_m; and

(C₁-C₆ alkyl)-C(O)-N(H)-S(O)₂-(C₁-C₈ alkylenyl)_m;


wherein each substituent on a carbon atom may further be independently selected from:

30 Halo; and

HO₂C;

wherein 2 substituents may be taken together with a carbon atom to which they are both bonded to form the group C=O;

wherein two adjacent, substantially sp^2 carbon atoms may be taken together with a diradical substituent to form a cyclic diradical selected from:



R is H or C_1 - C_6 alkyl;

G is CH_2 ; O, S, $S(O)$; or $S(O)_2$;

10 m is an integer of 0 or 1;

R^4 is independently selected from the groups:

15 H;
 CH_3 ;
 CH_3O ;
HO;
 CF_3 ; and
CN;

wherein each C_8 - C_{10} bicycloalkyl is a bicyclic carbocyclic ring that contains 8-, 9-, or 10-member carbon atoms which are 5,5-fused, 6,5-fused, or 6,6-fused bicyclic

rings, respectively, and wherein the ring is saturated or optionally contains one carbon-carbon double bond;

wherein each 8- to 10-membered heterobicycloalkyl is a bicyclic ring that contains carbon atoms and from 1 to 4 heteroatoms independently selected from 2 O, 1 S, 1 S(O), 1 S(O)₂, 1 N, 4 N(H), and 4 N(C₁-C₆ alkyl), and wherein when two O atoms or one O atom and one S atom are present, the two O atoms or one O atom and one S atom are not bonded to each other, and wherein the ring is saturated or optionally contains one carbon-carbon or carbon-nitrogen double bond, and wherein the heterobicycloalkyl is a 5,5-fused, 6,5-fused, or 6,6-fused bicyclic ring, respectively,

wherein each heterocycloalkyl is a ring that contains carbon atoms and from 1 to 4 heteroatoms independently selected from 2 O, 1 S, 1 S(O), 1 S(O)₂, 1 N, 4 N(H), and 4 N(C₁-C₆ alkyl), and wherein when two O atoms or one O atom and one S atom are present, the two O atoms or one O atom and one S atom are not bonded to each other, and wherein the ring is saturated or optionally contains one carbon-carbon or carbon-nitrogen double bond;

wherein each 5-membered heteroaryl contains carbon atoms and from 1 to 4 heteroatoms independently selected from 1 O, 1 S, 1 N(H), 1 N(C₁-C₆ alkyl), and 4 N, and each 6-membered heteroaryl contains carbon atoms and 1 or 2 heteroatoms independently selected from N, N(H), and N(C₁-C₆ alkyl), and 5- and 6-membered heteroaryl are monocyclic rings;

wherein each heterobiaryl contains carbon atoms and from 1 to 4 heteroatoms independently selected from 1 O, 1 S, 1 N(H), 1 N(C₁-C₆ alkyl), and 4 N, and where the 8-, 9-, and 10-membered heterobiaryl are 5,5-fused, 6,5-fused, and 6,6-fused bicyclic rings, respectively, and wherein at least 1 of the 2 fused rings of a bicyclic ring is aromatic, and wherein when the O and S atoms both are present, the O and S atoms are not bonded to each other;

wherein with any (C₁-C₆ alkyl)₂-N group, the C₁-C₆ alkyl groups may be optionally taken together with the nitrogen atom to which they are attached to form a 5- or 6-membered heterocycloalkyl; and

wherein each group and each substituent recited above is independently selected.

88. The compound according to Embodiment 87, selected from:
3-Benzyl-1-methyl-1,3,4,5,7,8-hexahydro-2H-pyrido[4,3-d]pyrimidine-6-
carboxylic acid benzyl ester; and
3-Benzyl-1-methyl-1,3,4,5,7,8-hexahydro-2H-pyrido[4,3-d]pyrimidine-6-
carboxylic acid pyridin-4-ylmethyl ester; or
a pharmaceutically acceptable salt thereof.
89. The compound according to Embodiment 87, selected from:
3-Benzyl-1-methyl-1,3,4,5,7,8-hexahydro-2H-pyrido[4,3-d]pyrimidine-6-
carboxylic acid 4-methoxy-benzyl ester;
3-Benzyl-1-methyl-1,3,4,5,7,8-hexahydro-2H-pyrido[4,3-d]pyrimidine-6-
carboxylic acid 3-methoxy-benzyl ester;
3-Benzyl-1-methyl-1,3,4,5,7,8-hexahydro-2H-pyrido[4,3-d]pyrimidine-6-
carboxylic acid 2-methoxy-pyridin-4-ylmethyl ester;
3-Benzyl-1-methyl-1,3,4,5,7,8-hexahydro-2H-pyrido[4,3-d]pyrimidine-6-
carboxylic acid 4-methyl-benzyl ester;
3-Benzyl-1-methyl-1,3,4,5,7,8-hexahydro-2H-pyrido[4,3-d]pyrimidine-6-
carboxylic acid 4-trifluoromethoxy-benzyl ester; and
3-Benzyl-1-methyl-1,3,4,5,7,8-hexahydro-2H-pyrido[4,3-d]pyrimidine-6-
carboxylic acid 4-dimethylamino-benzyl ester; or
a pharmaceutically acceptable salt thereof.
90. The compound according to Embodiment 87, selected from:
3-Benzyl-1-methyl-1,3,4,5,7,8-hexahydro-2H-pyrido[4,3-d]pyrimidine-6-
carboxylic acid pyrrolidin-1-ylmethyl ester;
3-Benzyl-1-methyl-1,3,4,5,7,8-hexahydro-2H-pyrido[4,3-d]pyrimidine-6-
carboxylic acid pyrrol-1-ylmethyl ester;
3-Benzyl-1-methyl-1,3,4,5,7,8-hexahydro-2H-pyrido[4,3-d]pyrimidine-6-
carboxylic acid imidazol-1-ylmethyl ester;
3-Benzyl-1-methyl-1,3,4,5,7,8-hexahydro-2H-pyrido[4,3-d]pyrimidine-6-
carboxylic acid 4-methyl-imidazol-1-ylmethyl ester;
3-Benzyl-1-methyl-1,3,4,5,7,8-hexahydro-2H-pyrido[4,3-d]pyrimidine-6-
carboxylic acid [1,2,4]triazol-4-ylmethyl ester;


3-Benzyl-1-methyl-1,3,4,5,7,8-hexahydro-2H-pyrido[4,3-d]pyrimidine-6-carboxylic acid 2,3-dihydro-tetrazol-1-ylmethyl ester; and
3-Benzyl-1-methyl-1,3,4,5,7,8-hexahydro-2H-pyrido[4,3-d]pyrimidine-6-carboxylic acid 1,3-dihydro-isoindol-2-ylmethyl ester; or
5 a pharmaceutically acceptable salt thereof.

91. The compound according to Embodiment 87, selected from:
3-Benzyl-1-methyl-1,3,4,5,7,8-hexahydro-2H-pyrido[4,3-d]pyrimidine-6-carboxylic acid 1H-indol-2-ylmethyl ester;
10 3-Benzyl-1-methyl-1,3,4,5,7,8-hexahydro-2H-pyrido[4,3-d]pyrimidine-6-carboxylic acid 1-methyl-1H-indol-2-ylmethyl ester;
3-Benzyl-1-methyl-1,3,4,5,7,8-hexahydro-2H-pyrido[4,3-d]pyrimidine-6-carboxylic acid benzo[b]thiophen-2-ylmethyl ester;
3-Benzyl-1-methyl-1,3,4,5,7,8-hexahydro-2H-pyrido[4,3-d]pyrimidine-6-carboxylic acid benzofuran-2-ylmethyl ester;
15 3-Benzyl-1-methyl-1,3,4,5,7,8-hexahydro-2H-pyrido[4,3-d]pyrimidine-6-carboxylic acid benzoxazol-2-ylmethyl ester;
3-Benzyl-1-methyl-1,3,4,5,7,8-hexahydro-2H-pyrido[4,3-d]pyrimidine-6-carboxylic acid 1H-benzimidazol-2-ylmethyl ester; and
20 3-Benzyl-1-methyl-1,3,4,5,7,8-hexahydro-2H-pyrido[4,3-d]pyrimidine-6-carboxylic acid benzothiazol-2-ylmethyl ester; or
a pharmaceutically acceptable salt thereof.

92. The compound according to Embodiment 87, selected from:
25 3-Benzyl-1-methyl-1,3,4,5,7,8-hexahydro-2H-pyrido[4,3-d]pyrimidine-6-carboxylic acid thiazol-2-ylmethyl ester;
3-Benzyl-1-methyl-1,3,4,5,7,8-hexahydro-2H-pyrido[4,3-d]pyrimidine-6-carboxylic acid furan-2-ylmethyl ester;
3-Benzyl-1-methyl-1,3,4,5,7,8-hexahydro-2H-pyrido[4,3-d]pyrimidine-6-carboxylic acid 1-methyl-1H-pyrrol-2-ylmethyl ester;
30 3-Benzyl-1-methyl-1,3,4,5,7,8-hexahydro-2H-pyrido[4,3-d]pyrimidine-6-carboxylic acid thiophen-2-ylmethyl ester;

- 3-Benzyl-1-methyl-1,3,4,5,7,8-hexahydro-2H-pyrido[4,3-d]pyrimidine-6-carboxylic acid 5-phenyl-thiophen-2-ylmethyl ester;
3-Benzyl-1-methyl-1,3,4,5,7,8-hexahydro-2H-pyrido[4,3-d]pyrimidine-6-carboxylic acid 5-methoxy-thiophen-2-ylmethyl ester;
5 3-Benzyl-1-methyl-1,3,4,5,7,8-hexahydro-2H-pyrido[4,3-d]pyrimidine-6-carboxylic acid 5-dimethylaminomethyl-furan-2-ylmethyl ester;
and
3-Benzyl-1-methyl-1,3,4,5,7,8-hexahydro-2H-pyrido[4,3-d]pyrimidine-6-carboxylic acid 5-chloro-thiophen-2-ylmethyl ester; or
10 a pharmaceutically acceptable salt thereof.

93. A compound of Formula IX



IX

- or a pharmaceutically acceptable salt thereof, or a pyrido-N-oxide thereof,
15 wherein:

R¹ is independently selected from:

- C₅ or C₆ cycloalkyl-(C₁-C₈ alkylenyl);
Substituted C₅ or C₆ cycloalkyl-(C₁-C₈ alkylenyl);
C₈-C₁₀ bicycloalkyl-(C₁-C₈ alkylenyl);
20 Substituted C₈-C₁₀ bicycloalkyl-(C₁-C₈ alkylenyl);
5- or 6-membered heterocycloalkyl-(C₁-C₈ alkylenyl);
Substituted 5- or 6-membered heterocycloalkyl-(C₁-C₈ alkylenyl);
8- to 10-membered heterobicycloalkyl-(C₁-C₈ alkylenyl);
Substituted 8- to 10-membered heterobicycloalkyl-(C₁-C₈ alkylenyl);
25 Phenyl-(C₁-C₈ alkylenyl);
Substituted phenyl-(C₁-C₈ alkylenyl);
Naphthyl-(C₁-C₈ alkylenyl);
Substituted naphthyl-(C₁-C₈ alkylenyl);
5- or 6-membered heteroaryl-(C₁-C₈ alkylenyl);

- Substituted 5- or 6-membered heteroaryl-(C₁-C₈ alkylene);
8- to 10-membered heterobiaryl-(C₁-C₈ alkylene);
Substituted 8- to 10-membered heterobiaryl-(C₁-C₈ alkylene);
Phenyl;
5 Substituted phenyl;
Naphthyl;
Substituted naphthyl;
5- or 6-membered heteroaryl;
Substituted 5- or 6-membered heteroaryl;
10 8- to 10-membered heterobiaryl; and
Substituted 8- to 10-membered heterobiaryl;

R² is independently selected from:

- H;
C₁-C₆ alkyl;
15 Phenyl-(C₁-C₈ alkylene);
Substituted phenyl-(C₁-C₈ alkylene);
Naphthyl-(C₁-C₈ alkylene);
Substituted naphthyl-(C₁-C₈ alkylene);
5- or 6-membered heteroaryl-(C₁-C₈ alkylene);
20 Substituted 5- or 6-membered heteroaryl-(C₁-C₈ alkylene);
8- to 10-membered heterobiaryl-(C₁-C₈ alkylene);
Substituted 8- to 10-membered heterobiaryl-(C₁-C₈ alkylene);
Phenyl-O-(C₁-C₈ alkylene);
Substituted phenyl-O-(C₁-C₈ alkylene);
25 Phenyl-S-(C₁-C₈ alkylene);
Substituted phenyl-S-(C₁-C₈ alkylene);
Phenyl-S(O)-(C₁-C₈ alkylene);
Substituted phenyl-S(O)-(C₁-C₈ alkylene);
Phenyl-S(O)₂-(C₁-C₈ alkylene); and
30 Substituted phenyl-S(O)₂-(C₁-C₈ alkylene);

Each substituted R¹ and R² group contains from 1 to 4 substituents, each independently on a carbon or nitrogen atom, independently selected from:

C₁-C₆ alkyl;


- CN;
 CF₃;
 HO;
 (C₁-C₆ alkyl)-O;
 5 (C₁-C₆ alkyl)-S(O)₂;
 H₂N;
 (C₁-C₆ alkyl)-N(H);
 (C₁-C₆ alkyl)₂-N;
 (C₁-C₆ alkyl)-C(O)O-(C₁-C₈ alkylenyl)_m;
 10 (C₁-C₆ alkyl)-C(O)O-(1- to 8-membered heteroalkylenyl)_m;
 (C₁-C₆ alkyl)-C(O)N(H)-(C₁-C₈ alkylenyl)_m;
 (C₁-C₆ alkyl)-C(O)N(H)-(1- to 8-membered heteroalkylenyl)_m;
 H₂NS(O)₂-(C₁-C₈ alkylenyl);
 (C₁-C₆ alkyl)-N(H)S(O)₂-(C₁-C₈ alkylenyl)_m;
 15 (C₁-C₆ alkyl)₂-NS(O)₂-(C₁-C₈ alkylenyl)_m;
 3- to 6-membered heterocycloalkyl-(G)_m;
 Substituted 3- to 6-membered heterocycloalkyl-(G)_m;
 5- or 6-membered heteroaryl-(G)_m;
 Substituted 5- or 6-membered heteroaryl-(G)_m;
 20 (C₁-C₆ alkyl)-S(O)₂-N(H)-C(O)-(C₁-C₈ alkylenyl)_m; and
 (C₁-C₆ alkyl)-C(O)-N(H)-S(O)₂-(C₁-C₈ alkylenyl)_m;


wherein each substituent on a carbon atom may further be independently selected from:

- Halo; and
 25 HO₂C;

wherein 2 substituents may be taken together with a carbon atom to which they are both bonded to form the group C=O;

wherein two adjacent, substantially sp² carbon atoms may be taken together with a diradical substituent to form a cyclic diradical selected from:





5 R is H or C₁-C₆ alkyl;

G is CH₂; O, S, S(O); or S(O)₂;

m is an integer of 0 or 1;

R⁴ is independently selected from the groups:

- 10 H;
CH₃;
CH₃O;
HO;
CF₃; and
CN;

15 wherein each C₈-C₁₀ bicycloalkyl is a bicyclic carbocyclic ring that contains 8-, 9-, or 10-member carbon atoms which are 5,5-fused, 6,5-fused, or 6,6-fused bicyclic rings, respectively, and wherein the ring is saturated or optionally contains one carbon-carbon double bond;

20 wherein each 8- to 10-membered heterobicycloalkyl is a bicyclic ring that contains carbon atoms and from 1 to 4 heteroatoms independently selected from 2 O, 1 S, 1 S(O), 1 S(O)₂, 1 N, 4 N(H), and 4 N(C₁-C₆ alkyl), and wherein when two

O atoms or one O atom and one S atom are present, the two O atoms or one O atom and one S atom are not bonded to each other, and wherein the ring is saturated or optionally contains one carbon-carbon or carbon-nitrogen double bond, and wherein the heterobicycloalkyl is a 5,5-fused, 6,5-fused, or 6,6-fused bicyclic ring, respectively,

wherein each heterocycloalkyl is a ring that contains carbon atoms and from 1 to 4 heteroatoms independently selected from 2 O, 1 S, 1 S(O), 1 S(O)₂, 1 N, 4 N(H), and 4 N(C₁-C₆ alkyl), and wherein when two O atoms or one O atom and one S atom are present, the two O atoms or one O atom and one S atom are not bonded to each other, and wherein the ring is saturated or optionally contains one carbon-carbon or carbon-nitrogen double bond;

wherein each 5-membered heteroaryl contains carbon atoms and from 1 to 4 heteroatoms independently selected from 1 O, 1 S, 1 N(H), 1 N(C₁-C₆ alkyl), and 4 N, and each 6-membered heteroaryl contains carbon atoms and 1 or 2 heteroatoms independently selected from N, N(H), and N(C₁-C₆ alkyl), and 5- and 6-membered heteroaryl are monocyclic rings;

wherein each heterobiaryl contains carbon atoms and from 1 to 4 heteroatoms independently selected from 1 O, 1 S, 1 N(H), 1 N(C₁-C₆ alkyl), and 4 N, and where the 8-, 9-, and 10-membered heterobiaryl are 5,5-fused, 6,5-fused, and 6,6-fused bicyclic rings, respectively, and wherein at least 1 of the 2 fused rings of a bicyclic ring is aromatic, and wherein when the O and S atoms both are present, the O and S atoms are not bonded to each other;

wherein with any (C₁-C₆ alkyl)₂-N group, the C₁-C₆ alkyl groups may be optionally taken together with the nitrogen atom to which they are attached to form a 5- or 6-membered heterocycloalkyl; and

wherein each group and each substituent recited above is independently selected.

94. The compound according to Embodiment 93, selected from:

3-Benzyl-1-methyl-6-(3-phenyl-prop-2-ynyl)-1,2,3,4,5,6,7,8-octahydro-pyrido[4,3-d]pyrimidine; and

3-Benzyl-1-methyl-6-(3-pyridin-4-yl-prop-2-ynyl)-1,2,3,4,5,6,7,8-octahydro-pyrido[4,3-d]pyrimidine; or

a pharmaceutically acceptable salt thereof.

95. The compound according to Embodiment 93, selected from:
3-Benzyl-6-[3-(4-methoxy-phenyl)-prop-2-ynyl]-1-methyl-1,2,3,4,5,6,7,8-
5 octahydro-pyrido[4,3-d]pyrimidine;
3-Benzyl-6-[3-(3-methoxy-phenyl)-prop-2-ynyl]-1-methyl-1,2,3,4,5,6,7,8-
octahydro-pyrido[4,3-d]pyrimidine;
3-Benzyl-6-[3-(2-methoxy-pyridin-4-yl)-prop-2-ynyl]-1-methyl-
1,2,3,4,5,6,7,8-octahydro-pyrido[4,3-d]pyrimidine; and
10 4-[3-(3-Benzyl-1-methyl-1,3,4,5,7,8-hexahydro-2H-pyrido[4,3-
d]pyrimidin-6-yl)-prop-1-ynyl]-pyridin-1-ol; or
a pharmaceutically acceptable salt thereof.
96. The compound according to Embodiment 93, selected from:
15 3-Benzyl-1-methyl-6-(3-pyrrolidin-1-yl-prop-2-ynyl)-1,2,3,4,5,6,7,8-
octahydro-pyrido[4,3-d]pyrimidine;
3-Benzyl-1-methyl-6-(3-piperidin-1-yl-prop-2-ynyl)-1,2,3,4,5,6,7,8-
octahydro-pyrido[4,3-d]pyrimidine; and
3-Benzyl-1-methyl-6-[3-(4-methyl-piperazin-1-yl)-prop-2-ynyl]-
20 1,2,3,4,5,6,7,8-octahydro-pyrido[4,3-d]pyrimidine; or
a pharmaceutically acceptable salt thereof.
97. The compound according to Embodiment 93, selected from:
3-Benzyl-1-methyl-6-[3-(1H-pyrrol-2-yl)-prop-2-ynyl]-1,2,3,4,5,6,7,8-
25 octahydro-pyrido[4,3-d]pyrimidine;
3-Benzyl-1-methyl-6-(3-thiophen-2-yl-prop-2-ynyl)-1,2,3,4,5,6,7,8-
octahydro-pyrido[4,3-d]pyrimidine;
3-Benzyl-6-(3-furan-2-yl-prop-2-ynyl)-1-methyl-1,2,3,4,5,6,7,8-
octahydro-pyrido[4,3-d]pyrimidine;
30 3-Benzyl-1-methyl-6-(3-oxazol-2-yl-prop-2-ynyl)-1,2,3,4,5,6,7,8-
octahydro-pyrido[4,3-d]pyrimidine;
3-Benzyl-1-methyl-6-(3-thiazol-2-yl-prop-2-ynyl)-1,2,3,4,5,6,7,8-
octahydro-pyrido[4,3-d]pyrimidine;

3-Benzyl-1-methyl-6-[3-(5-phenyl-thiazol-2-yl)-prop-2-ynyl]-
1,2,3,4,5,6,7,8-octahydro-pyrido[4,3-d]pyrimidine; and
3-Benzyl-6-{3-[5-(4-methoxy-phenyl)-thiazol-2-yl]-prop-2-ynyl}-1-
methyl-1,2,3,4,5,6,7,8-octahydro-pyrido[4,3-d]pyrimidine; or
5 a pharmaceutically acceptable salt thereof.

98. The compound according to Embodiment 93, selected from:

3-Benzyl-6-[3-(1,3-dihydro-isoindol-2-yl)-prop-2-ynyl]-1-methyl-
1,2,3,4,5,6,7,8-octahydro-pyrido[4,3-d]pyrimidine;

10 3-Benzyl-6-(3-imidazol-1-yl-prop-2-ynyl)-1-methyl-1,2,3,4,5,6,7,8-
octahydro-pyrido[4,3-d]pyrimidine;

3-Benzyl-1-methyl-6-(3-[1,2,4]triazol-4-yl-prop-2-ynyl)-1,2,3,4,5,6,7,8-
octahydro-pyrido[4,3-d]pyrimidine;

15 3-Benzyl-1-methyl-6-(3-tetrazol-1-yl-prop-2-ynyl)-1,2,3,4,5,6,7,8-
octahydro-pyrido[4,3-d]pyrimidine;

a pharmaceutically acceptable salt thereof.

99. The compound according to Embodiment 93, selected from:

3-Benzyl-6-[3-(1H-indol-2-yl)-prop-2-ynyl]-1-methyl-1,2,3,4,5,6,7,8-
20 octahydro-pyrido[4,3-d]pyrimidine;

6-(3-Benzo[b]thiophen-2-yl-prop-2-ynyl)-3-benzyl-1-methyl-
1,2,3,4,5,6,7,8-octahydro-pyrido[4,3-d]pyrimidine;

6-(3-Benzo[b]thiophen-2-yl-prop-2-ynyl)-3-benzyl-1-methyl-
1,2,3,4,5,6,7,8-octahydro-pyrido[4,3-d]pyrimidine;

25 6-(3-Benzofuran-2-yl-prop-2-ynyl)-3-benzyl-1-methyl-1,2,3,4,5,6,7,8-
octahydro-pyrido[4,3-d]pyrimidine;

6-(3-Benzooxazol-2-yl-prop-2-ynyl)-3-benzyl-1-methyl-1,2,3,4,5,6,7,8-
octahydro-pyrido[4,3-d]pyrimidine; and

30 6-[3-(1H-Benzoimidazol-2-yl)-prop-2-ynyl]-3-benzyl-1-methyl-
1,2,3,4,5,6,7,8-octahydro-pyrido[4,3-d]pyrimidine; or

a pharmaceutically acceptable salt thereof.

100. The compound according to Embodiment 93, selected from:

- 3-Benzyl-1-methyl-6-[3-(4-trifluoromethoxy-phenyl)-prop-2-ynyl]-
1,2,3,4,5,6,7,8-octahydro-pyrido[4,3-d]pyrimidine;
- 3-Benzyl-6-[3-(4-fluoro-phenyl)-prop-2-ynyl]-1-methyl-1,2,3,4,5,6,7,8-
octahydro-pyrido[4,3-d]pyrimidine;
- 5 3-Benzyl-6-[3-(4-chloro-phenyl)-prop-2-ynyl]-1-methyl-1,2,3,4,5,6,7,8-
octahydro-pyrido[4,3-d]pyrimidine;
- 3-Benzyl-6-[3-(4-bromo-phenyl)-prop-2-ynyl]-1-methyl-1,2,3,4,5,6,7,8-
octahydro-pyrido[4,3-d]pyrimidine;
- 10 3-Benzyl-6-[3-(4-iodo-phenyl)-prop-2-ynyl]-1-methyl-1,2,3,4,5,6,7,8-
octahydro-pyrido[4,3-d]pyrimidine;
- 3-Benzyl-1-methyl-6-[3-(4-trifluoromethyl-phenyl)-prop-2-ynyl]-
1,2,3,4,5,6,7,8-octahydro-pyrido[4,3-d]pyrimidine;
- 3-Benzyl-1-methyl-6-[3-(4-trichloromethyl-phenyl)-prop-2-ynyl]-
1,2,3,4,5,6,7,8-octahydro-pyrido[4,3-d]pyrimidine;
- 15 3-Benzyl-6-[3-(3,4-dimethoxy-phenyl)-prop-2-ynyl]-1-methyl-
1,2,3,4,5,6,7,8-octahydro-pyrido[4,3-d]pyrimidine; and
{ 4-[3-(3-Benzyl-1-methyl-1,3,4,5,7,8-hexahydro-2H-pyrido[4,3-
d]pyrimidin-6-yl)-prop-1-ynyl]-phenyl }-dimethyl-amine; or
a pharmaceutically acceptable salt thereof.
- 20
101. The compound according to Embodiment 93, selected from:
- 1-Methyl-3-(4-methyl-benzyl)-6-(3-phenyl-prop-2-ynyl)-1,2,3,4,5,6,7,8-
octahydro-pyrido[4,3-d]pyrimidine;
- 1-{ 4-[1-Methyl-6-(3-phenyl-prop-2-ynyl)-1,4,5,6,7,8-hexahydro-2H-
pyrido[4,3-d]pyrimidin-3-ylmethyl]-phenyl }-ethanone;
- 25 4-[1-Methyl-6-(3-phenyl-prop-2-ynyl)-1,4,5,6,7,8-hexahydro-2H-
pyrido[4,3-d]pyrimidin-3-ylmethyl]-benzoic acid;
- 4-[1-Methyl-6-(3-phenyl-prop-2-ynyl)-1,4,5,6,7,8-hexahydro-2H-
pyrido[4,3-d]pyrimidin-3-ylmethyl]-benzoic acid methyl ester;
- 30 3-(4-Fluoro-benzyl)-1-methyl-6-(3-phenyl-prop-2-ynyl)-1,2,3,4,5,6,7,8-
octahydro-pyrido[4,3-d]pyrimidine;
- 3-(4-Chloro-benzyl)-1-methyl-6-(3-phenyl-prop-2-ynyl)-1,2,3,4,5,6,7,8-
octahydro-pyrido[4,3-d]pyrimidine;


- 3-(4-Bromo-benzyl)-1-methyl-6-(3-phenyl-prop-2-ynyl)-1,2,3,4,5,6,7,8-octahydro-pyrido[4,3-d]pyrimidine;
- 3-(4-Iodo-benzyl)-1-methyl-6-(3-phenyl-prop-2-ynyl)-1,2,3,4,5,6,7,8-octahydro-pyrido[4,3-d]pyrimidine;
- 5 1-Methyl-6-(3-phenyl-prop-2-ynyl)-3-(4-trichloromethyl-benzyl)-1,2,3,4,5,6,7,8-octahydro-pyrido[4,3-d]pyrimidine;
- 1-Methyl-6-(3-phenyl-prop-2-ynyl)-3-(4-trifluoromethyl-benzyl)-1,2,3,4,5,6,7,8-octahydro-pyrido[4,3-d]pyrimidine;
- 3-(4-Methoxy-benzyl)-1-methyl-6-(3-phenyl-prop-2-ynyl)-1,2,3,4,5,6,7,8-octahydro-pyrido[4,3-d]pyrimidine; and
- 10 1-Methyl-6-(3-phenyl-prop-2-ynyl)-3-(4-trifluoromethoxy-benzyl)-1,2,3,4,5,6,7,8-octahydro-pyrido[4,3-d]pyrimidine; or
a pharmaceutically acceptable salt thereof.
- 15 102. The compound according to Embodiment 93, selected from:
- 1-Methyl-6-(3-phenyl-prop-2-ynyl)-3-pyridin-4-ylmethyl-1,2,3,4,5,6,7,8-octahydro-pyrido[4,3-d]pyrimidine;
- 1-Methyl-6-(3-phenyl-prop-2-ynyl)-3-pyridin-3-ylmethyl-1,2,3,4,5,6,7,8-octahydro-pyrido[4,3-d]pyrimidine;
- 20 3-(2-Methoxy-pyridin-4-ylmethyl)-1-methyl-6-(3-phenyl-prop-2-ynyl)-1,2,3,4,5,6,7,8-octahydro-pyrido[4,3-d]pyrimidine; and
- 3-Cyclohexylmethyl-1-methyl-6-(3-phenyl-prop-2-ynyl)-1,2,3,4,5,6,7,8-octahydro-pyrido[4,3-d]pyrimidine; or
a pharmaceutically acceptable salt thereof.
- 25 103. The compound according to Embodiment 93, selected from:
- 1-Methyl-6-(3-phenyl-prop-2-ynyl)-3-piperidin-1-ylmethyl-1,2,3,4,5,6,7,8-octahydro-pyrido[4,3-d]pyrimidine
- 1-Methyl-3-(4-methyl-piperazin-1-ylmethyl)-6-(3-phenyl-prop-2-ynyl)-1,2,3,4,5,6,7,8-octahydro-pyrido[4,3-d]pyrimidine
- 30 1-Methyl-6-(3-phenyl-prop-2-ynyl)-3-pyrrolidin-1-ylmethyl-1,2,3,4,5,6,7,8-octahydro-pyrido[4,3-d]pyrimidine

- 1-Methyl-6-(3-phenyl-prop-2-ynyl)-3-pyrrol-1-ylmethyl-1,2,3,4,5,6,7,8-octahydro-pyrido[4,3-d]pyrimidine
- 3-Imidazol-1-ylmethyl-1-methyl-6-(3-phenyl-prop-2-ynyl)-1,2,3,4,5,6,7,8-octahydro-pyrido[4,3-d]pyrimidine
- 5 1-Methyl-6-(3-phenyl-prop-2-ynyl)-3-[1,2,4]triazol-4-ylmethyl-1,2,3,4,5,6,7,8-octahydro-pyrido[4,3-d]pyrimidine
- 1-Methyl-6-(3-phenyl-prop-2-ynyl)-3-tetrazol-1-ylmethyl-1,2,3,4,5,6,7,8-octahydro-pyrido[4,3-d]pyrimidine; and
- 3-(1,3-Dihydro-isoindol-2-ylmethyl)-1-methyl-6-(3-phenyl-prop-2-ynyl)-1,2,3,4,5,6,7,8-octahydro-pyrido[4,3-d]pyrimidine; or
- 10 a pharmaceutically acceptable salt thereof.
104. The compound according to Embodiment 93, selected from:
- 1-Methyl-6-(3-phenyl-prop-2-ynyl)-3-(1H-pyrrol-2-ylmethyl)-1,2,3,4,5,6,7,8-octahydro-pyrido[4,3-d]pyrimidine;
- 15 1-Methyl-3-(1-methyl-1H-pyrrol-2-ylmethyl)-6-(3-phenyl-prop-2-ynyl)-1,2,3,4,5,6,7,8-octahydro-pyrido[4,3-d]pyrimidine;
- 1-Methyl-3-(1-methyl-1H-imidazol-2-ylmethyl)-6-(3-phenyl-prop-2-ynyl)-1,2,3,4,5,6,7,8-octahydro-pyrido[4,3-d]pyrimidine;
- 20 1-Methyl-3-oxazol-2-ylmethyl-6-(3-phenyl-prop-2-ynyl)-1,2,3,4,5,6,7,8-octahydro-pyrido[4,3-d]pyrimidine;
- 1-Methyl-6-(3-phenyl-prop-2-ynyl)-3-thiazol-2-ylmethyl-1,2,3,4,5,6,7,8-octahydro-pyrido[4,3-d]pyrimidine;
- 1-Methyl-6-(3-phenyl-prop-2-ynyl)-3-thiophen-2-ylmethyl-1,2,3,4,5,6,7,8-octahydro-pyrido[4,3-d]pyrimidine;
- 25 3-Furan-2-ylmethyl-1-methyl-6-(3-phenyl-prop-2-ynyl)-1,2,3,4,5,6,7,8-octahydro-pyrido[4,3-d]pyrimidine;
- 3-(5-Methoxy-furan-2-ylmethyl)-1-methyl-6-(3-phenyl-prop-2-ynyl)-1,2,3,4,5,6,7,8-octahydro-pyrido[4,3-d]pyrimidine; and
- 30 1-Methyl-3-(5-phenyl-furan-2-ylmethyl)-6-(3-phenyl-prop-2-ynyl)-1,2,3,4,5,6,7,8-octahydro-pyrido[4,3-d]pyrimidine; or
- a pharmaceutically acceptable salt thereof.

105. The compound according to Embodiment 93, selected from:
1-Methyl-3-(1-methyl-1H-benzimidazol-2-ylmethyl)-6-(3-phenyl-prop-2-ynyl)-1,2,3,4,5,6,7,8-octahydro-pyrido[4,3-d]pyrimidine;
3-Benzothiazol-2-ylmethyl-1-methyl-6-(3-phenyl-prop-2-ynyl)-
5 1,2,3,4,5,6,7,8-octahydro-pyrido[4,3-d]pyrimidine;
3-Benzo[b]thiophen-2-ylmethyl-1-methyl-6-(3-phenyl-prop-2-ynyl)-
1,2,3,4,5,6,7,8-octahydro-pyrido[4,3-d]pyrimidine;
3-Benzofuran-2-ylmethyl-1-methyl-6-(3-phenyl-prop-2-ynyl)-
1,2,3,4,5,6,7,8-octahydro-pyrido[4,3-d]pyrimidine; and
10 3-Benzooxazol-2-ylmethyl-1-methyl-6-(3-phenyl-prop-2-ynyl)-
1,2,3,4,5,6,7,8-octahydro-pyrido[4,3-d]pyrimidine; or
a pharmaceutically acceptable salt thereof.
106. The compound according to Embodiment 93, selected from:
15 3-(4-Methanesulfonyl-benzyl)-1-methyl-6-(3-phenyl-prop-2-ynyl)-
1,2,3,4,5,6,7,8-octahydro-pyrido[4,3-d]pyrimidine;
4-[1-Methyl-6-(3-phenyl-prop-2-ynyl)-1,4,5,6,7,8-hexahydro-2H-
pyrido[4,3-d]pyrimidin-3-ylmethyl]-benzenesulfonic acid;
4-[1-Methyl-6-(3-phenyl-prop-2-ynyl)-1,4,5,6,7,8-hexahydro-2H-
20 pyrido[4,3-d]pyrimidin-3-ylmethyl]-benzenesulfonamide;
N,N-Dimethyl-4-[1-methyl-6-(3-phenyl-prop-2-ynyl)-1,4,5,6,7,8-
hexahydro-2H-pyrido[4,3-d]pyrimidin-3-ylmethyl]-
benzenesulfonamide;
6-(3-phenyl-prop-2-ynyl)-1,2,3,4,5,6,7,8-octahydro-pyrido[4,3-
25 d]pyrimidine;
3-Benzo[b]thiophen-5-ylmethyl-1-methyl-6-(3-phenyl-prop-2-ynyl)-
1,2,3,4,5,6,7,8-octahydro-pyrido[4,3-d]pyrimidine;
3-Benzofuran-5-ylmethyl-1-methyl-6-(3-phenyl-prop-2-ynyl)-
1,2,3,4,5,6,7,8-octahydro-pyrido[4,3-d]pyrimidine;
30 Dimethyl-{4-[1-methyl-6-(3-phenyl-prop-2-ynyl)-1,4,5,6,7,8-hexahydro-
2H-pyrido[4,3-d]pyrimidin-3-ylmethyl]-phenyl}-amine;
3-(4-Aziridin-1-yl-benzyl)-1-methyl-6-(3-phenyl-prop-2-ynyl)-
1,2,3,4,5,6,7,8-octahydro-pyrido[4,3-d]pyrimidine; and

3-(Naphth-2-yl)-1-methyl-6-(3-phenyl-prop-2-ynyl)-1,2,3,4,5,6,7,8-octahydro-pyrido[4,3-d]pyrimidine; or
a pharmaceutically acceptable salt thereof.

5 107. A compound of Formula X



X

or a pharmaceutically acceptable salt thereof, or a pyrido-N-oxide thereof,
wherein:

R¹ is independently selected from:

- 10 C₅ or C₆ cycloalkyl-(C₁-C₈ alkylenyl);
Substituted C₅ or C₆ cycloalkyl-(C₁-C₈ alkylenyl);
C₈-C₁₀ bicycloalkyl-(C₁-C₈ alkylenyl);
Substituted C₈-C₁₀ bicycloalkyl-(C₁-C₈ alkylenyl);
5- or 6-membered heterocycloalkyl-(C₁-C₈ alkylenyl);
15 Substituted 5- or 6-membered heterocycloalkyl-(C₁-C₈ alkylenyl);
8- to 10-membered heterobicycloalkyl-(C₁-C₈ alkylenyl);
Substituted 8- to 10-membered heterobicycloalkyl-(C₁-C₈ alkylenyl);
Phenyl-(C₁-C₈ alkylenyl);
Substituted phenyl-(C₁-C₈ alkylenyl);
20 Naphthyl-(C₁-C₈ alkylenyl);
Substituted naphthyl-(C₁-C₈ alkylenyl);
5- or 6-membered heteroaryl-(C₁-C₈ alkylenyl);
Substituted 5- or 6-membered heteroaryl-(C₁-C₈ alkylenyl);
8- to 10-membered heterobiaryl-(C₁-C₈ alkylenyl);
25 Substituted 8- to 10-membered heterobiaryl-(C₁-C₈ alkylenyl);
Phenyl;
Substituted phenyl;
Naphthyl;
Substituted naphthyl;
30 5- or 6-membered heteroaryl;

Substituted 5- or 6-membered heteroaryl;

8- to 10-membered heterobiaryl; and

Substituted 8- to 10-membered heterobiaryl;


R^2 is independently selected from:


- 5 H;
 C_1 - C_6 alkyl;
 Phenyl- $(C_1$ - C_8 alkylenyl);
 Substituted phenyl- $(C_1$ - C_8 alkylenyl);
 Naphthyl- $(C_1$ - C_8 alkylenyl);
10 Substituted naphthyl- $(C_1$ - C_8 alkylenyl);
 5- or 6-membered heteroaryl- $(C_1$ - C_8 alkylenyl);
 Substituted 5- or 6-membered heteroaryl- $(C_1$ - C_8 alkylenyl);
 8- to 10-membered heterobiaryl- $(C_1$ - C_8 alkylenyl);
 Substituted 8- to 10-membered heterobiaryl- $(C_1$ - C_8 alkylenyl);
15 Phenyl-O- $(C_1$ - C_8 alkylenyl);
 Substituted phenyl-O- $(C_1$ - C_8 alkylenyl);
 Phenyl-S- $(C_1$ - C_8 alkylenyl);
 Substituted phenyl-S- $(C_1$ - C_8 alkylenyl);
 Phenyl-S(O)- $(C_1$ - C_8 alkylenyl);
20 Substituted phenyl-S(O)- $(C_1$ - C_8 alkylenyl);
 Phenyl-S(O)₂- $(C_1$ - C_8 alkylenyl); and
 Substituted phenyl-S(O)₂- $(C_1$ - C_8 alkylenyl);

Each substituted R^1 and R^2 group contains from 1 to 4 substituents, each independently on a carbon or nitrogen atom, independently selected from:

- 25 C_1 - C_6 alkyl;
 CN;
 CF₃;
 HO;
 $(C_1$ - C_6 alkyl)-O;
30 $(C_1$ - C_6 alkyl)-S(O)₂;
 H₂N;
 $(C_1$ - C_6 alkyl)-N(H);
 $(C_1$ - C_6 alkyl)₂-N;

- $(C_1-C_6 \text{ alkyl})-C(O)O-(C_1-C_8 \text{ alkylenyl})_m$;
 $(C_1-C_6 \text{ alkyl})-C(O)O-(1- \text{ to } 8\text{-membered heteroalkylenyl})_m$;
 $(C_1-C_6 \text{ alkyl})-C(O)N(H)-(C_1-C_8 \text{ alkylenyl})_m$;
 $(C_1-C_6 \text{ alkyl})-C(O)N(H)-(1- \text{ to } 8\text{-membered heteroalkylenyl})_m$;
5 $H_2NS(O)_2-(C_1-C_8 \text{ alkylenyl})$;
 $(C_1-C_6 \text{ alkyl})-N(H)S(O)_2-(C_1-C_8 \text{ alkylenyl})_m$;
 $(C_1-C_6 \text{ alkyl})_2-NS(O)_2-(C_1-C_8 \text{ alkylenyl})_m$;
3- to 6-membered heterocycloalkyl-(G)_m;
Substituted 3- to 6-membered heterocycloalkyl-(G)_m;
10 5- or 6-membered heteroaryl-(G)_m;
Substituted 5- or 6-membered heteroaryl-(G)_m;
 $(C_1-C_6 \text{ alkyl})-S(O)_2-N(H)-C(O)-(C_1-C_8 \text{ alkylenyl})_m$; and
 $(C_1-C_6 \text{ alkyl})-C(O)-N(H)-S(O)_2-(C_1-C_8 \text{ alkylenyl})_m$;
wherein each substituent on a carbon atom may further be independently selected
15 from:
Halo; and
HO₂C;
wherein 2 substituents may be taken together with a carbon atom to which they
are both bonded to form the group C=O;
20 wherein two adjacent, substantially sp² carbon atoms may be taken together with a
diradical substituent to form a cyclic diradical selected from:





R is H or C₁-C₆ alkyl;

G is CH₂; O, S, S(O); or S(O)₂;

5 m is an integer of 0 or 1;

wherein each C₈-C₁₀ bicycloalkyl is a bicyclic carbocyclic ring that contains 8-, 9-, or 10-member carbon atoms which are 5,5-fused, 6,5-fused, or 6,6-fused bicyclic rings, respectively, and wherein the ring is saturated or optionally contains one carbon-carbon double bond;

10 wherein each 8- to 10-membered heterobicycloalkyl is a bicyclic ring that contains carbon atoms and from 1 to 4 heteroatoms independently selected from 2 O, 1 S, 1 S(O), 1 S(O)₂, 1 N, 4 N(H), and 4 N(C₁-C₆ alkyl), and wherein when two O atoms or one O atom and one S atom are present, the two O atoms or one O atom and one S atom are not bonded to each other, and wherein the ring is
15 saturated or optionally contains one carbon-carbon or carbon-nitrogen double bond, and wherein the heterobicycloalkyl is a 5,5-fused, 6,5-fused, or 6,6-fused bicyclic ring, respectively,

wherein each heterocycloalkyl is a ring that contains carbon atoms and from 1 to 4
heteroatoms independently selected from 2 O, 1 S, 1 S(O), 1 S(O)₂, 1 N, 4
20 N(H), and 4 N(C₁-C₆ alkyl), and wherein when two O atoms or one O atom and one S atom are present, the two O atoms or one O atom and one S atom are not bonded to each other, and wherein the ring is saturated or optionally contains one carbon-carbon or carbon-nitrogen double bond;

wherein each 5-membered heteroaryl contains carbon atoms and from 1 to 4
25 heteroatoms independently selected from 1 O, 1 S, 1 N(H), 1 N(C₁-C₆ alkyl), and 4 N, and each 6-membered heteroaryl contains carbon atoms

and 1 or 2 heteroatoms independently selected from N, N(H), and N(C₁-C₆ alkyl), and 5- and 6-membered heteroaryl are monocyclic rings;

wherein each heterobiaryl contains carbon atoms and from 1 to 4 heteroatoms independently selected from 1 O, 1 S, 1 N(H), 1 N(C₁-C₆ alkyl), and 4 N, and where the 8-, 9-, and 10-membered heterobiaryl are 5,5-fused, 6,5-fused, and 6,6-fused bicyclic rings, respectively, and wherein at least 1 of the 2 fused rings of a bicyclic ring is aromatic, and wherein when the O and S atoms both are present, the O and S atoms are not bonded to each other;

wherein with any (C₁-C₆ alkyl)₂-N group, the C₁-C₆ alkyl groups may be optionally taken together with the nitrogen atom to which they are attached to form a 5- or 6-membered heterocycloalkyl; and

wherein each group and each substituent recited above is independently selected.

108. The compound according to Embodiment 107, selected from:

3-Benzyl-6-(3-phenyl-prop-2-ynyl)-3,4,5,6,7,8-hexahydro-pyrido[4,3-d]pyrimidine; and

3-Benzyl-6-(3-pyridin-4-yl-prop-2-ynyl)-3,4,5,6,7,8-hexahydro-pyrido[4,3-d]pyrimidine; or

a pharmaceutically acceptable salt thereof.

109. The compound according to Embodiment 107, selected from:

3-Benzyl-6-[3-(2-methoxy-pyridin-4-yl)-prop-2-ynyl]-3,4,5,6,7,8-hexahydro-pyrido[4,3-d]pyrimidine;

3-Benzyl-6-[3-(3-methoxy-phenyl)-prop-2-ynyl]-3,4,5,6,7,8-hexahydro-pyrido[4,3-d]pyrimidine;

3-Benzyl-6-[3-(4-methoxy-phenyl)-prop-2-ynyl]-3,4,5,6,7,8-hexahydro-pyrido[4,3-d]pyrimidine;

3-Benzyl-6-[3-(4-trifluoromethoxy-phenyl)-prop-2-ynyl]-3,4,5,6,7,8-hexahydro-pyrido[4,3-d]pyrimidine;

3-Benzyl-6-[3-(4-trichloromethoxy-phenyl)-prop-2-ynyl]-3,4,5,6,7,8-hexahydro-pyrido[4,3-d]pyrimidine;

- 3-Benzyl-6-[3-(4-fluoro-phenyl)-prop-2-ynyl]-3,4,5,6,7,8-hexahydro-pyrido[4,3-d]pyrimidine;
- 3-Benzyl-6-[3-(4-chloro-phenyl)-prop-2-ynyl]-3,4,5,6,7,8-hexahydro-pyrido[4,3-d]pyrimidine;
- 5 3-Benzyl-6-[3-(4-bromo-phenyl)-prop-2-ynyl]-3,4,5,6,7,8-hexahydro-pyrido[4,3-d]pyrimidine;
- 3-Benzyl-6-[3-(4-iodo-phenyl)-prop-2-ynyl]-3,4,5,6,7,8-hexahydro-pyrido[4,3-d]pyrimidine;
- 10 3-Benzyl-6-(3-cyclohexyl-prop-2-ynyl)-3,4,5,6,7,8-hexahydro-pyrido[4,3-d]pyrimidine; and
- 3-Benzyl-6-(4,4-dimethyl-pent-2-ynyl)-3,4,5,6,7,8-hexahydro-pyrido[4,3-d]pyrimidine; or
- a pharmaceutically acceptable salt thereof.
- 15 110. The compound according to Embodiment 107, selected from:
- 3-Benzyl-6-(3-piperidin-1-yl-prop-2-ynyl)-3,4,5,6,7,8-hexahydro-pyrido[4,3-d]pyrimidine;
- 3-Benzyl-6-[3-(4-methyl-piperazin-1-yl)-prop-2-ynyl]-3,4,5,6,7,8-hexahydro-pyrido[4,3-d]pyrimidine;
- 20 3-Benzyl-6-(3-morpholin-4-yl-prop-2-ynyl)-3,4,5,6,7,8-hexahydro-pyrido[4,3-d]pyrimidine;
- 3-Benzyl-6-(3-pyrrolidin-1-yl-prop-2-ynyl)-3,4,5,6,7,8-hexahydro-pyrido[4,3-d]pyrimidine;
- 25 3-Benzyl-6-(3-pyrrol-1-yl-prop-2-ynyl)-3,4,5,6,7,8-hexahydro-pyrido[4,3-d]pyrimidine;
- 3-Benzyl-6-(3-imidazol-1-yl-prop-2-ynyl)-3,4,5,6,7,8-hexahydro-pyrido[4,3-d]pyrimidine;
- 3-Benzyl-6-[3-(4-methyl-imidazol-1-yl)-prop-2-ynyl]-3,4,5,6,7,8-hexahydro-pyrido[4,3-d]pyrimidine;
- 30 3-Benzyl-6-(3-[1,2,4]triazol-4-yl-prop-2-ynyl)-3,4,5,6,7,8-hexahydro-pyrido[4,3-d]pyrimidine;
- 3-Benzyl-6-(3-tetrazol-1-yl-prop-2-ynyl)-3,4,5,6,7,8-hexahydro-pyrido[4,3-d]pyrimidine; and

[3-(3-Benzyl-3,5,7,8-tetrahydro-4H-pyrido[4,3-d]pyrimidin-6-yl)-prop-1-ynyl]-dimethyl-amine; or
a pharmaceutically acceptable salt thereof.

- 5 111. The compound according to Embodiment 107, selected from:
3-Benzyl-6-[3-(1H-pyrrol-2-yl)-prop-2-ynyl]-3,4,5,6,7,8-hexahydro-
pyrido[4,3-d]pyrimidine;
3-Benzyl-6-[3-(1H-imidazol-2-yl)-prop-2-ynyl]-3,4,5,6,7,8-hexahydro-
pyrido[4,3-d]pyrimidine;
10 3-Benzyl-6-[3-(1-methyl-1H-imidazol-2-yl)-prop-2-ynyl]-3,4,5,6,7,8-
hexahydro-pyrido[4,3-d]pyrimidine;
3-Benzyl-6-(3-thiazol-2-yl-prop-2-ynyl)-3,4,5,6,7,8-hexahydro-pyrido[4,3-
d]pyrimidine;
3-Benzyl-6-(3-oxazol-2-yl-prop-2-ynyl)-3,4,5,6,7,8-hexahydro-pyrido[4,3-
15 d]pyrimidine;
3-Benzyl-6-[3-(1H-tetrazol-5-yl)-prop-2-ynyl]-3,4,5,6,7,8-hexahydro-
pyrido[4,3-d]pyrimidine;
3-Benzyl-6-(3-thiophen-2-yl-prop-2-ynyl)-3,4,5,6,7,8-hexahydro-
pyrido[4,3-d]pyrimidine;
20 3-Benzyl-6-(3-furan-2-yl-prop-2-ynyl)-3,4,5,6,7,8-hexahydro-pyrido[4,3-
d]pyrimidine;
3-Benzyl-6-[3-(5-methyl-furan-2-yl)-prop-2-ynyl]-3,4,5,6,7,8-hexahydro-
pyrido[4,3-d]pyrimidine; and
3-Benzyl-6-[3-(5-phenyl-furan-2-yl)-prop-2-ynyl]-3,4,5,6,7,8-hexahydro-
25 pyrido[4,3-d]pyrimidine; or
a pharmaceutically acceptable salt thereof.

112. The compound according to Embodiment 107, selected from:
6-(3-Benzofuran-2-yl-prop-2-ynyl)-3-benzyl-3,4,5,6,7,8-hexahydro-
30 pyrido[4,3-d]pyrimidine;
6-(3-Benzooxazol-2-yl-prop-2-ynyl)-3-benzyl-3,4,5,6,7,8-hexahydro-
pyrido[4,3-d]pyrimidine;

6-(3-Benzothiazol-2-yl-prop-2-ynyl)-3-benzyl-3,4,5,6,7,8-hexahydro-pyrido[4,3-d]pyrimidine;
6-(3-Benzo[b]thiophen-2-yl-prop-2-ynyl)-3-benzyl-3,4,5,6,7,8-hexahydro-pyrido[4,3-d]pyrimidine; and
3-Benzyl-6-[3-(1H-indol-2-yl)-prop-2-ynyl]-3,4,5,6,7,8-hexahydro-pyrido[4,3-d]pyrimidine; or
a pharmaceutically acceptable salt thereof.

113. The compound according to Embodiment 107, selected from:

6-[3-(1H-Benzoimidazol-2-yl)-prop-2-ynyl]-3-benzyl-3,4,5,6,7,8-hexahydro-pyrido[4,3-d]pyrimidine;
6-[3-(1H-Benzoimidazol-2-yl)-prop-2-ynyl]-3-(4-methyl-benzyl)-3,4,5,6,7,8-hexahydro-pyrido[4,3-d]pyrimidine;
6-[3-(1H-Benzoimidazol-2-yl)-prop-2-ynyl]-3-pentafluorophenylmethyl-3,4,5,6,7,8-hexahydro-pyrido[4,3-d]pyrimidine;
6-[3-(1H-Benzoimidazol-2-yl)-prop-2-ynyl]-3-(4-fluoro-benzyl)-3,4,5,6,7,8-hexahydro-pyrido[4,3-d]pyrimidine;
6-[3-(1H-Benzoimidazol-2-yl)-prop-2-ynyl]-3-(4-chloro-benzyl)-3,4,5,6,7,8-hexahydro-pyrido[4,3-d]pyrimidine;
6-[3-(1H-Benzoimidazol-2-yl)-prop-2-ynyl]-3-(4-bromo-benzyl)-3,4,5,6,7,8-hexahydro-pyrido[4,3-d]pyrimidine;
6-[3-(1H-Benzoimidazol-2-yl)-prop-2-ynyl]-3-(4-iodo-benzyl)-3,4,5,6,7,8-hexahydro-pyrido[4,3-d]pyrimidine;
(4-{6-[3-(1H-Benzoimidazol-2-yl)-prop-2-ynyl]-5,6,7,8-tetrahydro-4H-pyrido[4,3-d]pyrimidin-3-ylmethyl}-phenyl)-dimethyl-amine;
6-[3-(1H-Benzoimidazol-2-yl)-prop-2-ynyl]-3-(4-pyrrolidin-1-yl-benzyl)-3,4,5,6,7,8-hexahydro-pyrido[4,3-d]pyrimidine;
6-[3-(1H-Benzoimidazol-2-yl)-prop-2-ynyl]-3-(4-piperidin-1-yl-benzyl)-3,4,5,6,7,8-hexahydro-pyrido[4,3-d]pyrimidine;
6-[3-(1H-Benzoimidazol-2-yl)-prop-2-ynyl]-3-[4-(4-methyl-piperazin-1-yl)-benzyl]-3,4,5,6,7,8-hexahydro-pyrido[4,3-d]pyrimidine;
6-[3-(1H-Benzoimidazol-2-yl)-prop-2-ynyl]-3-(4-morpholin-4-yl-benzyl)-3,4,5,6,7,8-hexahydro-pyrido[4,3-d]pyrimidine;

6-[3-(1H-Benzoimidazol-2-yl)-prop-2-ynyl]-3-(4-cyclohexyl-benzyl)-
3,4,5,6,7,8-hexahydro-pyrido[4,3-d]pyrimidine;
4-{6-[3-(1H-Benzoimidazol-2-yl)-prop-2-ynyl]-5,6,7,8-tetrahydro-4H-
pyrido[4,3-d]pyrimidin-3-ylmethyl}-benzoic acid methyl ester; and
5 4-{6-[3-(1H-Benzoimidazol-2-yl)-prop-2-ynyl]-5,6,7,8-tetrahydro-4H-
pyrido[4,3-d]pyrimidin-3-ylmethyl}-benzoic acid; or
a pharmaceutically acceptable salt thereof.

114. The compound according to Embodiment 107, selected from:
10 6-[3-(1H-Benzoimidazol-2-yl)-prop-2-ynyl]-3-(4-methoxy-benzyl)-
3,4,5,6,7,8-hexahydro-pyrido[4,3-d]pyrimidine;
6-[3-(1H-Benzoimidazol-2-yl)-prop-2-ynyl]-3-(3-methoxy-benzyl)-
3,4,5,6,7,8-hexahydro-pyrido[4,3-d]pyrimidine;
6-[3-(1H-Benzoimidazol-2-yl)-prop-2-ynyl]-3-pyridin-4-ylmethyl-
15 3,4,5,6,7,8-hexahydro-pyrido[4,3-d]pyrimidine; and
6-[3-(1H-Benzoimidazol-2-yl)-prop-2-ynyl]-3-(2-methoxy-pyridin-4-
ylmethyl)-3,4,5,6,7,8-hexahydro-pyrido[4,3-d]pyrimidine; or
a pharmaceutically acceptable salt thereof.

20 115. The compound according to Embodiment 107, selected from:
6-[3-(1H-Benzoimidazol-2-yl)-prop-2-ynyl]-3-biphenyl-4-ylmethyl-
3,4,5,6,7,8-hexahydro-pyrido[4,3-d]pyrimidine; and
6-[3-(1H-Benzoimidazol-2-yl)-prop-2-ynyl]-3-naphthalen-2-ylmethyl-
3,4,5,6,7,8-hexahydro-pyrido[4,3-d]pyrimidine; or
25 a pharmaceutically acceptable salt thereof.


116. The compound according to Embodiment 107, selected from:
6-[3-(1H-Benzoimidazol-2-yl)-prop-2-ynyl]-3-(1H-pyrrol-2-ylmethyl)-
3,4,5,6,7,8-hexahydro-pyrido[4,3-d]pyrimidine;
30 6-[3-(1H-Benzoimidazol-2-yl)-prop-2-ynyl]-3-(1H-imidazol-2-ylmethyl)-
3,4,5,6,7,8-hexahydro-pyrido[4,3-d]pyrimidine;
6-[3-(1H-Benzoimidazol-2-yl)-prop-2-ynyl]-3-(4H-[1,2,4]triazol-3-
ylmethyl)-3,4,5,6,7,8-hexahydro-pyrido[4,3-d]pyrimidine;

- 6-[3-(1H-Benzoimidazol-2-yl)-prop-2-ynyl]-3-(1H-tetrazol-5-ylmethyl)-
3,4,5,6,7,8-hexahydro-pyrido[4,3-d]pyrimidine;
6-[3-(1H-Benzoimidazol-2-yl)-prop-2-ynyl]-3-(1-methyl-1H-pyrrol-2-
ylmethyl)-3,4,5,6,7,8-hexahydro-pyrido[4,3-d]pyrimidine;
5 6-[3-(1H-Benzoimidazol-2-yl)-prop-2-ynyl]-3-thiophen-2-ylmethyl-
3,4,5,6,7,8-hexahydro-pyrido[4,3-d]pyrimidine;
6-[3-(1H-Benzoimidazol-2-yl)-prop-2-ynyl]-3-thiazol-2-ylmethyl-
3,4,5,6,7,8-hexahydro-pyrido[4,3-d]pyrimidine;
6-[3-(1H-Benzoimidazol-2-yl)-prop-2-ynyl]-3-oxazol-2-ylmethyl-
10 3,4,5,6,7,8-hexahydro-pyrido[4,3-d]pyrimidine;
6-[3-(1H-Benzoimidazol-2-yl)-prop-2-ynyl]-3-furan-2-ylmethyl-
3,4,5,6,7,8-hexahydro-pyrido[4,3-d]pyrimidine;
6-[3-(1H-Benzoimidazol-2-yl)-prop-2-ynyl]-3-pyrrol-1-ylmethyl-
3,4,5,6,7,8-hexahydro-pyrido[4,3-d]pyrimidine; and
15 6-[3-(1H-Benzoimidazol-2-yl)-prop-2-ynyl]-3-imidazol-1-ylmethyl-
3,4,5,6,7,8-hexahydro-pyrido[4,3-d]pyrimidine; or
a pharmaceutically acceptable salt thereof.
117. The compound according to Embodiment 107, selected from:
20 6-[3-(1H-Benzoimidazol-2-yl)-prop-2-ynyl]-3-benzooxazol-2-ylmethyl-
3,4,5,6,7,8-hexahydro-pyrido[4,3-d]pyrimidine;
3-Benzofuran-2-ylmethyl-6-[3-(1H-benzoimidazol-2-yl)-prop-2-ynyl]-
3,4,5,6,7,8-hexahydro-pyrido[4,3-d]pyrimidine;
25 3-Benzo[b]thiophen-2-ylmethyl-6-[3-(1H-benzoimidazol-2-yl)-prop-2-
ynyl]-3,4,5,6,7,8-hexahydro-pyrido[4,3-d]pyrimidine;
3-(1H-Benzoimidazol-2-ylmethyl)-6-[3-(1H-benzoimidazol-2-yl)-prop-2-
ynyl]-3,4,5,6,7,8-hexahydro-pyrido[4,3-d]pyrimidine; and
3-(1H-Benzoimidazol-2-ylmethyl)-6-[3-(1H-benzoimidazol-2-yl)-prop-2-
ynyl]-3-(1,3-dihydro-isindol-2-ylmethyl)-3,4,5,6,7,8-hexahydro-
30 pyrido[4,3-d]pyrimidine; or
a pharmaceutically acceptable salt thereof.
118. The compound according to Embodiment 107, selected from:

6-[3-(1H-Benzoimidazol-2-yl)-prop-2-ynyl]-3-cyclohexylmethyl-
3,4,5,6,7,8-hexahydro-pyrido[4,3-d]pyrimidine;
6-[3-(1H-Benzoimidazol-2-yl)-prop-2-ynyl]-3-piperidin-1-ylmethyl-
3,4,5,6,7,8-hexahydro-pyrido[4,3-d]pyrimidine; and
5 6-[3-(1H-Benzoimidazol-2-yl)-prop-2-ynyl]-3-but-2-ynyl-3,4,5,6,7,8-
hexahydro-pyrido[4,3-d]pyrimidine; or
a pharmaceutically acceptable salt thereof.

119. A compound of Formula XI

10



XI

or a pharmaceutically acceptable salt thereof, or a pyrido-N-oxide thereof,
wherein:

R¹ is independently selected from:

- C₅ or C₆ cycloalkyl-(C₁-C₈ alkylenyl);
- 15 Substituted C₅ or C₆ cycloalkyl-(C₁-C₈ alkylenyl);
- C₈-C₁₀ bicycloalkyl-(C₁-C₈ alkylenyl);
- Substituted C₈-C₁₀ bicycloalkyl-(C₁-C₈ alkylenyl);
- 5- or 6-membered heterocycloalkyl-(C₁-C₈ alkylenyl);
- Substituted 5- or 6-membered heterocycloalkyl-(C₁-C₈ alkylenyl);
- 20 8- to 10-membered heterobicycloalkyl-(C₁-C₈ alkylenyl);
- Substituted 8- to 10-membered heterobicycloalkyl-(C₁-C₈ alkylenyl);
- Phenyl-(C₁-C₈ alkylenyl);
- Substituted phenyl-(C₁-C₈ alkylenyl);
- Naphthyl-(C₁-C₈ alkylenyl);
- 25 Substituted naphthyl-(C₁-C₈ alkylenyl);
- 5- or 6-membered heteroaryl-(C₁-C₈ alkylenyl);
- Substituted 5- or 6-membered heteroaryl-(C₁-C₈ alkylenyl);
- 8- to 10-membered heterobiaryl-(C₁-C₈ alkylenyl);
- Substituted 8- to 10-membered heterobiaryl-(C₁-C₈ alkylenyl);

Phenyl;
Substituted phenyl;
Naphthyl;
Substituted naphthyl;
5 5- or 6-membered heteroaryl;
Substituted 5- or 6-membered heteroaryl;
8- to 10-membered heterobiaryl; and
Substituted 8- to 10-membered heterobiaryl;

R² is independently selected from:

10 H;
C₁-C₆ alkyl;
Phenyl-(C₁-C₈ alkylene);
Substituted phenyl-(C₁-C₈ alkylene);
Naphthyl-(C₁-C₈ alkylene);
15 Substituted naphthyl-(C₁-C₈ alkylene);
5- or 6-membered heteroaryl-(C₁-C₈ alkylene);
Substituted 5- or 6-membered heteroaryl-(C₁-C₈ alkylene);
8- to 10-membered heterobiaryl-(C₁-C₈ alkylene);
Substituted 8- to 10-membered heterobiaryl-(C₁-C₈ alkylene);
20 Phenyl-O-(C₁-C₈ alkylene);
Substituted phenyl-O-(C₁-C₈ alkylene);
Phenyl-S-(C₁-C₈ alkylene);
Substituted phenyl-S-(C₁-C₈ alkylene);
Phenyl-S(O)-(C₁-C₈ alkylene);
25 Substituted phenyl-S(O)-(C₁-C₈ alkylene);
Phenyl-S(O)₂-(C₁-C₈ alkylene); and
Substituted phenyl-S(O)₂-(C₁-C₈ alkylene);

Each substituted R¹ and R² group contains from 1 to 4 substituents, each independently on a carbon or nitrogen atom, independently selected from:


30 C₁-C₆ alkyl;
CN;
CF₃;
HO;


- (C₁-C₆ alkyl)-O;
 (C₁-C₆ alkyl)-S(O)₂;
 H₂N;
 (C₁-C₆ alkyl)-N(H);
 5 (C₁-C₆ alkyl)₂-N;
 (C₁-C₆ alkyl)-C(O)O-(C₁-C₈ alkylenyl)_m;
 (C₁-C₆ alkyl)-C(O)O-(1- to 8-membered heteroalkylenyl)_m;
 (C₁-C₆ alkyl)-C(O)N(H)-(C₁-C₈ alkylenyl)_m;
 (C₁-C₆ alkyl)-C(O)N(H)-(1- to 8-membered heteroalkylenyl)_m;
 10 H₂NS(O)₂-(C₁-C₈ alkylenyl);
 (C₁-C₆ alkyl)-N(H)S(O)₂-(C₁-C₈ alkylenyl)_m;
 (C₁-C₆ alkyl)₂-NS(O)₂-(C₁-C₈ alkylenyl)_m;
 3- to 6-membered heterocycloalkyl-(G)_m;
 Substituted 3- to 6-membered heterocycloalkyl-(G)_m;
 15 5- or 6-membered heteroaryl-(G)_m;
 Substituted 5- or 6-membered heteroaryl-(G)_m;
 (C₁-C₆ alkyl)-S(O)₂-N(H)-C(O)-(C₁-C₈ alkylenyl)_m; and
 (C₁-C₆ alkyl)-C(O)-N(H)-S(O)₂-(C₁-C₈ alkylenyl)_m;

wherein each substituent on a carbon atom may further be independently selected
 20 from:

Halo; and
 HO₂C;

- wherein 2 substituents may be taken together with a carbon atom to which they
 are both bonded to form the group C=O;
 25 wherein two adjacent, substantially sp² carbon atoms may be taken together with a
 diradical substituent to form a cyclic diradical selected from:





R is H or C₁-C₆ alkyl;

5 G is CH₂; O, S, S(O); or S(O)₂;

m is an integer of 0 or 1;

wherein each C₈-C₁₀ bicycloalkyl is a bicyclic carbocyclic ring that contains 8-, 9-, or 10-member carbon atoms which are 5,5-fused, 6,5-fused, or 6,6-fused bicyclic rings, respectively, and wherein the ring is saturated or optionally contains one carbon-carbon double bond;

10

wherein each 8- to 10-membered heterobicycloalkyl is a bicyclic ring that contains carbon atoms and from 1 to 4 heteroatoms independently selected from 2 O, 1 S, 1 S(O), 1 S(O)₂, 1 N, 4 N(H), and 4 N(C₁-C₆ alkyl), and wherein when two O atoms or one O atom and one S atom are present, the two O atoms or one O atom and one S atom are not bonded to each other, and wherein the ring is saturated or optionally contains one carbon-carbon or carbon-nitrogen double bond, and wherein the heterobicycloalkyl is a 5,5-fused, 6,5-fused, or 6,6-fused bicyclic ring, respectively,

15

wherein each heterocycloalkyl is a ring that contains carbon atoms and from 1 to 4 heteroatoms independently selected from 2 O, 1 S, 1 S(O), 1 S(O)₂, 1 N, 4 N(H), and 4 N(C₁-C₆ alkyl), and wherein when two O atoms or one O atom and one S atom are present, the two O atoms or one O atom and one

20

S atom are not bonded to each other, and wherein the ring is saturated or optionally contains one carbon-carbon or carbon-nitrogen double bond; wherein each 5-membered heteroaryl contains carbon atoms and from 1 to 4 heteroatoms independently selected from 1 O, 1 S, 1 N(H), 1 N(C₁-C₆ alkyl), and 4 N, and each 6-membered heteroaryl contains carbon atoms and 1 or 2 heteroatoms independently selected from N, N(H), and N(C₁-C₆ alkyl), and 5- and 6-membered heteroaryl are monocyclic rings; wherein each heterobiaryl contains carbon atoms and from 1 to 4 heteroatoms independently selected from 1 O, 1 S, 1 N(H), 1 N(C₁-C₆ alkyl), and 4 N, and where the 8-, 9-, and 10-membered heterobiaryl are 5,5-fused, 6,5-fused, and 6,6-fused bicyclic rings, respectively, and wherein at least 1 of the 2 fused rings of a bicyclic ring is aromatic, and wherein when the O and S atoms both are present, the O and S atoms are not bonded to each other; wherein with any (C₁-C₆ alkyl)₂-N group, the C₁-C₆ alkyl groups may be optionally taken together with the nitrogen atom to which they are attached to form a 5- or 6-membered heterocycloalkyl; and wherein each group and each substituent recited above is independently selected.

120. The compound according to Embodiment 119, selected from:
- 3-(4-Carboxy-benzyl)-3,5,7,8-tetrahydro-4H-pyrido[4,3-d]pyrimidine-6-carboxylic acid benzyl ester;
- 3-(4-Carboxy-benzyl)-3,5,7,8-tetrahydro-4H-pyrido[4,3-d]pyrimidine-6-carboxylic acid pyridin-4-ylmethyl ester;
- 3-(4-Carboxy-benzyl)-3,5,7,8-tetrahydro-4H-pyrido[4,3-d]pyrimidine-6-carboxylic acid 2-methoxy-pyridin-4-ylmethyl ester;
- 3-(4-Carboxy-benzyl)-3,5,7,8-tetrahydro-4H-pyrido[4,3-d]pyrimidine-6-carboxylic acid 3-methoxy-benzyl ester; and
- 3-(4-Carboxy-benzyl)-3,5,7,8-tetrahydro-4H-pyrido[4,3-d]pyrimidine-6-carboxylic acid 4-methoxy-benzyl ester; or
- a pharmaceutically acceptable salt thereof.

121. The compound according to Embodiment 119, selected from:

- 3-(4-Carboxy-benzyl)-3,5,7,8-tetrahydro-4H-pyrido[4,3-d]pyrimidine-6-carboxylic acid piperidin-1-ylmethyl ester;
3-(4-Carboxy-benzyl)-3,5,7,8-tetrahydro-4H-pyrido[4,3-d]pyrimidine-6-carboxylic acid morpholin-4-ylmethyl ester; and
5 3-(4-Carboxy-benzyl)-3,5,7,8-tetrahydro-4H-pyrido[4,3-d]pyrimidine-6-carboxylic acid 4-methyl-piperazin-1-ylmethyl ester; or
a pharmaceutically acceptable salt thereof.
122. The compound according to Embodiment 119, selected from:
10 3-(4-Carboxy-benzyl)-3,5,7,8-tetrahydro-4H-pyrido[4,3-d]pyrimidine-6-carboxylic acid pyrrolidin-1-ylmethyl ester
3-(4-Carboxy-benzyl)-3,5,7,8-tetrahydro-4H-pyrido[4,3-d]pyrimidine-6-carboxylic acid pyrrol-1-ylmethyl ester
3-(4-Carboxy-benzyl)-3,5,7,8-tetrahydro-4H-pyrido[4,3-d]pyrimidine-6-carboxylic acid imidazol-1-ylmethyl ester
15 3-(4-Carboxy-benzyl)-3,5,7,8-tetrahydro-4H-pyrido[4,3-d]pyrimidine-6-carboxylic acid [1,2,4]triazol-4-ylmethyl ester
3-(4-Carboxy-benzyl)-3,5,7,8-tetrahydro-4H-pyrido[4,3-d]pyrimidine-6-carboxylic acid tetrazol-1-ylmethyl ester
20 a pharmaceutically acceptable salt thereof.
123. The compound according to Embodiment 119, selected from:
3-(4-Carboxy-benzyl)-3,5,7,8-tetrahydro-4H-pyrido[4,3-d]pyrimidine-6-carboxylic acid 1,3-dihydro-isoindol-2-ylmethyl ester;
25 3-(4-Carboxy-benzyl)-3,5,7,8-tetrahydro-4H-pyrido[4,3-d]pyrimidine-6-carboxylic acid 1H-indol-2-ylmethyl ester;
3-(4-Carboxy-benzyl)-3,5,7,8-tetrahydro-4H-pyrido[4,3-d]pyrimidine-6-carboxylic acid 1H-benzoimidazol-2-ylmethyl ester;
3-(4-Carboxy-benzyl)-3,5,7,8-tetrahydro-4H-pyrido[4,3-d]pyrimidine-6-carboxylic acid benzothiazol-2-ylmethyl ester;
30 3-(4-Carboxy-benzyl)-3,5,7,8-tetrahydro-4H-pyrido[4,3-d]pyrimidine-6-carboxylic acid benzo[b]thiophen-2-ylmethyl ester;

3-(4-Carboxy-benzyl)-3,5,7,8-tetrahydro-4H-pyrido[4,3-d]pyrimidine-6-carboxylic acid benzofuran-2-ylmethyl ester; and
3-(4-Carboxy-benzyl)-3,5,7,8-tetrahydro-4H-pyrido[4,3-d]pyrimidine-6-carboxylic acid benzooxazol-2-ylmethyl ester; or
5 a pharmaceutically acceptable salt thereof.

124. The compound according to Embodiment 119, selected from:

3-(4-Carboxy-benzyl)-3,5,7,8-tetrahydro-4H-pyrido[4,3-d]pyrimidine-6-carboxylic acid 4-fluoro-benzyl ester;

10 3-(4-Carboxy-benzyl)-3,5,7,8-tetrahydro-4H-pyrido[4,3-d]pyrimidine-6-carboxylic acid 4-chloro-benzyl ester;

3-(4-Carboxy-benzyl)-3,5,7,8-tetrahydro-4H-pyrido[4,3-d]pyrimidine-6-carboxylic acid 4-bromo-benzyl ester;

15 3-(4-Carboxy-benzyl)-3,5,7,8-tetrahydro-4H-pyrido[4,3-d]pyrimidine-6-carboxylic acid 4-iodo-benzyl ester;

3-(4-Carboxy-benzyl)-3,5,7,8-tetrahydro-4H-pyrido[4,3-d]pyrimidine-6-carboxylic acid 4-dimethylamino-benzyl ester;

3-(4-Carboxy-benzyl)-3,5,7,8-tetrahydro-4H-pyrido[4,3-d]pyrimidine-6-carboxylic acid 4-pyrrolidin-1-yl-benzyl ester;

20 3-(4-Carboxy-benzyl)-3,5,7,8-tetrahydro-4H-pyrido[4,3-d]pyrimidine-6-carboxylic acid pentafluorophenylmethyl ester;

3-(4-Carboxy-benzyl)-3,5,7,8-tetrahydro-4H-pyrido[4,3-d]pyrimidine-6-carboxylic acid 4-methanesulfonyl-benzyl ester;

25 3-(4-Carboxy-benzyl)-3,5,7,8-tetrahydro-4H-pyrido[4,3-d]pyrimidine-6-carboxylic acid 4-dimethylsulfamoyl-benzyl ester; and

3-(4-Carboxy-benzyl)-3,5,7,8-tetrahydro-4H-pyrido[4,3-d]pyrimidine-6-carboxylic acid 4-cyano-benzyl ester; or
a pharmaceutically acceptable salt thereof.

30 125. The compound according to Embodiment 119, selected from:

3-(4-Carboxy-benzyl)-3,5,7,8-tetrahydro-4H-pyrido[4,3-d]pyrimidine-6-carboxylic acid benzo[b]thiophen-6-ylmethyl ester; and

3-(4-Carboxy-benzyl)-3,5,7,8-tetrahydro-4H-pyrido[4,3-d]pyrimidine-6-carboxylic acid naphthalen-2-ylmethyl ester; or
a pharmaceutically acceptable salt thereof.

5 126. The compound according to Embodiment 119, selected from:
 3-(4-Carboxy-benzyl)-3,5,7,8-tetrahydro-4H-pyrido[4,3-d]pyrimidine-6-
 carboxylic acid cyclohexylmethyl ester; and
 3-(4-Carboxy-benzyl)-3,5,7,8-tetrahydro-4H-pyrido[4,3-d]pyrimidine-6-
 carboxylic acid but-2-ynyl ester; or
 10 a pharmaceutically acceptable salt thereof.

127. The compound according to Embodiment 119, selected from:
3-(4-Carboxy-benzyl)-3,5,7,8-tetrahydro-4H-pyrido[4,3-d]pyrimidine-6-carboxylic acid furan-2-ylmethyl ester;
15 3-(4-Carboxy-benzyl)-3,5,7,8-tetrahydro-4H-pyrido[4,3-d]pyrimidine-6-carboxylic acid 5-dimethylaminomethyl-furan-2-ylmethyl ester;
3-(4-Carboxy-benzyl)-3,5,7,8-tetrahydro-4H-pyrido[4,3-d]pyrimidine-6-carboxylic acid oxazol-2-ylmethyl ester; and
3-(4-Carboxy-benzyl)-3,5,7,8-tetrahydro-4H-pyrido[4,3-d]pyrimidine-6-carboxylic acid thiazol-2-ylmethyl ester; or
20 a pharmaceutically acceptable salt thereof.

128. A pharmaceutical composition, comprising a compound of Formula I according to Embodiment 1, or a pharmaceutically acceptable salt thereof, admixed with a pharmaceutically acceptable carrier, excipient, or diluent.

129. The pharmaceutical composition according to Embodiment 128,
comprising a compound of Formula I according to any one of Embodiments 2 to
127, or a pharmaceutically acceptable salt thereof, admixed with a
30 pharmaceutically acceptable carrier, excipient, or diluent.

130. A method for inhibiting an MMP-13 enzyme in an animal, comprising administering to the animal an MMP-13 inhibiting amount of a compound of

Formula I according to Embodiment 1, or a pharmaceutically acceptable salt thereof.

131. The method according to Embodiment 130, wherein the compound of
5 Formula I is according to any one of Embodiments 2 to 127, or a pharmaceutically acceptable salt thereof.

132. A method for treating a disease mediated by an MMP-13 enzyme,
comprising administering to a patient suffering from such a disease a nontoxic
10 effective amount of a compound of Formula I according to Embodiment 1, or a pharmaceutically acceptable salt thereof.

133. The method according to Embodiment 132, wherein the compound of
Formula I is according to any one of Embodiments 2 to 127, or a pharmaceutically
15 acceptable salt thereof.

134. A method for treating arthritis, comprising administering to a patient
suffering from an arthritis disease a nontoxic antiarthritic effective amount of a
compound of Formula I according to Embodiment 1, or a pharmaceutically
20 acceptable salt thereof.

135. The method according to Embodiment 134, wherein the compound of
Formula I is according to any one of Embodiments 2 to 127, or a pharmaceutically
acceptable salt thereof.

25 136. A method for treating osteoarthritis, comprising administering to a patient
suffering from osteoarthritis a nontoxic effective amount of a compound of
Formula I according to Embodiment 1, or a pharmaceutically acceptable salt
thereof.

30 137. The method according to Embodiment 136, wherein the compound of
Formula I is according to any one of Embodiments 2 to 127, or a pharmaceutically
acceptable salt thereof.

138. A method for treating rheumatoid arthritis, comprising administering to a patient suffering from rheumatoid arthritis a nontoxic effective amount of a compound of Formula I according to Embodiment 1, or a pharmaceutically acceptable salt thereof.

5

139. The method according to Embodiment 138, wherein the compound of Formula I is according to any one of Embodiments 2 to 127, or a pharmaceutically acceptable salt thereof.

10

140. A method for treating psoriatic arthritis, comprising administering to a patient suffering from psoriatic arthritis a nontoxic effective amount of a compound of Formula I according to Embodiment 1, or a pharmaceutically acceptable salt thereof.

15

141. The method according to Embodiment 140, wherein the compound of Formula I is according to any one of Embodiments 2 to 127, or a pharmaceutically acceptable salt thereof.

142. A method for treating a cancer, comprising administering to a patient suffering from a cancer a nontoxic anti-cancer effective amount of a compound of Formula I according to Embodiment 1, or a pharmaceutically acceptable salt thereof.

143. The method according to Embodiment 142, wherein the compound of Formula I is according to any one of Embodiments 2 to 127, or a pharmaceutically acceptable salt thereof.

144. A method for treating breast carcinoma, comprising administering to a patient suffering from breast carcinoma a nontoxic effective amount of a compound of Formula I according to Embodiment 1, or a pharmaceutically acceptable salt thereof.

30

145. The method according to Embodiment 144, wherein the compound of Formula I is according to any one of Embodiments 2 to 127, or a pharmaceutically acceptable salt thereof.

5 146. A method for treating atherosclerosis, comprising administering to a patient suffering from atherosclerosis a nontoxic effective amount of a compound of Formula I according to Embodiment 1, or a pharmaceutically acceptable salt thereof.

10 147. The method according to Embodiment 146, wherein the compound of Formula I is according to any one of Embodiments 2 to 127, or a pharmaceutically acceptable salt thereof.

15 148. A method for treating inflammation, comprising administering to a patient suffering from inflammation a nontoxic effective amount of a compound of Formula I according to Embodiment 1, or a pharmaceutically acceptable salt thereof.

20 149. The method according to Embodiment 148, wherein the compound of Formula I is according to any one of Embodiments 2 to 127, or a pharmaceutically acceptable salt thereof.

25 150. A method for treating heart failure, comprising administering to a patient suffering from heart failure a nontoxic effective amount of a compound of Formula I according to Embodiment 1, or a pharmaceutically acceptable salt thereof.

30 151. The method according to Embodiment 150, wherein the compound of Formula I is according to any one of Embodiments 2 to 127, or a pharmaceutically acceptable salt thereof.

152. A method for treating age-related macular degeneration, comprising administering to a patient suffering from age-related macular degeneration a

nontoxic effective amount of a compound of Formula I according to Embodiment 1, or a pharmaceutically acceptable salt thereof.

5 153. The method according to Embodiment 152, wherein the compound of Formula I is according to any one of Embodiments 2 to 127, or a pharmaceutically acceptable salt thereof.

10 154. A method for treating chronic obstructive pulmonary disease, comprising administering to a patient suffering from chronic obstructive pulmonary disease a nontoxic effective amount of a compound of Formula I according to Embodiment 1, or a pharmaceutically acceptable salt thereof.

15 155. The method according to Embodiment 154, wherein the compound of Formula I is according to any one of Embodiments 2 to 127, or a pharmaceutically acceptable salt thereof.

20 156. A method for treating heart disease, comprising administering to a patient suffering from heart disease a nontoxic effective amount of a compound of Formula I according to Embodiment 1, or a pharmaceutically acceptable salt thereof.

25 157. The method according to Embodiment 156, wherein the compound of Formula I is according to any one of Embodiments 2 to 127, or a pharmaceutically acceptable salt thereof.

30 158. A method for treating multiple sclerosis, comprising administering to a patient suffering from multiple sclerosis a nontoxic effective amount of a compound of Formula I according to Embodiment 1, or a pharmaceutically acceptable salt thereof.

159. The method according to Embodiment 158, wherein the compound of Formula I is according to any one of Embodiments 2 to 127, or a pharmaceutically acceptable salt thereof.

160. A method for treating psoriasis, comprising administering to a patient suffering from psoriasis a nontoxic effective amount of a compound of Formula I according to Embodiment 1, or a pharmaceutically acceptable salt thereof.

5

161. The method according to Embodiment 160, wherein the compound of Formula I is according to any one of Embodiments 2 to 127, or a pharmaceutically acceptable salt thereof.

10

162. A method for treating asthma, comprising administering to a patient suffering from asthma a nontoxic effective amount of a compound of Formula I according to Embodiment 1, or a pharmaceutically acceptable salt thereof.

15

163. The method according to Embodiment 162, wherein the compound of Formula I is according to any one of Embodiments 2 to 127, or a pharmaceutically acceptable salt thereof.

20

164. A method for treating cardiac insufficiency, comprising administering to a patient suffering from cardiac insufficiency a nontoxic effective amount of a compound of Formula I according to Embodiment 1, or a pharmaceutically acceptable salt thereof.

25

165. The method according to Embodiment 164, wherein the compound of Formula I is according to any one of Embodiments 2 to 127, or a pharmaceutically acceptable salt thereof.


30

166. A method for treating inflammatory bowel disease, comprising administering to a patient suffering from inflammatory bowel disease a nontoxic effective amount of a compound of Formula I according to Embodiment 1, or a pharmaceutically acceptable salt thereof.

167. The method according to Embodiment 166, wherein the compound of Formula I is according to any one of Embodiments 2 to 127, or a pharmaceutically acceptable salt thereof.
- 5 168. A method for treating osteoporosis, comprising administering to a patient suffering from osteoporosis a nontoxic effective amount of a compound of Formula I according to Embodiment 1, or a pharmaceutically acceptable salt thereof.
- 10 169. The method according to Embodiment 168, wherein the compound of Formula I is according to any one of Embodiments 2 to 127, or a pharmaceutically acceptable salt thereof.
- 15 170. A method for treating periodontal diseases, comprising administering to a patient suffering from periodontal diseases a nontoxic effective amount of a compound of Formula I according to Embodiment 1, or a pharmaceutically acceptable salt thereof.
- 20 171. The method according to Embodiment 170, wherein the compound of Formula I is according to any one of Embodiments 2 to 127, or a pharmaceutically acceptable salt thereof.
- 25 172. The method according to any one of Embodiments 130 to 171, wherein the compound of Formula I according to Embodiment 1, or a pharmaceutically acceptable salt thereof, is administered as a pharmaceutical composition according to Embodiment 128 or 129.

DETAILED DESCRIPTION OF THE INVENTION

This invention provides compounds defined by Formula I



I

or a pharmaceutically acceptable salt thereof, or a pyrido-N-oxide thereof, wherein R^1 , Q , Y^1 , Y^2 , Y^3 , and R^2 are as defined above.

5 The invention also provides pharmaceutical compositions comprising a compound of Formula I, or a pharmaceutically acceptable salt thereof, as defined above, together with a pharmaceutically acceptable carrier, diluent, or excipient.

The invention also provides methods of inhibiting an MMP-13 enzyme in an animal, comprising administering to the animal a compound of Formula I, or a pharmaceutically acceptable salt thereof.

10 The invention also provides methods of treating a disease mediated by an MMP-13 enzyme in a patient, comprising administering to the patient a compound of Formula I, or a pharmaceutically acceptable salt thereof, either alone or in a pharmaceutical composition.

15 The invention also provides methods of treating diseases such as heart disease, multiple sclerosis, osteo- and rheumatoid arthritis, arthritis other than osteo- or rheumatoid arthritis, cardiac insufficiency, inflammatory bowel disease, heart failure, age-related macular degeneration, chronic obstructive pulmonary disease, asthma, periodontal diseases, psoriasis, atherosclerosis, and osteoporosis in a patient, comprising administering to the patient a compound of Formula I, or
20 a pharmaceutically acceptable salt thereof, either alone or in a pharmaceutical composition.

The invention also provides combinations, comprising a compound of Formula I, or a pharmaceutically acceptable salt thereof, together with another pharmaceutically active component as described.

25 As seen above, the groups of Formula I include "C₁-C₆ alkyl" groups. C₁-C₆ alkyl groups are straight and branched carbon chains having from 1 to 6 carbon atoms. Examples of C₁-C₆ alkyl groups include methyl, ethyl, 1-propyl, 2-propyl, 1-butyl, 2-butyl, 2,2-dimethylethyl, 1-pentyl, 2-pentyl, 2,2-dimethylpropyl, and 1-hexyl.

The phrase "substituted C₁-C₆ alkyl" means a C₁-C₆ alkyl group as defined above that is substituted with from 1 to 4 substituents independently selected from the list above. Illustrative examples of substituted C₁-C₆ alkyl groups include CH₂OH, CF₂OH, CH₂C(CH₃)₂CO₂CH₃, CF₃, C(O)CF₃, C(O)-
5 CH₃, (CH₂)₄-S-CH₃, CH(CO₂H)CH₂CH₂C(O)NMe₂, (CH₂)₅NH-C(O)-NH₂, CH₂-CH₂-C(H)-(4-fluorophenyl), CH(OCH₃)CH₂CH₃, CH₂SO₂NH₂, and CH(CH₃)CH₂CH₂OC(O)CH₃.

The term "C₂-C₆ alkenyl" means a straight or branched, unsubstituted hydrocarbon group having from 2 to 6 carbon atoms and 1 or 2 carbon-carbon
10 double bonds, and include allenyl groups. Typical examples of C₂-C₆ alkenyl groups include ethenyl, 1-propen-1-yl, 1-propen-2-yl, 2-propen-1-yl, 1-buten-3-yl, 2-penten-2-yl, and 1-hexen-6-yl.

The phrase "substituted C₂-C₆ alkenyl" means a C₂-C₆ alkenyl as defined above, which is substituted with from 1 to 4 substituents independently selected
15 from the list above. Illustrative examples of substituted C₂-C₆ alkenyl groups include C(H)=C(H)CH₂OH, CH=CF₂, CH₂C(H)=C(H)-(CH₂)₂CF₂OH, CH₂C(=CH₂)CO₂CH₃, C(H)=C(H)-CF₃, CH₂-CH₂-C(H)=C(H)-C(O)-CH₃, C(H)=C(CH₃)-S-CH₃, C(H)=C(H)-C(H)=C(CH₃)-CO₂Me, and C(H)=C=C(H)OC(O)CH₃.

20 The term "C₂-C₆ alkynyl" means a straight or branched, unsubstituted hydrocarbon group having from 2 to 6 carbon atoms and 1 or 2 carbon-carbon triple bonds. Typical examples of C₂-C₆ alkynyl groups include ethynyl, 1-propyn-1-yl, 1-propyn-3-yl, 1-butyne-3-yl, 2-pentyne-1-yl, and 1-hexyn-6-yl.

The phrase "substituted C₂-C₆ alkynyl" means a C₂-C₆ alkynyl as defined
25 above, which is substituted with from 1 to 4 substituents independently selected from the list above. Illustrative examples of substituted C₂-C₆ alkynyl groups include C≡CCH₂OH, C≡CF, CH₂C≡C-(CH₂)₂CF₂OH, C≡C-CH₂CO₂CH₃, CH₂C≡C-CF₃, CH₂-CH₂-C≡C-C(O)-CH₃, C≡C-S-CH₃, and C≡C-C(O)OC(O)CH₃.

The term "C₃-C₆ cycloalkyl" means an unsubstituted cyclic hydrocarbon group having from 3 to 6 carbon atoms. C₃-C₆ cycloalkyl may optionally contain one carbon-carbon double bond. The group C₃-C₆ cycloalkyl includes cyclopropyl, cyclobutyl, cyclopentyl, cyclopenten-1-yl, cyclopenten-4-yl, and cyclohexyl.

The phrase "substituted C₃-C₆ cycloalkyl" means a C₃-C₆ cycloalkyl as defined above, which is substituted with from 1 to 4 substituents independently selected from the list above. Illustrative examples of substituted C₃-C₆ cycloalkyl groups include 1-hydroxy-cyclopropyl, cyclobutanon-3-yl, 3-(3-phenyl-ureido)-cyclopent-1-yl, and 4-carboxy-cyclohexyl.

The phrase "3- to 6-membered heterocycloalkyl" means an unsubstituted saturated cyclic group having carbon atoms and 1 or 2 heteroatoms independently selected from 2 O, 1 S, 1 S(O), 1 S(O)₂, 1 N, 2 N(H), and 2 N(C₁-C₆ alkyl), wherein when two O atoms or one O atom and one S atom are present, the two O atoms or one O atom and one S atom are not bonded to each other. Optionally, a 3- to 6-membered heterocycloalkyl may contain one carbon-carbon or carbon-nitrogen double bond. Illustrative examples of 3- to 6-membered heterocycloalkyl includes aziridin-1-yl, 1-oxa-cyclobutan-2-yl, tetrahyrdofuran-3-yl, morpholin-4-yl, 2-thiacyclohex-1-yl, 2-oxo-2-thiacyclohe-1-yl, 2,2-dioxo-2-thiacyclohex-1-yl, and 4-methyl-piperazin-2-yl.

The phrase "substituted 3- to 6-membered heterocycloalkyl" means a 3- to 6-membered heterocycloalkyl as defined above, which is substituted with from 1 to 4 substituents independently selected from the list above. Illustrative examples of substituted 3- to 6-membered heterocycloalkyl include 2-hydroxy-aziridin-1-yl, 3-oxo-1-oxacyclobutan-2-yl, 2,2-dimethyl-tetrahydrofuran-3-yl, 3-carboxy-morpholin-4-yl, and 1-cyclopropyl-4-methyl-piperazin-2-yl.

The term "C₁-C₈ alkylenyl" means a saturated hydrocarbon diradical that is straight or branched and has from 1 to 8 carbon atoms. C₁-C₈ alkylenyl having from 2 to 8 carbon atoms may optionally independently contain one carbon-carbon double bond. Illustrative examples of C₁-C₈ alkylenyl include CH₂, CH₂CH₂, C(CH₃)H, C(H)(CH₃)CH₂CH₂, and CH₂C(H)=C(H)CH₂CH₂CH₂CH₂CH₂.

The term "1- to 8-membered heteroalkylenyl" means a saturated diradical chain that is straight or branched and contains from 1 to 7 carbon atoms and 1 heteroatom selected from O, S, N(H), and N(C₁-C₆ alkyl). 2- to 8-membered heteroalkylenyl, having from 2 to 8 chain atoms, may optionally independently
5 contain one carbon-carbon double bond. Illustrative examples of 1- to 8-membered heteroalkylenyl include OCH₂, CH₂CH₂O, C(CH₃)HS, and CH₂C(H)=C(H)CH₂N(H)CH₂CH₂CH₂.

The phrase "C₃-C₆ cycloalkyl-(C₁-C₈ alkylenyl)" means a C₃-C₆ cycloalkyl, as defined above, bonded through a C₁-C₈ alkylenyl, as defined above.
10 Illustrative examples of C₃-C₆ cycloalkyl-(C₁-C₈ alkylenyl include cyclopropylmethyl, 1-cyclopentyl-hex-2-yl, and 2-cyclobutyl-but-2-yl.

The phrase "Substituted C₃-C₆ cycloalkyl-(C₁-C₈ alkylenyl)" means a C₃-C₆ cycloalkyl-(C₁-C₈ alkylenyl), as defined above, substituted on C₃-C₆ cycloalkyl and/or C₁-C₈ alkylenyl with from 1 to 4 substituents, as defined above. Illustrative
15 examples of substituted C₃-C₆ cycloalkyl-(C₁-C₈ alkylenyl include cyclopropylcarbonyl and 1-(1-aminomethyl-cyclopentyl)-hex-2-yl.

The phrase "C₅ or C₆ cycloalkyl-(C₁-C₈ alkylenyl)" means a cyclopentyl or cyclohexyl bonded through a C₁-C₈ alkylenyl, as defined above, wherein the cycloalkyl optionally contains 1 carbon-carbon double bond.

20 The phrase "Substituted C₅ or C₆ cycloalkyl-(C₁-C₈ alkylenyl)" means a substituted cyclopentyl or cyclohexyl, wherein the substituents are as defined above, bonded through a C₁-C₈ alkylenyl, as defined above, wherein the cycloalkyl optionally contains 1 carbon-carbon double bond.

The phrase "C₈-C₁₀ bicycloalkyl-(C₁-C₈ alkylenyl)" means a cyclopentyl or cyclohexyl fused to another cyclopentyl or cyclohexyl to give a 5,5-, 5,6-, or 6,6-fused bicyclic carbocyclic group, which is bonded through a C₁-C₈ alkylenyl, as defined above, wherein the bicycloalkyl optionally contains 1 carbon-carbon double bond.

30 The phrase "Substituted C₈-C₁₀ bicycloalkyl-(C₁-C₈ alkylenyl)" means a C₈-C₁₀ bicycloalkyl, as defined above, substituted with from 1 to 4 substituents, as defined above, bonded through a C₁-C₈ alkylenyl, as defined above.

The phrase "5- or 6-membered heterocycloalkyl-(C₁-C₈ alkylenyl)" means a 5- or 6-membered ring containing carbon atoms and 1 or 2 heteroatoms selected

from 1 O, 1 S, 1 N, 2 N(H), and 2 N(C₁-C₆ alkyl), bonded through a C₁-C₈ alkylenyl, as defined above.

The phrase "Substituted 5- or 6-membered heterocycloalkyl-(C₁-C₆ alkylenyl)" means a 5- or 6-membered heterocycloalkyl, as defined above,
5 substituted with from 1 to 4 substituents, as defined above, bonded through a C₁-C₈ alkylenyl, as defined above.

The phrase "8- to 10-membered heterobicycloalkyl-(C₁-C₈ alkylenyl)" means a 5- or 6-membered ring fused to another 5- or 6-membered ring to give a 5,5-, 5,6-, or 6,6-fused bicyclic group containing carbon atoms and from 1 to 4
10 heteroatoms independently selected from 2 O, 1 S, 1 S(O), 1 S(O)₂, 1 N, 4 N(H), and 4 N(C₁-C₆ alkyl), bonded through a C₁-C₈ alkylenyl, as defined above, wherein the bicycloalkyl optionally contains 1 carbon-carbon double bond or 1 carbon-nitrogen double bond.

The phrase "Substituted 8- to 10-membered heterobicycloalkyl-(C₁-C₆ alkylenyl)" means an 8- to 10-membered heterobicycloalkyl, as defined above,
15 substituted with from 1 to 4 substituents, as defined above, bonded through a C₁-C₈ alkylenyl, as defined above.

The phrase "3- to 6-membered heterocycloalkyl-(C₁-C₈ alkylenyl)" means a 3- to 6-membered heterocycloalkyl, as defined above, bonded through a C₁-C₈
20 alkylenyl, as defined above.

The phrase "Substituted 3- to 6-membered heterocycloalkyl-(C₁-C₈ alkylenyl)" means a substituted 3- to 6-membered heterocycloalkyl, as defined above, bonded through a C₁-C₈ alkylenyl, as defined above.

The phrase "Phenyl-(C₁-C₈ alkylenyl)" means a phenyl group bonded
25 through a C₁-C₈ alkylenyl diradical, wherein C₁-C₈ alkylenyl is as defined above. Illustrative examples of phenyl-(C₁-C₈ alkylenyl) include benzyl, 2-phenylethyl, 1-phenyl-prop-1-yl, and 3-phenyl-heptyl.

The phrase "Substituted phenyl-(C₁-C₈ alkylenyl)" means a phenyl-(C₁-C₈ alkylenyl) as defined above, which is substituted on phenyl and/or C₁-C₈ alkylenyl
30 with from 1 to 4 substituents independently selected from the list above. Illustrative examples of substituted phenyl-(C₁-C₈ alkylenyl) include 4-fluorophenylmethyl, 2-(4-carboxy-phenyl)-ethyl, 1-(2,4-dimethoxy-phenyl)-2-oxopropyl, and 1-phenyl-5,5-difluoro-oct-3-yl.

The term “naphthyl” includes 1-naphthyl and 2-naphthyl.

The phrase “Naphthyl-(C₁-C₈ alkylene)” means a naphthyl group as defined above bonded through a C₁-C₈ alkylene diradical, wherein C₁-C₈ alkylene is as defined above. Illustrative examples of naphthyl-(C₁-C₈ alkylene) include naphth-1-ylmethyl, 2-(naphth-1-yl)ethyl, and 3-(naphth-2-yl)-1-heptyl.

The phrase “Substituted naphthyl-(C₁-C₈ alkylene)” means a naphthyl-(C₁-C₈ alkylene) as defined above, which is substituted on naphthyl and/or C₁-C₈ alkylene with from 1 to 4 substituents independently selected from the list above. Illustrative examples of substituted phenyl-(C₁-C₈ alkylene) include 4-fluoro-(naphth-1-yl)methyl, 2-(4-carboxy-(naphth-1-yl))-ethyl, 1-(2,4-dimethoxy-(naphth-1-yl))-2-oxo-propyl, and 1-(naphth-2-yl)-5,5-difluorohept-2-yl.

The phrase “5- or 6-membered heteroaryl” means a 5-membered, monocyclic heteroaryl having carbon atoms and from 1 to 4 heteroatoms independently selected from 1 O, 1 S, 1 N(H), 1 N(C₁-C₆ alkyl), and 4 N, or a 6-membered, monocyclic heteroaryl having carbon atoms and 1 or 2 heteroatoms selected from 2 N, and wherein:


(i) The phrase “5-membered, monocyclic heteroaryl” means a 5-membered, monocyclic, aromatic ring group as defined above having carbon atoms and from 1 to 4 heteroatoms selected from 1 O, 1 S, 1 N(H), 1 N(C₁-C₆ alkyl), and 4 N. Illustrative examples of a 5-membered, monocyclic heteroaryl include thiophen-2-yl, furan-2-yl, pyrrol-3-yl, pyrrol-1-yl, imidazol-4-yl, isoxazol-3-yl, oxazol-2-yl, thiazol-4-yl, tetrazol-1-yl, 1,2,4-oxadiazol-3-yl, 1,2,4-triazol-1-yl, and pyrazol-3-yl; and

(ii) The phrase “6-membered, monocyclic heteroaryl” means a 6-membered, monocyclic, aromatic ring group as defined above having carbon atoms and 1 or 2 N. Illustrative examples of a 6-membered, monocyclic heteroaryl include pyridin-2-yl, pyridin-4-yl, pyrimidin-2-yl, pyridazin-4-yl, and pyrazin-2-yl.

The phrase “8- to 10-membered heterobiaryl” means an 8-membered, 5,5-fused bicyclic heteroaryl, a 9-membered, 6,5-fused bicyclic heteroaryl, or a 10-membered, 6,6-fused bicyclic heteroaryl, having carbon atoms and from 1 to 4 heteroatoms independently selected from 1 O, 1 S, 1 N(H), 1 N(C₁-C₆ alkyl), and 4 N, wherein at least one of the 2 fused rings is aromatic, and wherein when the O

and S atoms both are present, the O and S atoms are not bonded to each other, which are as defined below:

- (i) The phrase “8-membered, 5,5-fused bicyclic heteroaryl” means a
an 8-membered aromatic, fused-bicyclic ring group as defined above having
5 carbon atoms and from 1 to 4 heteroatoms selected from 1 O, 1 S, 1 N(H), 1 N(C₁-
C₆ alkyl), and 4 N. Illustrative examples of an 8-membered, fused-bicyclic
heteroaryl include



- (ii) The phrase “9-membered, 6,5-fused bicyclic heteroaryl” means a
10 9-membered aromatic, fused-bicyclic ring group as defined above having carbon
atoms and from 1 to 4 heteroatoms selected from 1 O, 1 S, 1 N(H), 1 N(C₁-C₆
alkyl), and 4 N. Illustrative examples of a 9-membered, fused-bicyclic heteroaryl
include indol-2-yl, indol-6-yl, iso-indol-2-yl, benzimidazol-2-yl,
benzimidazol-1-yl, benztriazol-1-yl, benztriazol-5-yl, benzoxazol-2-yl,
15 benzothiophen-5-yl, and benzofuran-3-yl; and

- (iii) The phrase “10-membered, 6,5-fused bicyclic heteroaryl” means a
10-membered aromatic, fused-bicyclic ring group as defined above having carbon
atoms and from 1 to 4 heteroatoms selected from 1 O, 1 S, 1 N(H), 1 N(C₁-C₆
alkyl), and 4 N. Illustrative examples of a 10-membered, fused-bicyclic heteroaryl
20 include quinolin-2-yl, isoquinolin-7-yl, and benzopyrimidin-2-yl.


The phrases “substituted 5- or 6-membered heteroaryl” and “substituted 8-
to 10-membered heterobiaryl” means a 5- or 6-membered heteroaryl, as defined
above, or an 8- to 10-membered heterobiaryl, as defined above, respectively,
which is substituted on a carbon (CH) atom, and/or nitrogen [N(H)] atom in the
25 case of 5-, 8- to 10-membered heterobiaryl, with from 1 to 4 substituents
independently selected from the list above.

Illustrative examples of substituted 5-membered, monocyclic heteroaryl
groups include 2-hydroxy-oxazol-4-yl, 5-chloro-thiophen-2-yl,

1-methylimidazol-5-yl, 1-propyl-pyrrol-2-yl, 1-acetyl-pyrazol-4-yl, 1-methyl-1,2,4-triazol-3-yl, and 2-hexyl-tetrazol-5-yl.

Illustrative examples of substituted 6-membered, monocyclic heteroaryl groups include 4-acetyl-pyridin-2-yl, 3-fluoro-pyridin-4-yl, 5-carboxy-pyrimidin-2-yl, 6-tertiary butyl-pyridazin-4-yl, and 5-hydroxymethyl-pyrazin-2-yl.

Illustrative examples of substituted 8-membered, 5,5-fused bicyclic heteroaryl include:



Illustrative examples of substituted 9-membered, 5,6-fused bicyclic heteroaryl include 3-(2-aminomethyl)-indol-2-yl, 2-carboxy-indol-6-yl, 1-(methanesulfonyl)-iso-indol-2-yl, 5-trifluoromethyl-6,7-difluoro-4-hydroxymethyl-benzimidazol-2-yl, 4-(3-methylureido)-2-cyano-benzimidazol-1-yl, 1-methylbenzimidazol-6-yl, 1-acetylbenztriazol-7-yl, 1-methanesulfonyl-indol-3-yl, 1-cyano-6-aza-indol-5-yl, and 1-(2,6-dichlorophenylmethyl)-benzpyrazol-3-yl.

Illustrative examples of substituted 10-membered, 6,6-fused bicyclic heteroaryl include 5,7-dichloro-quinolin-2-yl, isoquinolin-7-yl-1-carboxylic acid ethyl ester, and 3-bromo-benzopyrimidin-2-yl.

The phrase “5- or 6-membered heteroaryl-(C₁-C₈ alkylenyl)” means a 5- or 6-membered heteroaryl, as defined above, bonded through a C₁-C₈ alkylenyl, as defined above.

The phrase “Substituted 5- or 6-membered heteroaryl-(C₁-C₈ alkylenyl)” means a 5- or 6-membered heteroaryl-(C₁-C₈ alkylenyl), as defined above, which is substituted on 5- or 6-membered heteroaryl and/or C₁-C₈ alkylenyl with from 1 to 4 substituents independently selected from the list above.


Illustrative examples of substituted 5-membered heteroaryl-(C₁-C₈ alkylenyl) groups include 2-hydroxy-oxazol-4-ylmethyl, 4-(5-chloro-thiophen-2-yl)-hex-1-yl, and 2-tetrazol-5-yl-octyl.

Illustrative examples of substituted 6-membered heteroaryl-(C₁-C₈ alkylenyl) groups include 4-acetyl-pyridin-2-ylmethyl, 7-(3-fluoro-pyridin-4-yl)-hept-2-yl, and 2-(5-hydroxymethyl-pyrazin-2-yl)-1,1-difluoro-2-hydroxy-prop-2-yl.

5 The phrase “8- to 10-membered heterobiaryl-(C₁-C₈ alkylenyl)” means an 8- to 10-membered heterobiaryl, as defined above, bonded through a C₁-C₈ alkylenyl, as defined above.

 The phrase “Substituted 8- to 10-membered heterobiaryl-(C₁-C₈ alkylenyl)” means an 8- to 10-membered heterobiaryl-(C₁-C₈ alkylenyl), as
10 defined above, which is substituted on 8- to 10-membered heterobiaryl and/or C₁-C₈ alkylenyl with from 1 to 4 substituents independently selected from the list above.

Illustrative examples of substituted 8-membered heterobiaryl-(C₁-C₈ alkylenyl) include:



Illustrative examples of substituted 9-membered heterobiaryl-(C₁-C₈ alkylenyl) include 3-(2-aminomethyl)-indol-2-ylmethyl, and 1-(1-(2,6-dichlorophenylmethyl)-benzpyrazol-3-yl)-prop3-yl.

20 Illustrative examples of substituted 10-membered heterobiaryl-(C₁-C₈ alkylenyl) include 5,7-dichloro-quinolin-2-ylmethyl, and 5-(3-bromo-benzopyrimidin-2-yl)-oct-2-yl.

The phrase “(C₁-C₆ alkyl)-O” means a C₁-C₆ alkyl group, as defined above, bonded through an oxygen atom.

25 The phrase “(C₁-C₆ alkyl)-S” means a C₁-C₆ alkyl group, as defined above, bonded through an sulfur atom.

The phrase “(C₁-C₆ alkyl)-S(O)₂” means a C₁-C₆ alkyl group, as defined above, bonded through a sulfur atom, which sulfur atom is substituted with two oxygen atoms.

The phrase “(C₁-C₆ alkyl)-N(H)” means a C₁-C₆ alkyl group, as defined above, bonded through a nitrogen atom, which is bonded to a hydrogen atom.

5 The phrase “(C₁-C₆ alkyl)₂-N” means two independently selected C₁-C₆ alkyl groups, as defined above, including cyclic groups wherein the two C₁-C₆ alkyl groups are taken together with the nitrogen atom to which they are both bonded to form a 5- or 6-membered heterocycloalkyl, bonded through a nitrogen atom.

The phrase “(C₁-C₆ alkyl)-OC(O)” means a C₁-C₆ alkyl, as defined above, bonded through an oxygen atom-carbonyl carbon atom.

10 The phrase “(C₁-C₆ alkyl)-C(O)O-(C₁-C₈ alkylene)_m”, wherein m is an integer of 0 or 1, means when, m is 0, a C₁-C₆ alkyl group, as defined above, bonded through a carbonyl carbon atom-oxygen atom, and, when m is 1, a C₁-C₆ alkyl group, as defined above, bonded through a carbonyl carbon atom-oxygen atom-(C₁-C₈ alkylene), wherein C₁-C₈ alkylene is as defined above.

15 The phrase “(C₁-C₆ alkyl)-C(O)O-(1- to 8-membered heteroalkylene)_m”, wherein m is an integer of 0 or 1, means, when m is 0, a C₁-C₆ alkyl group, as defined above, bonded through a carbonyl carbon atom-oxygen atom, and, when m is 1, a C₁-C₆ alkyl group, as defined above, bonded through a carbonyl carbon atom-oxygen atom-(1- to 8-membered heteroalkylene), wherein 1- to 8-membered heteroalkylene is as defined above.

20 The phrase “(C₁-C₆ alkyl)-C(O)N(H)-(C₁-C₈ alkylene)_m”, wherein m is an integer of 0 or 1, means, when m is 0, a C₁-C₆ alkyl group, as defined above, bonded through a carbonyl carbon atom-nitrogen atom, which is bonded to a hydrogen atom, and, when m is 1, a C₁-C₆ alkyl group, as defined above, bonded through a carbonyl carbon atom-nitrogen atom-(C₁-C₈ alkylene), wherein C₁-C₈ alkylene is as defined above and the nitrogen atom is bonded to a hydrogen atom.

25 The phrase “(C₁-C₆ alkyl)-C(O)N(H)-(1- to 8-membered heteroalkylene)_m”, wherein m is an integer of 0 or 1, means when, m is 0, a C₁-C₆ alkyl group, as defined above, bonded through a carbonyl carbon atom-nitrogen atom, which is bonded to a hydrogen atom, and, when m is 1, a C₁-C₆ alkyl group, as defined above, bonded through a carbonyl carbon atom-nitrogen atom-(1- to 8-membered heteroalkylene), wherein 1- to 8-membered heteroalkylene is as defined above and the nitrogen atom is bonded to a hydrogen atom.

The phrase " $\text{H}_2\text{NS(O)}_2\text{-(C}_1\text{-C}_8\text{ alkylenyl)}$ " means an amino bonded through a sulfur atom-($\text{C}_1\text{-C}_8\text{ alkylenyl}$), wherein the $\text{C}_1\text{-C}_8\text{ alkylenyl}$ is as defined above and the sulfur atom is bonded to two oxygen atoms.

5 The phrase " $\text{(C}_1\text{-C}_6\text{ alkyl)-N(H)S(O)}_2\text{-(C}_1\text{-C}_8\text{ alkylenyl)}_m$ ", wherein m is an integer of 0 or 1, means, when m is 0, a $\text{C}_1\text{-C}_6\text{ alkyl}$, as defined above, bonded through a nitrogen atom-sulfur atom, and, when m is 1, a $\text{C}_1\text{-C}_6\text{ alkyl}$, as defined above, bonded through a nitrogen atom-sulfur atom-($\text{C}_1\text{-C}_8\text{ alkylenyl}$), wherein the nitrogen atom is bonded to a hydrogen atom, the sulfur atom is bonded to two oxygen atoms, and $\text{C}_1\text{-C}_8\text{ alkylenyl}$ is as defined above.

10 The phrase " $\text{(C}_1\text{-C}_6\text{ alkyl)}_2\text{-NS(O)}_2\text{-(C}_1\text{-C}_8\text{ alkylenyl)}_m$ ", wherein m is an integer of 0 or 1, means, when m is 0, two $\text{C}_1\text{-C}_6\text{ alkyl}$ groups, as defined above, including cyclic groups wherein the two $\text{C}_1\text{-C}_6\text{ alkyl}$ groups are taken together with the nitrogen atom to which they are both bonded to form a 5- or 6-membered heterocycloalkyl, each bonded through a nitrogen atom-sulfur atom, and, when m
15 is 1, two $\text{C}_1\text{-C}_6\text{ alkyl}$ groups, as defined above, each bonded through a nitrogen atom-sulfur atom-($\text{C}_1\text{-C}_8\text{ alkylenyl}$), wherein the nitrogen atom is bonded to a hydrogen atom, the sulfur atom is bonded to two oxygen atoms, and $\text{C}_1\text{-C}_8\text{ alkylenyl}$ is as defined above.

The phrase " $\text{3- to 6-membered heterocycloalkyl-(G)}_m$ ", wherein m is an
20 integer of 0 or 1, means, when m is 0, a 3- to 6-membered heterocycloalkyl, as defined above, and, when m is 1, a 3- to 6-membered heterocycloalkyl, as defined above, bonded through a group G, as defined above.

The phrase " $\text{Substituted 3- to 6-membered heterocycloalkyl-(G)}_m$ ", wherein m is an integer of 0 or 1, means, when m is 0, a substituted 3- to 6-
25 membered heterocycloalkyl, as defined above, and, when m is 1, a substituted 3- to 6-membered heterocycloalkyl, as defined above, bonded through a group G, as defined above.

The phrase " $\text{5- or 6-membered heteroaryl-(G)}_m$ ", wherein m is an integer of 0 or 1, means, when m is 0, a 5- or 6-membered heteroaryl, as defined above,
30 and, when m is 1, a 5- or 6-membered heteroaryl, as defined above, bonded through a group G, as defined above.

The phrase " $\text{Substituted 5- or 6-membered heteroaryl-(G)}_m$ ", wherein m is an integer of 0 or 1, means, when m is 0, a substituted 5- or 6-membered

heteroaryl, as defined above, and, when m is 1, a substituted 5- or 6-membered heteroaryl, as defined above, bonded through a group G, as defined above.

The term "Phenyl-O-(C₁-C₈ alkylenyl)" means a phenyl bonded through an oxygen atom, which is bonded through a C₁-C₈ alkylenyl, wherein C₁-C₈ alkylenyl is as defined above. Illustrative examples of phenyl-O-(C₁-C₈ alkylenyl) include phenoxymethyl and 2-phenoxyethyl.

The term "Substituted phenyl-O-(C₁-C₈ alkylenyl)" means a phenyl-O-(C₁-C₈ alkylenyl) group, as defined above, that is substituted with from 1 to 4 substituents as defined above for R². Illustrative examples of substituted phenyl-O-(C₁-C₈ alkylenyl) include 4-fluorophenoxymethyl and 2-phenoxy-methylcarbonyl.

The term "Phenyl-S-(C₁-C₈ alkylenyl)" means a phenyl bonded through an sulfur atom, which is bonded through a C₁-C₈ alkylenyl, wherein C₁-C₈ alkylenyl is as defined above. Illustrative examples of phenyl-S-(C₁-C₈ alkylenyl) include thiophenoxymethyl and 2-thiophenoxyethyl.

The term "Substituted phenyl-S-(C₁-C₈ alkylenyl)" means a phenyl-S-(C₁-C₈ alkylenyl) group, as defined above, that is substituted with from 1 to 4 substituents as defined above for R². Illustrative examples of substituted phenyl-S-(C₁-C₈ alkylenyl) include 4-fluorothiophenoxymethyl and 2-thiophenoxy-methylcarbonyl.

The term "Phenyl-S(O)-(C₁-C₈ alkylenyl)" means a phenyl bonded through an sulfur atom, which is bonded through a C₁-C₈ alkylenyl, wherein C₁-C₈ alkylenyl is as defined above and the sulfur atom is also bonded to an oxygen atom. Illustrative examples of phenyl-S(O)-(C₁-C₈ alkylenyl) include phenyl-S(=O)-CH₂ and phenyl-S(=O)-CH₂CH₂.

The term "Substituted phenyl-S(O)-(C₁-C₈ alkylenyl)" means a phenyl-S(O)-(C₁-C₈ alkylenyl) group, as defined above, that is substituted with from 1 to 4 substituents as defined above for R². Illustrative examples of substituted phenyl-S(O)-(C₁-C₈ alkylenyl) include (4-Fluoro-phenyl)-S(=O)-CH₂ and phenyl-S(=O)-CH₂C(=O).

The term "Phenyl-S(O)₂-(C₁-C₈ alkylenyl)" means a phenyl bonded through an sulfur atom, which is bonded through a C₁-C₈ alkylenyl, wherein C₁-C₈ alkylenyl is as defined above and the sulfur atom is also bonded to two oxygen

atoms. Illustrative examples of phenyl-S(O)₂-(C₁-C₈ alkylenyl) include phenyl-S(=O)₂-CH₂ and phenyl-S(=O)₂-CH₂CH₂.

The term "Substituted phenyl-S(O)₂-(C₁-C₈ alkylenyl)" means a phenyl-S(O)₂-(C₁-C₈ alkylenyl) group, as defined above, that is substituted with from 1 to 4 substituents as defined above for R². Illustrative examples of substituted phenyl-S(O)₂-(C₁-C₈ alkylenyl) include (4-Fluoro-phenyl)-S(=O)₂-CH₂ and phenyl-S(=O)₂-CH₂C(=O).


The term "(C₁-C₆ alkyl)-S(O)₂-N(H)-C(O)-(C₁-C₈ alkylenyl)_m", wherein m is an integer of 0 or 1, means, when m is 0, a C₁-C₆ alkyl group, as defined above, bonded through a sulfur atom, which is bonded through a nitrogen atom, which is bonded through a carbon atom, wherein the sulfur atom is bonded to two oxygen atoms, the nitrogen atom is bonded to a hydrogen atom, and the carbon atom is doubly bonded to an oxygen atom to form a carbonyl group; and when m is 1, the term means a C₁-C₆ alkyl group, as defined above, bonded through a sulfur atom, which is bonded through a nitrogen atom, which is bonded through a carbon atom, which is bonded through a C₁-C₈ alkylenyl group, as defined above, wherein the sulfur atom is bonded to two oxygen atoms, the nitrogen atom is bonded to a hydrogen atom, and the carbon atom is doubly bonded to an oxygen atom to form a carbonyl group. Illustrative examples of (C₁-C₆ alkyl)-S(=O)₂-N(H)-C(O)-(C₁-C₈ alkylenyl)_m include CH₃-S(O)₂-N(H)-C(=O) and CH₃-S(O)₂-N(H)-C(=O)-CH₂.

The term "(C₁-C₆ alkyl)-C(O)-N(H)-S(O)₂-(C₁-C₈ alkylenyl)_m", wherein m is an integer of 0 or 1, means, when m is 0, a C₁-C₆ alkyl group, as defined above, bonded through a carbon atom, which is bonded through a nitrogen atom, which is bonded through a sulfur atom, wherein the sulfur atom is bonded to two oxygen atoms, the nitrogen atom is bonded to a hydrogen atom, and the carbon atom is doubly bonded to an oxygen atom to form a carbonyl group; and when m is 1, the term means a C₁-C₆ alkyl group, as defined above, bonded through a carbon atom, which is bonded through a nitrogen atom, which is bonded through a sulfur atom, which is bonded through a C₁-C₈ alkylenyl group, as defined above, wherein the sulfur atom is bonded to two oxygen atoms, the nitrogen atom is bonded to a hydrogen atom, and the carbon atom is doubly bonded to an oxygen atom to form a carbonyl group. Illustrative examples of (C₁-C₆ alkyl)-C(O)-N(H)-S(O)₂-(C₁-C₈ alkylenyl)_m include CH₃-C(=O)-N(H)-S(=O)₂ and CH₃-C(=O)-N(H)-S(=O)₂-CH₂.

Preferred substituents for substituted phenyl, substituted naphthyl (i.e., substituted 1-naphthyl or substituted 2-naphthyl), and preferred substituents at carbon atoms for substituted 5-membered, monocyclic heteroaryl, substituted 6-membered, monocyclic heteroaryl, and substituted 9- or 10-membered, fused-bicyclic heteroaryl are C₁-C₄ alkyl, halo, OH, O-C₁-C₄ alkyl, 1,2-methylenedioxy, oxo (“=O”), CN, NO₂, N₃, NH₂, N(H)CH₃, N(CH₃)₂, C(O)CH₃, OC(O)-C₁-C₄ alkyl, C(O)-H, CO₂H, CO₂-(C₁-C₄ alkyl), C(O)-N(H)OH, C(O)NH₂, C(O)NHMe, C(O)N(Me)₂, NHC(O)CH₃, N(H)C(O)NH₂, SH, S-C₁-C₄ alkyl, C≡CH, C(=NOH)-H, C(=NOH)-CH₃, CH₂OH, CH₂NH₂, CH₂N(H)CH₃, CH₂N(CH₃)₂, C(H)F-OH, CF₂-OH, S(O)₂NH₂, S(O)₂N(H)CH₃, S(O)₂N(CH₃)₂, S(O)-CH₃, S(O)₂CH₃, S(O)₂CF₃, or NHS(O)₂CH₃.

Especially preferred substituents are 1,2-methylenedioxy, methoxy, ethoxy, -O-C(O)CH₃, carboxy, carbomethoxy, and carboethoxy.

The term “1,2-methylenedioxy” means the diradical group -O-CH₂-O-, wherein the substituent 1,2-methylenedioxy is bonded to adjacent carbon atoms of the group which is substituted to form a 5-membered ring. Illustrative examples of groups substituted by 1,2-methylenedioxy include 1,3-benzoxazol-5-yl of formula B



B

which is a phenyl group substituted by 1,2-methylenedioxy.

A fused-bicyclic group is a group wherein two ring systems share two, and only two, atoms.

It should be appreciated that the groups heteroaryl or heterocycloalkyl may not contain two ring atoms bonded to each other which atoms are oxygen and/or sulfur atoms.

The term “oxo” means =O. Oxo is attached at a carbon atom unless otherwise noted. Oxo, together with the carbon atom to which it is attached forms a carbonyl group (i.e., C=O).


The term "heteroatom" includes O, S, S(O), S(O)₂, N, N(H), and N(C₁-C₆ alkyl).

The term "halo" includes fluoro, chloro, bromo, and iodo.

The term "amino" means NH₂.

5 The phrase "two adjacent, substantially sp² carbon atoms" means carbon atoms that comprise a carbon-carbon double bond that is capable of being substituted on each carbon atom, wherein the carbon-carbon double bond is contained in an aromatic or nonaromatic, cyclic or acyclic, or carbocyclic or heterocyclic group.

10 The phrase "pyrido-N-oxide" means an N-oxide of a 6-membered heteroaryl which contains a pyridinyl radical or an N-oxide of a substituted 6-membered heteroaryl which contains a substituted pyridinyl radical. A pyrido-N-oxide is optionally present in a compound of Formula I when R¹ and/or R² contains a 6-membered heteroaryl or a substituted 6-membered heteroaryl which
15 is a pyridinyl radical or substituted pyridinyl radical, respectively. R¹ and/or R² contains a 6-membered heteroaryl or a substituted 6-membered heteroaryl when, for example, R¹ and/or R² is substituted 5- or 6-membered heteroaryl-(C₁-C₈ alkylenyl) and/or 5- or 6-membered heteroaryl-(G)_m. An unsubstituted 4-pyrido-N-oxide is drawn below for illustration purposes:



25 The phrase "tertiary organic amine" means a trisubstituted nitrogen group wherein the 3 substituents are independently selected from C₁-C₁₂ alkyl, C₃-C₁₂ cycloalkyl, benzyl, or wherein two of the substituents are taken together with the nitrogen atom to which they are bonded to form a 5- or 6-membered, monocyclic heterocycle containing one nitrogen atom and carbon atoms, and the third substituent is selected from C₁-C₁₂ alkyl and benzyl, or wherein the three
30 substituents are taken together with the nitrogen atom to which they are bonded to form a 7- to 12-membered bicyclic heterocycle containing 1 or 2 nitrogen atoms and carbon atoms, and optionally a C=N double bond when 2 nitrogen atoms are present. Illustrative examples of tertiary organic amine include triethylamine,

diisopropylethylamine, benzyl diethylamino, dicyclohexylmethyl-amine,
1,8-diazabicyclo[5.4.0]undec-7-ene (DBU), 1,4-diazabicyclo[2.2.2]octane (TED),
and 1,5-diazabicyclo[4.3.0]non-5-ene.

5 The phrase “pharmaceutical composition” means a composition suitable
for administration in medical or veterinary use.

 The term “admixed” and the phrase “in admixture” are synonymous and
mean in a state of being in a homogeneous or heterogeneous mixture. Preferred is
a homogeneous mixture.

10 The term “patient” means a mammal. Preferred patients are humans, cats,
dogs, cows, horses, pigs, and sheep.

 The term “animal” means a mammal, as defined above. Preferred animals
include humans, cats, dogs, horses, pigs, sheep, cows, monkeys, rats, mice, guinea
pigs, and rabbits.

15 The term “mammal” includes humans, companion animals such as cats
and dogs, primates such as monkeys and chimpanzees, and livestock animals such
as horses, cows, pigs, and sheep.

20 The phrase “livestock animals” as used herein refers to domesticated
quadrupeds, which includes those being raised for meat and various byproducts,
e.g., a bovine animal including cattle and other members of the genus *Bos*, a
porcine animal including domestic swine and other members of the genus *Sus*, an
ovine animal including sheep and other members of the genus *Ovis*, domestic
goats and other members of the genus *Capra*; domesticated quadrupeds being
raised for specialized tasks such as use as a beast of burden, e.g., an equine animal
including domestic horses and other members of the family *Equidae*, genus
25 *Equus*, or for searching and sentinel duty, e.g., a canine animal including domestic
dogs and other members of the genus *Canis*; and domesticated quadrupeds being
raised primarily for recreational purposes, e.g., members of *Equus* and *Canis*, as
well as a feline animal including domestic cats and other members of the family
Felidae, genus *Felis*.

30 The phrase “anticancer effective amount” means an amount of invention
compound, or a pharmaceutically acceptable salt thereof, or a tautomer thereof,
sufficient to inhibit, halt, or cause regression of the cancer being treated in a
particular patient or patient population. For example in humans or other mammals,

an anticancer effective amount can be determined experimentally in a laboratory or clinical setting, or may be the amount required by the guidelines of the United States Food and Drug Administration, or equivalent foreign agency, for the particular cancer and patient being treated.

5 The phrase “anti-arthritic effective amount” means an amount of invention compound, or a pharmaceutically acceptable salt thereof, or a tautomer thereof, sufficient to inhibit, halt, or cause regression of the arthritis being treated in a particular patient or patient population. For example in humans or other mammals, an anti-arthritic effective amount can be determined experimentally in a laboratory
10 or clinical setting, or may be the amount required by the guidelines of the United States Food and Drug Administration, or equivalent foreign agency, for the particular arthritis and patient being treated.

 The phrase “MMP-13 inhibiting amount” means an amount of invention compound, or a pharmaceutically acceptable salt thereof, or a tautomer thereof,
15 sufficient to inhibit an enzyme matrix metalloproteinase-13, including a truncated form thereof, including a catalytic domain thereof, in a particular animal or animal population. For example in a human or other mammal, an MMP-13 inhibiting amount can be determined experimentally in a laboratory or clinical setting, or may be the amount required by the guidelines of the United States Food and Drug
20 Administration, or equivalent foreign agency, for the particular MMP-13 enzyme and patient being treated.


 It should be appreciated that determination of proper dosage forms, dosage amounts, and routes of administration, is within the level of ordinary skill in the pharmaceutical and medical arts, and is described below.

25 The phrases “effective amount” and “therapeutically effective amount” are synonymous and mean an amount of a compound of the present invention, a pharmaceutically acceptable salt thereof, or a solvate thereof, sufficient to effect an improvement of the condition being treated when administered to a patient suffering from a disease that is mediated by MMP-13 and optionally from 0 to
30 12 additional MMP enzymes.

 The term “tautomer” means a form of invention compound existing in a state of equilibrium with an isomeric form of the invention compound, wherein


the invention compound is able to react according to either form by virtue of the ability of the forms to interconvert by isomerization in situ, including in a reaction mixture, in an in vitro biological assay, or in vivo.

The term “(E)” means entgegen, and designates that the conformation about the double bond to which the term refers is the conformation having the two higher ranking substituent groups, as determined according to the Cahn-Ingold-Prelog ranking system, on opposite sides of the double bond. An (E) double bond is illustrated below by the compound of Formula (W)



, wherein the two higher-ranking substituents are groups A and D.

The term “(Z)” means zusammen, and designates that the conformation about the double bond to which the term refers is the conformation having the two higher ranking substituent groups, as determined according to the Cahn-Ingold-Prelog ranking system, on the same side of the double bond. A (Z) double bond is illustrated below by the compound of Formula (X)



, wherein the two higher-ranking substituents are groups A and D.

It should be appreciated that the S1' site of MMP-13 was previously thought to be a grossly linear channel which contained an opening at the top that allowed an amino acid side chain from a substrate molecule to enter during binding, and was closed at the bottom. Applicants has discovered that the S1' site is actually composed of an S1' channel angularly connected to a newly discovered pocket which applicant calls the S1" site. The S1" site is open to solvent at the bottom, which can expose a functional group of Applicants' invention compounds to solvent. For illustrative purposes, the S1' site of the MMP-13 enzyme can now be thought of as being like a sock with a hole in the toes, wherein the S1' channel is the region from approximately the opening to the ankle, and the S1" site is the foot region below the ankle, which foot region is angularly connected to the ankle region.

More particularly, the S1' channel is a specific part of the S1' site and is formed largely by Leu218, Val219, His222 and by residues from Leu239 to Tyr244. The S1" binding site which has been newly discovered is defined by residues from Tyr246 to Pro255. The S1" site contains at least two hydrogen bond donors and aromatic groups which interact with an invention compound.

Without wishing to be bound by any particular theory, the inventors believe that the S1" site could be a recognition site for triple helix collagen, the natural substrate for MMP-13. It is possible that the conformation of the S1" site is modified only when an appropriate compound binds to MMP-13, thereby interfering with the collagen recognition process. This newly discovered pattern of binding offers the possibility of greater selectivity than what is achievable with the binding pattern of known selective inhibitors of MMP-13, wherein the known binding pattern requires ligation of the catalytic zinc atom at the active site and occupation the S1' channel, but not the S1" site.

The term "Thr245" means threonine 245 of an MMP-13 enzyme.

The term "Thr247" means threonine 247 of an MMP-13 enzyme.

The term "Met253" means methionine 253 of an MMP-13 enzyme.

The term "His251" means histidine 251 of an MMP-13 enzyme.

It should be appreciated that the matrix metalloproteinases include, but are not limited to, the following enzymes:

MMP-1, also known as interstitial collagenase, collagenase-1, or fibroblast-type collagenase;

MMP-2, also known as gelatinase A or 72 kDa Type IV collagenase;

MMP-3, also known as stromelysin or stromelysin-1;

MMP-7, also known as matrilysin or PUMP-1;

MMP-8, also known as collagenase-2, neutrophil collagenase or polymorphonuclear-type ("PMN-type") collagenase;

MMP-9, also known as gelatinase B or 92 kDa Type IV collagenase;

MMP-10, also known as stromelysin-2;

MMP-11, also known as stromelysin-3;

MMP-12, also known as metalloelastase;

MMP-13, also known as collagenase-3;

MMP-14, also known as membrane-type ("MT") 1-MMP or MT1-MMP;

MMP-15, also known as MT2-MMP;

MMP-16, also known as MT3-MMP;

MMP-17, also known as MT4-MMP;

MMP-18; and

5 MMP-19.

Other known MMPs include MMP-26 (Matrilysin-2).

For the purposes of this invention, the term "arthritis", which is synonymous with the phrase "arthritic condition", includes osteoarthritis, rheumatoid arthritis, degenerative joint disease, spondyloarthropathies, gouty
10 arthritis, systemic lupus erythematosus, juvenile arthritis, and psoriatic arthritis. An allosteric inhibitor of MMP-13 having an anti-arthritic effect is a compound as defined above that inhibits the progress, prevents further progress, or reverses progression, in part or in whole, of any one or more symptoms of any one of the arthritic diseases and disorders listed above.

15 The term "IC₅₀" means the concentration of a compound, usually expressed as micromolar or nanomolar, required to inhibit an enzyme's catalytic activity by 50%.

The term "ED₄₀" means the concentration of a compound, usually expressed as micromolar or nanomolar, required to treat a disease in about 40% of a patient
20 group.

The term "ED₃₀" means the concentration of a compound, usually expressed as micromolar or nanomolar, required to treat a disease in 30% of a patient group.

The phrase "pharmaceutical composition" means a composition suitable for administration in medical or veterinary use.

25 The term "admixed" and the phrase "in admixture" are synonymous and mean in a state of being in a homogeneous or heterogeneous mixture. Preferred is a homogeneous mixture.

As used herein, the phrase "cartilage damage" means a disorder of hyaline cartilage and subchondral bone characterized by hypertrophy of tissues in and
30 around the involved joints, which may or may not be accompanied by deterioration of hyaline cartilage surface.


The phrase "treating", which is related to the terms "treat" and "treated", means administration of an invention combination as defined above that inhibits

the progress, prevents further progress, or reverses progression, in part or in whole, of any one or more symptoms of any one of the diseases and disorders listed above.


5 The phrase "invention compound" means a compound of Formula I, or a pharmaceutically acceptable salt thereof, as fully defined above.

The term "nontoxic" means the efficacious dose is 10 times or greater than the dose at which a toxic effect is observed in 10% or more of a patient population.

10 The term "celecoxib" means the compound named 4-(5-(4-methylphenyl)-3-(trifluoromethyl)-1H-pyrazol-1-yl)-benzenesulfonamide. Celecoxib is a selective cyclooxygenase-2 ("COX-2") inhibitor currently approved by the FDA for the treatment of osteoarthritis, rheumatoid arthritis, and Polyposis-familial adenomatus. Celecoxib is marketed under the tradename "Celebrex". Celecoxib is currently in clinical trials for the treatment of bladder cancer, chemopreventative-
15 lung cancer, and post-operative pain, and is registered for the treatment of dysmenorrhea. Celecoxib has the structure drawn below:



20 The term "valdecoxib" means the compound named 4-(5-methyl-3-phenyl-4-isoxazolyl)-benzenesulfonamide. Valdecoxib is a selective COX-2 inhibitor that has been approved by the FDA for treating osteoarthritis, rheumatoid arthritis, dysmenorrhea, and general pain, and is marketed under the tradename "Bextra". Valdecoxib is in clinical trials for the treatment of migraine. Valdecoxib has the structure drawn below:



It should be appreciated that COX-2 is also known as prostaglandin synthase-2 and prostaglandin PGH₂ synthase.

A selective inhibitor of COX-2 means compounds that inhibit COX-2 selectively versus COX-1 such that a ratio of IC₅₀ for a compound with COX-1 divided by a ratio of IC₅₀ for the compound with COX-2 is greater than, or equal to, 5, where the ratios are determined in one or more assays. All that is required to determine whether a compound is a selective COX-2 inhibitor is to assay a compound in one of a number of well know assays in the art.

The term “NSAID” is an acronym for the phrase “nonsteroidal anti-inflammatory drug”, which means any compound which inhibits cyclooxygenase-1 (“COX-1”) and cyclooxygenase-2. Most NSAIDs fall within one of the following five structural classes: (1) propionic acid derivatives, such as ibuprofen, naproxen, naprosyn, diclofenac, and ketoprofen; (2) acetic acid derivatives, such as tolmetin and sulindac; (3) fenamic acid derivatives, such as mefenamic acid and meclofenamic acid; (4) biphenylcarboxylic acid derivatives, such as diflunisal and flufenisal; and (5) oxicams, such as piroxim, peroxicam, sudoxicam, and isoxicam. Other useful NSAIDs include aspirin, acetaminophen, indomethacin, and phenylbutazone. Selective inhibitors of cyclooxygenase-2 as described above may be considered to be NSAIDs also.

The term “drugs”, which is synonymous with the phrases “active components”, “active compounds”, and “active ingredients”, includes celecoxib, or a pharmaceutically acceptable salt thereof, valdecoxib, or a pharmaceutically acceptable salt thereof, and an allosteric inhibitor of MMP-13, and may further include one or two of the other therapeutic agents described above.

The compounds of Formula I, or pharmaceutically acceptable salts thereof, or tautomers thereof, include compounds which are invention compounds. An allosteric inhibitor of MMP-13 is any compound of Formula I that binds allosterically into the S1' site of the MMP-13 enzyme, including the S1' channel, and a newly discovered S1" site, without ligating, coordinating, or binding the catalytic zinc of the MMP-13.

An invention compound that is an allosteric inhibitor of MMP-13 may be readily identified by one of ordinary skill in the pharmaceutical or medical arts by assaying an alkyne test compound for inhibition of MMP-13 as described below in Biological Methods 1 or 2, and for allosteric inhibition of MMP-13 by assaying the test invention compound for inhibition of MMP-13 in the presence of an inhibitor to the catalytic zinc of MMP-13 as described below in Biological Methods 3 or 4.

Further, an invention compound having an anti-inflammatory, an analgesic, anti-arthritic, or a cartilage damage inhibiting effect, or any combination of these effects, may be readily identified by one of ordinary skill in the pharmaceutical or medical arts by assaying the invention compound in any number of well known assays for measuring determining the invention compound's effects on cartilage damage, arthritis, inflammation, or pain. These assays include in vitro assays that utilize cartilage samples and in vivo assays in whole animals that measure cartilage degradation, inhibition of inflammation, or pain alleviation.

For example with regard to assaying cartilage damage in vitro, an amount of an invention compound or control vehicle may be administered with a cartilage damaging agent to cartilage, and the cartilage damage inhibiting effects in both tests studied by gross examination or histopathologic examination of the cartilage, or by measurement of biological markers of cartilage damage such as, for example, proteoglycan content or hydroxyproline content. Further, in vivo assays to assay cartilage damage may be performed as follows: an amount of an invention compound or control vehicle may be administered with a cartilage damaging agent to an animal, and the effects of the invention compound being assayed on cartilage in the animal may be evaluated by gross examination or histopathologic examination of the cartilage, by observation of the effects in an

acute model on functional limitations of the affected joint that result from cartilage damage, or by measurement of biological markers of cartilage damage such as, for example, proteoglycan content or hydroxyproline content.

5 Several methods of identifying an invention compound with cartilage damage inhibiting properties are described below. The amount to be administered in an assay is dependent upon the particular assay employed, but in any event is not higher than the well known maximum amount of a compound that the particular assay can effectively accommodate.

10 Similarly, invention compounds having pain-alleviating properties may be identified using any one of a number of in vivo animal models of pain.

 Still similarly, invention compounds having anti-inflammatory properties may be identified using any one of a number of in vivo animal models of inflammation. For example, for an example of inflammation models, see United States patent number 6, 329,429, which is incorporated herein by reference.

15 Still similarly, invention compounds having anti-arthritis properties may be identified using any one of a number of in vivo animal models of arthritis. For example, for an example of arthritis models, see also United States patent number 6, 329,429.

20 Other mammalian diseases and disorders which are treatable by administration of an invention combination alone, or contained in a pharmaceutical composition as defined below, include: fever (including rheumatic fever and fever associated with influenza and other viral infections), common cold, dysmenorrhea, menstrual cramps, inflammatory bowel disease, Crohn's disease, emphysema, acute respiratory distress syndrome, asthma, bronchitis, chronic
25 obstructive pulmonary disease, Alzheimer's disease, organ transplant toxicity, cachexia, allergic reactions, allergic contact hypersensitivity, cancer (such as solid tumor cancer including colon cancer, breast cancer, lung cancer and prostate cancer; hematopoietic malignancies including leukemias and lymphomas; Hodgkin's disease; aplastic anemia, skin cancer and familial adenomatous
30 polyposis), tissue ulceration, peptic ulcers, gastritis, regional enteritis, ulcerative colitis, diverticulitis, recurrent gastrointestinal lesion, gastrointestinal bleeding, coagulation, anemia, synovitis, gout, ankylosing spondylitis, restenosis, periodontal disease, epidermolysis bullosa, osteoporosis, loosening of artificial joint implants,

atherosclerosis (including atherosclerotic plaque rupture), aortic aneurysm (including abdominal aortic aneurysm and brain aortic aneurysm), periarteritis nodosa, congestive heart failure, myocardial infarction, stroke, cerebral ischemia, head trauma, spinal cord injury, neuralgia, neuro-degenerative disorders (acute and chronic), autoimmune disorders, Huntington's disease, Parkinson's disease, migraine, depression, peripheral neuropathy, pain (including low back and neck pain, headache and toothache), gingivitis, cerebral amyloid angiopathy, nootropic or cognition enhancement, amyotrophic lateral sclerosis, multiple sclerosis, ocular angiogenesis, corneal injury, macular degeneration, conjunctivitis, abnormal wound healing, muscle or joint sprains or strains, tendonitis, skin disorders (such as psoriasis, eczema, scleroderma and dermatitis), myasthenia gravis, polymyositis, myositis, bursitis, burns, diabetes (including types I and II diabetes, diabetic retinopathy, neuropathy and nephropathy), tumor invasion, tumor growth, tumor metastasis, corneal scarring, scleritis, immunodeficiency diseases (such as AIDS in humans and FLV, FIV in cats), sepsis, premature labor, hypoprothrombinemia, hemophilia, thyroiditis, sarcoidosis, Behcet's syndrome, hypersensitivity, kidney disease, Rickettsial infections (such as Lyme disease, Erlichiosis), Protozoan diseases (such as malaria, giardia, coccidia), reproductive disorders (preferably in livestock), epilepsy, convulsions, and septic shock.

Other aspects of the present invention are compounds of Formula I, or a pharmaceutically acceptable salt thereof, that are ≥ 10 , ≥ 20 , ≥ 50 , ≥ 100 , or ≥ 1000 times more potent versus MMP-13 than versus at least two of any other MMP enzyme or TACE.

Still other aspects of the present invention are compounds of Formula I, or a pharmaceutically acceptable salt thereof, that are selective inhibitors of MMP-13 versus 2, 3, 4, 5, 6, or 7 other MMP enzymes, or versus TACE and 1, 2, 3, 4, 5, 6, or 7 other MMP enzymes.

It should be appreciated that selectivity of a compound of Formula I, or a pharmaceutically acceptable salt thereof, is a multidimensional characteristic that includes the number of other MMP enzymes and TACE over which selectivity for MMP-13 inhibition is present and the degree of selectivity of inhibition of MMP-13 over another particular MMP or TACE, as measured by, for example, the IC_{50}

in micromolar concentration of the compound for the inhibition of the other MMP enzyme or TACE divided by the IC_{50} in micromolar concentration of the compound for the inhibition of MMP-13.

As discussed above, one aspect of the present invention is novel
5 compounds that are selective inhibitors of the enzyme MMP-13. A selective inhibitor of MMP-13, as used in the present invention, is a compound that is $\geq 5X$ more potent *in vitro* versus MMP-13 than versus at least one other matrix metalloproteinase enzyme such as, for example, MMP-1, MMP-2, MMP-3, MMP-7, MMP-8, MMP-9, or MMP-14, or versus tumor necrosis factor alpha
10 convertase ("TACE"). A preferred aspect of the present invention is novel compounds that are selective inhibitors of MMP-13 versus MMP-1.

The invention provides a compound of Formula I, or a pharmaceutically acceptable salt thereof, which has an IC_{50} with any MMP enzyme that is less than or equal to 50 micromolar. Preferred are compounds of Formula I, or a
15 pharmaceutically acceptable salt thereof, which have an IC_{50} with a human full-length MMP-13 ("hMMP-13FL") or a human MMP-13 catalytic domain ("hMMP-13CD") that is less than or equal to 50 micromolar. More preferred are compounds of Formula I, or a pharmaceutically acceptable salt thereof, which have an IC_{50} with a human full-length MMP-13 ("hMMP-13FL") or a human
20 MMP-13 catalytic domain ("hMMP-13CD") that is less than or equal to 10 micromolar.

The compound named 4-[1-oxo-7-(3-[1,2,3]triazol-1-yl)prop-1-ynyl]-1H-isoquinolin-2-ylmethyl]benzoic acid is excluded from the invention because it was assayed with hMMP-13CD and was found to have an IC_{50} that exceeded 50
25 micromolar.

Examples of biological methods useful for determining IC_{50} s for the invention compounds with an MMP are described below in Biological Methods 1 to 4. Any compound of Formula I, or a pharmaceutically acceptable salt thereof, or any form thereof as defined above, that does not have an IC_{50} with any MMP
30 enzyme that is less than, or equal to, 10 micromolar is excluded from this invention.

Some of the invention compounds are capable of further forming nontoxic pharmaceutically acceptable salts, including, but not limited to, acid addition

and/or base salts. The acid addition salts are formed from basic invention compounds, whereas the base addition salts are formed from acidic invention compounds. All of these forms are within the scope of the compounds useful in the invention.

5 Pharmaceutically acceptable acid addition salts of the basic invention compounds include nontoxic salts derived from inorganic acids such as hydrochloric, nitric, phosphoric, sulfuric, hydrobromic, hydroiodic, hydrofluoric, phosphorous, and the like, as well nontoxic salts derived from organic acids, such as aliphatic mono- and dicarboxylic acids, phenyl-substituted alkanolic acids,
10 hydroxy alkanolic acids, alkanedioic acids, aromatic acids, aliphatic and aromatic sulfonic acids, etc. Such salts thus include sulfate, pyrosulfate, bisulfate, sulfite, bisulfite, nitrate, phosphate, monohydrogenphosphate, dihydrogenphosphate, metaphosphate, pyrophosphate, chloride, bromide, iodide, acetate, trifluoroacetate, propionate, caprylate, isobutyrate, oxalate, malonate, succinate,
15 suberate, sebacate, fumarate, maleate, mandelate, benzoate, chlorobenzoate, methylbenzoate, dinitrobenzoate, phthalate, benzenesulfonate, toluenesulfonate, phenylacetate, citrate, lactate, malate, tartrate, methanesulfonate, and the like. Also contemplated are salts of amino acids such as arginate and the like and gluconate, galacturonate (see, for example, Berge S.M. et al., "Pharmaceutical
20 Salts," J. of Pharma. Sci., 1977;66:1).

 An acid addition salt of a basic invention compound is prepared by contacting the free base form of the compound with a sufficient amount of a desired acid to produce a nontoxic salt in the conventional manner. The free base form of the compound may be regenerated by contacting the acid addition salt so
25 formed with a base, and isolating the free base form of the compound in the conventional manner. The free base forms of compounds prepared according to a process of the present invention differ from their respective acid addition salt forms somewhat in certain physical properties such as solubility, crystal structure, hygroscopicity, and the like, but otherwise free base forms of the invention
30 compounds and their respective acid addition salt forms are equivalent for purposes of the present invention.

 A nontoxic pharmaceutically acceptable base addition salt of an acidic invention compound may be prepared by contacting the free acid form of the

compound with a metal cation such as an alkali or alkaline earth metal cation, or an amine, especially an organic amine. Examples of suitable metal cations include sodium cation (Na^+), potassium cation (K^+), magnesium cation (Mg^{2+}), calcium cation (Ca^{2+}), and the like. Examples of suitable amines are
5 N,N'-dibenzylethylenediamine, chlorprocaine, choline, diethanolamine, dicyclohexylamine, ethylenediamine, N-methylglucamine, and procaine (see, for example, Berge, supra., 1977).

A base addition salt of an acidic invention compound may be prepared by contacting the free acid form of the compound with a sufficient amount of a
10 desired base to produce the salt in the conventional manner. The free acid form of the compound may be regenerated by contacting the salt form so formed with an acid, and isolating the free acid of the compound in the conventional manner. The free acid forms of the invention compounds differ from their respective salt forms somewhat in certain physical properties such as solubility, crystal structure,
15 hygroscopicity, and the like, but otherwise the salts are equivalent to their respective free acid for purposes of the present invention.

Certain invention compounds can exist in unsolvated forms as well as solvated forms, including hydrated forms. In general, the solvated forms, including hydrated forms, are equivalent to unsolvated forms and are
20 encompassed within the scope of the present invention.

Certain of the invention compounds possess one or more chiral centers, and each center may exist in the R or S configuration. An invention compound includes any diastereomeric, enantiomeric, or epimeric form of the compound, as well as mixtures thereof.

25 Additionally, certain invention compounds may exist as geometric isomers such as the entgegen (E) and zusammen (Z) isomers of 1,2-disubstituted alkenyl groups or cis and trans isomers of disubstituted cyclic groups. An invention compound includes any cis, trans, syn, anti, entgegen (E), or zusammen (Z) isomer of the compound, as well as mixtures thereof.

30 Certain invention compounds can exist as two or more tautomeric forms. Tautomeric forms of the invention compounds may interchange, for example, via enolization/de-enolization, 1,2-hydride, 1,3-hydride, or 1,4-hydride shifts, and the

like. An invention compound includes any tautomeric form of the compound, as well as mixtures thereof.

Some compounds of the present invention have alkenyl groups, which may exist as *entgegen* or *zusammen* conformations, in which case all geometric forms thereof, both *entgegen* and *zusammen*, *cis* and *trans*, and mixtures thereof, are within the scope of the present invention.

Some compounds of the present invention have cycloalkyl groups, which may be substituted at more than one carbon atom, in which case all geometric forms thereof, both *cis* and *trans*, and mixtures thereof, are within the scope of the present invention.

It should be appreciated that in the Summary of the Invention above, the term "Embodiment" refers to an aspect of this invention. The Embodiments in the Summary of the Invention are numbered for ease of referral.

The invention compounds also include isotopically-labelled compounds, which are identical to those recited above, but for the fact that one or more atoms are replaced by an atom having an atomic mass or mass number different from the atomic mass or mass number usually found in nature. Examples of isotopes that can be incorporated into compounds of the invention include isotopes of hydrogen, carbon, nitrogen, oxygen, phosphorous, fluorine and chlorine, such as ^2H , ^3H , ^{13}C , ^{14}C , ^{15}N , ^{18}O , ^{17}O , ^{31}P , ^{32}P , ^{35}S , ^{18}F and ^{36}Cl , respectively. Compounds of the present invention and pharmaceutically acceptable salts of said compounds which contain the aforementioned isotopes and/or other isotopes of other atoms are within the scope of this invention. Certain isotopically labelled compounds of the present invention, for example those into which radioactive isotopes such as ^3H and ^{14}C are incorporated, are useful in drug and/or substrate tissue distribution assays. Tritiated, i.e., ^3H and carbon-14, i.e., ^{14}C , isotopes are particularly preferred for their ease of preparation and detectability. Further, substitution with heavier isotopes such as deuterium, i.e., ^2H , can afford certain therapeutic advantages resulting from greater metabolic stability, for example increased in vivo half-life or reduced dosage requirements and, hence, may be preferred in some circumstances. Isotopically labelled compounds of those described above in this invention can generally be prepared by carrying out the procedures

incorporated by reference above or disclosed in the Schemes and/or in the Examples and Preparations below, by substituting a readily available isotopically labelled reagent for a non-isotopically labelled reagent.

5 All of the above-describe forms of an invention compound are included by the phrase “invention compound”, a “compound of Formula I”, a “compound of Formula I, or a pharmaceutically acceptable salt thereof”, or any named species thereof, unless specifically excluded therefrom.

10 One of ordinary skill in the art will appreciate that the compounds of the invention are useful in treating a diverse array of diseases. One of ordinary skill in the art will also appreciate that when using the compounds of the invention in the treatment of a specific disease that the compounds of the invention may be combined with various existing therapeutic agents used for that disease.

15 For the treatment of rheumatoid arthritis, the compounds of the invention may be combined with agents such as TNF- α inhibitors such as anti-TNF monoclonal antibodies and TNF receptor immunoglobulin molecules (such as Enbrel®), low dose methotrexate, lefunimide, hydroxychloroquine, d-penicillamine, auranofin or parenteral or oral gold.

20 The compounds of the invention can also be used in combination with existing therapeutic agents for the treatment of osteoarthritis. Suitable agents to be used in combination include standard non-steroidal anti-inflammatory agents (hereinafter NSAID's) such as piroxicam, diclofenac, propionic acids such as naproxen, flurbiprofen, fenoprofen, ketoprofen and ibuprofen, fenamates such as mefenamic acid, indomethacin, sulindac, apazone, pyrazolones such as phenylbutazone, salicylates such as aspirin, COX-2 inhibitors such as etoricoxib
25 and rofecoxib, analgesics and intraarticular therapies such as corticosteroids and hyaluronic acids such as hyalgan and synvisc.

30 This invention also relates to a method of or a pharmaceutical composition for treating inflammatory processes and diseases comprising administering a compound of this invention to a mammal, including a human, cat, livestock or dog, wherein said inflammatory processes and diseases are defined as above and said inhibitory compound is used in combination with one or more other therapeutically active agents under the following conditions:

A.) where a joint has become seriously inflamed as well as infected at the same time by bacteria, fungi, protozoa and/or virus, said inhibitory compound is administered in combination with one or more antibiotic, antifungal, antiprotozoal and/or antiviral therapeutic agents;

5 B.) where a multi-fold treatment of pain and inflammation is desired, said inhibitory compound is administered in combination with inhibitors of other mediators of inflammation, comprising one or more members independently selected from the group consisting essentially of:

- (1) NSAIDs;
- 10 (2) H₁-receptor antagonists;
- (3) kinin-B₁ - and B₂-receptor antagonists;
- (4) prostaglandin inhibitors selected from the group consisting of PGD-, PGF-, PGI₂ - and PGE-receptor antagonists;
- (5) thromboxane A₂ (TXA₂-) inhibitors;
- 15 (6) 5-, 12- and 15-lipoxygenase inhibitors;
- (7) leukotriene LTC₄ -, LTD₄/LTE₄ - and LTB₄ -inhibitors;
- (8) PAF-receptor antagonists;
- (9) gold in the form of an aurothio group together with one or more hydrophilic groups;
- 20 (10) immunosuppressive agents selected from the group consisting of cyclosporine, azathioprine and methotrexate;
- (11) anti-inflammatory glucocorticoids;
- (12) penicillamine;
- (13) hydroxychloroquine;
- 25 (14) anti-gout agents including colchicine; xanthine oxidase inhibitors including allopurinol; and uricosuric agents selected from probenecid, sulfinpyrazone and benzbromarone;

C. where older mammals are being treated for disease conditions, syndromes and symptoms found in geriatric mammals, said inhibitory compound is administered in combination with one or more members independently selected from the group consisting essentially of:

- (1) cognitive therapeutics to counteract memory loss and impairment;

(2) anti-hypertensives and other cardiovascular drugs intended to offset the consequences of atherosclerosis, hypertension, myocardial ischemia, angina, congestive heart failure and myocardial infarction, selected from the group consisting of:

- 5 a. diuretics;
- b. vasodilators;
- c. β -adrenergic receptor antagonists;
- d. angiotensin-II converting enzyme inhibitors (ACE-inhibitors), alone or optionally together with neutral endopeptidase inhibitors;
- 10 e. angiotensin II receptor antagonists;
- f. renin inhibitors;
- g. calcium channel blockers;
- h. sympatholytic agents;
- i. α_2 -adrenergic agonists;
- 15 j. α -adrenergic receptor antagonists; and
- k. HMG-CoA-reductase inhibitors (anti-hypercholesterolemics);
- (3) antineoplastic agents selected from:
 - a. antimitotic drugs selected from:
 - i. vinca alkaloids selected from:
- 20 [1] vinblastine and
- [2] vincristine;
- (4) growth hormone secretagogues;
- (5) strong analgesics;
- (6) local and systemic anesthetics; and
- 25 (7) H_2 -receptor antagonists, proton pump inhibitors and other gastroprotective agents.

The active ingredient of the present invention may be administered in combination with inhibitors of other mediators of inflammation, comprising one or more members selected from the group consisting essentially of the classes of

30 such inhibitors and examples thereof which include, matrix metalloproteinase inhibitors, aggrecanase inhibitors, TACE inhibitors, leucotriene receptor antagonists, IL-1 processing and release inhibitors, ILra, H_1 -receptor antagonists;

kinin-B₁ - and B₂ -receptor antagonists; prostaglandin inhibitors such as PGD-, PGF- PGI₂ - and PGE-receptor antagonists; thromboxane A₂ (TXA₂-) inhibitors; 5- and 12-lipoxygenase inhibitors; leukotriene LTC₄ -, LTD₄/LTE₄ - and LTB₄ - inhibitors; PAF-receptor antagonists; gold in the form of an aurothio group together with various hydrophilic groups; immunosuppressive agents, e.g., cyclosporine, azathioprine and methotrexate; anti-inflammatory glucocorticoids; penicillamine; hydroxychloroquine; anti-gout agents, e.g., colchicine, xanthine oxidase inhibitors, e.g., allopurinol and uricosuric agents, e.g., probenecid, sulfinpyrazone and benzbromarone.

10 The compounds of the present invention may also be used in combination with anticancer agents such as endostatin and angiostatin or cytotoxic drugs such as adriamycin, daunomycin, cis-platinum, etoposide, taxol, taxotere and alkaloids, such as vincristine and antimetabolites such as methotrexate.

15 The compounds of the present invention may also be used in combination with anti-hypertensives and other cardiovascular drugs intended to offset the consequences of atherosclerosis, including hypertension, myocardial ischemia including angina, congestive heart failure and myocardial infarction, selected from vasodilators such as hydralazine, β -adrenergic receptor antagonists such as propranolol, calcium channel blockers such as nifedipine, α_2 -adrenergic agonists
20 such as clonidine, α -adrenergic receptor antagonists such as prazosin and HMG-CoA-reductase inhibitors (anti-hypercholesterolemics) such as lovastatin or atorvastatin.

25 The compounds of the present invention may also be administered in combination with one or more antibiotic, antifungal, antiprotozoal, antiviral or similar therapeutic agents.

30 The compounds of the present invention may also be used in combination with CNS agents such as antidepressants (such as sertraline), anti-Parkinsonian drugs (such as L-dopa, requip, mirapex, MAOB inhibitors such as selegine and rasagiline, comP inhibitors such as Tasmar, A-2 inhibitors, dopamine reuptake inhibitors, NMDA antagonists, nicotine agonists, dopamine agonists and inhibitors of neuronal nitric oxide synthase) and anti-Alzheimer's drugs such as donepezil, tacrine, COX-2 inhibitors, propentofylline or metryfonate.

The compounds of the present invention may also be used in combination with osteoporosis agents such as roloxifene, lasofoxifene, droloxifene or fosomax and immunosuppressant agents such as FK-506 and rapamycin.

5 The invention compounds may be used in combination with a COX-2 selective inhibitor, more preferably celecoxib (e.g., CELEBREX®), valdecoxib (e.g., BEXTRA®), parecoxib, lumiracoxib (e.g., PREXIGE®), or rofecoxib (e.g., VIOXX®), or with compounds such as etanercept (e.g., ENBREL®), infliximab (e.g., REMICADE®), leflunomide, (e.g., ARAVA®) or methotrexate, and the like.

10 The invention compounds may be used in combination with biological therapeutics useful for treating arthritic conditions, including CP-870, etanercept (a tumor necrosis factor alpha ("TNF-alpha") receptor immunoglobulin molecule; trade names ENBREL® and ENBREL ENTANERCEPT® by Immunex Corporation, Seattle, Washington), infliximab (an anti-TNF-alpha chimeric IgG
15 1K monoclonal antibody; tradename REMICADE® by Centocor, Inc., Malvern, Pennsylvania), methotrexate (tradename RHEUMATREX® by American Cyanamid Company, Wayne, New Jersey), and adalimumab (a human monoclonal anti-TNF-alpha antibody; tradename HUMIRA® by Abbott Laboratories, Abbott Park, Illinois).

20 The present invention also relates to the formulation of a compound of the present invention alone or with one or more other therapeutic agents which are to form the intended combination, including wherein said different drugs have varying half-lives, by creating controlled-release forms of said drugs with different release times which achieves relatively uniform dosing; or, in the case of
25 non-human patients, a medicated feed dosage form in which said drugs used in the combination are present together in admixture in the feed composition. There is further provided in accordance with the present invention co-administration in which the combination of drugs is achieved by the simultaneous administration of said drugs to be given in combination; including co-administration by means of
30 different dosage forms and routes of administration; the use of combinations in accordance with different but regular and continuous dosing schedules whereby desired plasma levels of said drugs involved are maintained in the patient being

treated, even though the individual drugs making up said combination are not being administered to said patient simultaneously.

The invention method is useful in human and veterinary medicines for treating mammals suffering from one or more of the above-listed diseases and disorders.

All that is required to practice a method of this invention is to administer a compound of Formula I, or a pharmaceutically acceptable salt thereof, in an amount that is therapeutically effective for preventing, inhibiting, or reversing the condition being treated. The invention compound can be administered directly or in a pharmaceutical composition as described below.

A therapeutically effective amount, or, simply, effective amount, of an invention compound will generally be from about 1 to about 300 mg/kg of subject body weight of the compound of Formula I, or a pharmaceutically acceptable salt thereof. Typical doses will be from about 10 to about 5000 mg/day for an adult subject of normal weight for each component of the combination. In a clinical setting, regulatory agencies such as, for example, the Food and Drug Administration ("FDA") in the U.S. may require a particular therapeutically effective amount.

In determining what constitutes a nontoxic effective amount or a therapeutically effective amount of an invention compound for treating, preventing, or reversing one or more symptoms of any one of the diseases and disorders described above that are being treated according to the invention methods, a number of factors will generally be considered by the medical practitioner or veterinarian in view of the experience of the medical practitioner or veterinarian, including the Food and Drug Administration guidelines, or guidelines from an equivalent agency, published clinical studies, the subject's (e.g., mammal's) age, sex, weight and general condition, as well as the type and extent of the disease, disorder or condition being treated, and the use of other medications, if any, by the subject. As such, the administered dose may fall within the ranges or concentrations recited above, or may vary outside them, ie, either below or above those ranges, depending upon the requirements of the individual subject, the severity of the condition being treated, and the particular therapeutic formulation being employed. Determination of a proper dose for a particular

situation is within the skill of the medical or veterinary arts. Generally, treatment may be initiated using smaller dosages of the invention compound that are less than optimum for a particular subject. Thereafter, the dosage can be increased by small increments until the optimum effect under the circumstance is reached. For
5 convenience, the total daily dosage may be divided and administered in portions during the day, if desired.

Pharmaceutical compositions, described briefly here and more fully below, of an invention combination may be produced by formulating the invention combination in dosage unit form with a pharmaceutical carrier. Some examples of
10 dosage unit forms are tablets, capsules, pills, powders, aqueous and nonaqueous oral solutions and suspensions, and parenteral solutions packaged in containers containing either one or some larger number of dosage units and capable of being subdivided into individual doses. Alternatively, the invention compounds may be formulated separately.

Some examples of suitable pharmaceutical carriers, including pharmaceutical diluents, are gelatin capsules; sugars such as lactose and sucrose; starches such as corn starch and potato starch; cellulose derivatives such as sodium carboxymethyl cellulose, ethyl cellulose, methyl cellulose, and cellulose acetate phthalate; gelatin; talc; stearic acid; magnesium stearate; vegetable oils
15 such as peanut oil, cottonseed oil, sesame oil, olive oil, corn oil, and oil of theobroma; propylene glycol, glycerin; sorbitol; polyethylene glycol; water; agar; alginic acid; isotonic saline, and phosphate buffer solutions; as well as other compatible substances normally used in pharmaceutical formulations.

The compositions to be employed in the invention can also contain other
25 components such as coloring agents, flavoring agents, and/or preservatives. These materials, if present, are usually used in relatively small amounts. The compositions can, if desired, also contain other therapeutic agents commonly employed to treat any of the above-listed diseases and disorders.

The percentage of the active ingredients of a compound of Formula I, or a
30 pharmaceutically acceptable salt thereof, in the foregoing compositions can be varied within wide limits, but for practical purposes it is preferably present in a total concentration of at least 10% in a solid composition and at least 2% in a primary liquid composition. The most satisfactory compositions are those in

which a much higher proportion of the active ingredients are present, for example, up to about 95%.

Preferred routes of administration of an invention compound are oral or parenteral. However, another route of administration may be preferred depending upon the condition being treated. For example, topical administration or administration by injection may be preferred for treating conditions localized to the skin or a joint. Administration by transdermal patch may be preferred where, for example, it is desirable to effect sustained dosing.

It should be appreciated that the different routes of administration may require different dosages. For example, a useful intravenous ("IV") dose is between 5 and 50 mg, and a useful oral dosage is between 20 and 800 mg, of a compound of Formula I, or a pharmaceutically acceptable salt thereof. The dosage is within the dosing range used in treatment of the above-listed diseases, or as would be determined by the needs of the patient as described by the physician.

The invention compounds may be administered in any form. Preferably, administration is in unit dosage form. A unit dosage form of the invention compound to be used in this invention may also comprise other compounds useful in the therapy of diseases described above. A further description of pharmaceutical formulations useful for administering the invention compounds and invention combinations is provided below.

The active components of the invention combinations, may be formulated together or separately and may be administered together or separately. The particular formulation and administration regimens used may be tailored to the particular patient and condition being treated by a practitioner of ordinary skill in the medical or pharmaceutical arts.

The advantages of using an invention compound in a method of the instant invention include the nontoxic nature of the compounds at and substantially above therapeutically effective doses, their ease of preparation, the fact that the compounds are well-tolerated, and the ease of topical, IV, or oral administration of the drugs.

Another important advantage is that the present invention compounds more effectively target a particular disease that is responsive to inhibition of MMP-13 with fewer undesirable side effects than similar compounds that inhibit

MMP-13 that are not invention compounds. This is so because the instant invention compounds of Formula I, or a pharmaceutically acceptable salt thereof, do not directly, or indirectly via a bridging water molecule, ligate, coordinate to, or bind to the catalytic zinc cation of MMP-13, but instead bind at a different location from where natural substrate binds to MMP-13. The binding requirements of an allosteric MMP-13 binding site are unique to MMP-13, and account for the specificity of the invention compounds for inhibiting MMP-13 over any other MMP enzyme. This binding mode has not been reported in the art. Indeed, prior art inhibitors of MMP-13 bind to the catalytic zinc cations of other MMP enzymes as well as to the catalytic zinc cation of MMP-13, and are consequently significantly less selective inhibitors of MMP-13 enzyme.

The invention compounds which are invention compounds, and pharmaceutically acceptable salts thereof, are thus therapeutically superior to other inhibitors of MMP-13, or even tumor necrosis factor-alpha converting enzyme ("TACE"), because of fewer undesirable side effects from inhibition of the other MMP enzymes or TACE. For example, virtually all prior art MMP inhibitors tested clinically to date have exhibited an undesirable side effect known as musculoskeletal syndrome ("MSS"). MSS is associated with administering an inhibitor of multiple MMP enzymes or an inhibitor of a particular MMP enzyme such as MMP-1. MSS will be significantly reduced in type and severity by administering the invention compound instead of any prior art MMP-13 inhibitor, or a pharmaceutically acceptable salt thereof. The invention compounds are superior to similar compounds that interact with the catalytic zinc cation of the MMP-13 enzyme as discussed above, even if similar compounds show some selectivity for the MMP-13.

It is expected that nearly all, if not all, compounds of Formula I, or pharmaceutically acceptable salts thereof, are invention compounds.

This advantage of the instant compounds will also significantly increase the likelihood that agencies which regulate new drug approvals, such as the United States Food and Drug Administration, will approve the instant compounds versus a competing similar compound that does not allosterically bind to MMP-13 as discussed above even in the unlikely event that the two compounds behaved similarly in clinical trials. These regulatory agencies are increasingly aware that

clinical trials, which test drug in limited population groups, do not always uncover safety problems with a drug, and thus all other things being equal, the agencies will favor the drug with the lowest odds of producing undesirable side effects.

Another important advantage is that the disease modifying properties of the invention compounds provide patients suffering from cartilage damage, arthritis, preferably osteoarthritis, inflammation and/or pain with both relief of symptoms and prevention or inhibition of the underlying disease pathology such as cartilage degradation. There is no currently approved drug for disease modification of cartilage damage, including in osteoarthritis.

Any invention compound is readily available, either commercially, or by synthetic methodology, well known to those skilled in the art of organic chemistry. For specific syntheses, see the examples below and the preparations of invention compound outlined in the Schemes below.

Intermediates for the synthesis of a compound of Formula I, or a pharmaceutically acceptable salt thereof, may be prepared by one of ordinary skill in the art of organic chemistry by adapting various synthetic procedures incorporated by reference above or that are well-known in the art of organic chemistry. These synthetic procedures may be found in the literature in, for example, Reagents for Organic Synthesis, by Fieser and Fieser, John Wiley & Sons, Inc, New York, 2000; Comprehensive Organic Transformations, by Richard C. Larock, VCH Publishers, Inc, New York, 1989; the series Compendium of Organic Synthetic Methods, 1989, by Wiley-Interscience; the text Advanced Organic Chemistry, 4th edition, by Jerry March, Wiley-Interscience, New York, 1992; or the Handbook of Heterocyclic Chemistry by Alan R. Katritzky, Pergamon Press Ltd, London, 1985, to name a few. Alternatively, a skilled artisan may find methods useful for preparing the intermediates in the chemical literature by searching widely available databases such as, for example, those available from the Chemical Abstracts Service, Columbus, Ohio, or MDL Information Systems GmbH (formerly Beilstein Information Systems GmbH), Frankfurt, Germany.

Preparations of the invention compounds may use starting materials, reagents, solvents, and catalysts that may be purchased from commercial sources

or they may be readily prepared by adapting procedures in the references or resources cited above. Commercial sources of starting materials, reagents, solvents, and catalysts useful in preparing invention compounds include, for example, The Aldrich Chemical Company, and other subsidiaries of Sigma-
5 Aldrich Corporation, St. Louis, Missouri, BACHEM, BACHEM A.G., Switzerland, or Lancaster Synthesis Ltd, United Kingdom.

Syntheses of some invention compounds may utilize starting materials, intermediates, or reaction products that contain a reactive functional group. During chemical reactions, a reactive functional group may be protected from
10 reacting by a protecting group that renders the reactive functional group substantially inert to the reaction conditions employed. A protecting group is introduced onto a starting material prior to carrying out the reaction step for which a protecting group is needed. Once the protecting group is no longer needed, the protecting group can be removed. It is well within the ordinary skill in the art to
15 introduce protecting groups during a synthesis of a compound of Formula I, or a pharmaceutically acceptable salt thereof, and then later remove them. Procedures for introducing and removing protecting groups are known and referenced such as, for example, in Protective Groups in Organic Synthesis, 2nd ed., Greene T.W. and Wuts P.G., John Wiley & Sons, New York: New York, 1991, which is hereby
20 incorporated by reference.


Thus, for example, protecting groups such as the following may be utilized to protect amino, hydroxyl, and other groups: carboxylic acyl groups such as, for example, formyl, acetyl, and trifluoroacetyl; alkoxycarbonyl groups such as, for example, ethoxycarbonyl, tert-butoxycarbonyl (BOC), β,β,β -
25 trichloroethoxycarbonyl (TCEC), and β -iodoethoxycarbonyl; aralkyloxycarbonyl groups such as, for example, benzyloxycarbonyl (CBZ), paramethoxybenzyloxycarbonyl, and 9-fluorenylmethyloxycarbonyl (Fmoc); trialkylsilyl groups such as, for example, trimethylsilyl (TMS) and tert-butyltrimethylsilyl (TBDMS); and other groups such as, for example,
30 triphenylmethyl (trityl), tetrahydropyranyl, vinyloxycarbonyl, ortho-nitrophenylsulfenyl, diphenylphosphinyl, para-toluenesulfonyl (Ts), mesyl, trifluoromethanesulfonyl, and benzyl. Examples of procedures for removal of

protecting groups include hydrogenolysis of CBZ groups using, for example, hydrogen gas at 50 psi in the presence of a hydrogenation catalyst such as 10% palladium on carbon, acidolysis of BOC groups using, for example, hydrogen chloride in dichloromethane, trifluoroacetic acid (TFA) in dichloromethane, and
5 the like, reaction of silyl groups with fluoride ions, and reductive cleavage of TCEC groups with zinc metal.

A general synthesis of the compounds of Formula I wherein Y^1 is $C(O)$ and Y^3 and Y^2 are taken together to form $C=C(R^3)$ is outlined below in Scheme 1.

In Scheme 1, commercially available compound (1) is allowed to react
10 with formamidine, for example, formamidine acetate, at elevated temperatures to give cyclic compound (2). Compound (2) is alkylated at N-3 to give compound (3), and the benzyl protecting group is removed to give compound (4). Compound (4) may be used to prepare 6-(3-substituted prop-2-ynyl) compounds of Formula I such as compound (5) or, alternatively, carbonylated derivatives of Formula I such
15 as compound (6).

Scheme 1.



wherein R¹ is as defined above for Formula I and R^{2a} is a subset of the group R² for Formula I.


5 The synthesis of a compound of Formula I wherein Y¹ is C(O), Y² is C(O), and Y³ is N(R⁴) is outlined below in Scheme 2.

In Scheme 2, commercially available compound (1) is optionally allowed to react with formaldehyde under reducing conditions to give cyclic compound (2). Compound (2) is cyclized to give compound (3). Compound (3) is alkylated at N-3 to give compound (4), and the benzyl protecting group is removed to give compound (5). Compound (5) may be used to prepare 6-(3-substituted prop-2-

10

ynyl) compounds of Formula I such as compound (6) or, alternatively, carbonylated derivatives of Formula I such as compound (7).

Scheme 2.




wherein R¹ is as defined above for Formula I and R^{2a} is a subset of the group R² for Formula I.





- 5 The synthesis of a compound of Formula I wherein Y¹ is CH₂, Y² is C(O), and Y³ is N(R⁴) is outlined below in Scheme 3.

5 In Scheme 3, commercially available compound (1) is optionally allowed to react with formaldehyde under reducing conditions to give cyclic compound (2). Compound (2) is cyclized to give compound (3). Compound (3) is alkylated at N-3 to give compound (4), and the benzyl protecting group is removed to give compound (5). Compound (5) may be used to prepare 6-(3-substituted prop-2-ynyl) compounds of Formula I such as compound (6) or, alternatively, carbonylated derivatives of Formula I such as compound (7).

Scheme 3.



Commercially available




wherein R¹ is as defined above for Formula I and R^{2a} is a subset of the group R² for Formula I.

The synthesis of a compound of Formula I wherein Y^1 is CH_2 and Y^3 and Y^2 are taken together to form $C=C(R^3)$ is outlined below in Scheme 4.

5 In Scheme 4, commercially available compound (1) is hydrolyzed to give compound (2). Compound (2) is cyclized to give compound (3). Compound (3) is alkylated at N-3 to give compound (4), and the benzyl protecting group is removed to give compound (5). Compound (5) may be used to prepare 6-(3-substituted prop-2-ynyl) compounds of Formula I such as compound (6) or, alternatively, carbonylated derivatives of Formula I such as compound (7).

Scheme 4.




wherein R¹ is as defined above for Formula I and R^{2a} is a subset of the group R² for Formula I.

5

It should be appreciated that when Q is trans-(H)C=C(H), cis-(H)C=C(H), C≡C, CH₂C≡C, or CF₂C≡C and is bonded to a sp² carbon atom in Formula I, a

palladium catalyzed coupling of the corresponding terminal olefin or alkyne of formulas R^1 -(trans-(H)C=CH₂), R^1 -(cis-(H)C=CH₂), R^1 -C≡CH, R^1 -CH₂C≡CH, or R^1 -CF₂C≡CH, wherein R^1 is as defined above, with a bromo- or iodo-substituted sp^2 carbon atom of formula:




5

in the presence of a suitable base will yield a compound of Formula I wherein Q is trans-(H)C=C(H), cis-(H)C=C(H), C≡C, CH₂C≡C, or CF₂C≡C and D is a group that is bonded to Q at a sp^2 carbon atom, and R^1 , V, and R^2 are as defined above for Formula I. Illustrative examples of the coupling reagents and catalysts include palladium tetrakis(triphenylphosphine) or palladium(II) acetate as catalyst, a tertiary organic amine base such as triethylamine or diisopropylethylamine, a suitable solvent such as dimethylformamide ("DMF") or tetrahydrofuran ("THF"), and optionally a co-catalyst such as copper(I)iodide, at a suitable temperature such as from 0°C to 100°C, for a suitable time such as from 30 minutes to 2 days, and under an inert atmosphere such as an atmosphere of nitrogen or argon gas.

10

15


Alternatively, a corresponding aldehyde of formula



prepared as described below, may be coupled with a phosphonium ylide under Wittig olefination, or Horner-Emmons olefination, conditions to give a compound of Formula I wherein Q is trans-(H)C=C(H).

20

The bromo or iodo intermediates described above may be converted by conventional means to the corresponding carboxylic acid of formula




and the carboxylic acid converted by conventional means to compounds of Formula I wherein Q is OC(O), CH(R⁶)C(O), OC(NR⁶), CH(R⁶)C(NR⁶), N(R⁶)C(O), N(R⁶)C(S), N(R⁶)C(NR⁶), SC(O), CH(R⁶)C(S), or SC(NR⁶).

Illustrative examples include coupling of the carboxylic acid with an amine to


5 provide a compound of Formula I wherein Q is N(R⁶)C(O), and optionally sulfurating the resulting amide with, for example P₂S₅ to provide a compound of Formula I wherein Q is N(R⁶)C(S). Alternatively, the carboxylic acid may be coupled with an alcohol to provide a compound of Formula I wherein Q is OC(O).

10 Alternatively, the carboxylic acid may be reduced to the corresponding hydroxymethyl compound of formula



and the hydroxymethyl converted to a compound of Formula I wherein Q is OCH₂ or N(R⁶)CH₂ by conventional means.


15 Alternatively, the hydroxymethyl compound may be oxidized to the corresponding aldehyde of formula



and the aldehyde coupled with hydroxylamine to give a corresponding oxime. The oxime may be chlorinated, and the chlorooxime cyclized with an olefin or alkyne to give a compound of Formula I wherein Q is a 5-membered heteroarylene.


20 Alternatively, the aldehyde may be prepared from the corresponding carboxylic acid by coupling the carboxylic acid with N,O-dimethylhydroxylamine and reducing the resulting dimethylhydroxamide with a suitable hydride reducing agent such as sodium borohydride or lithium aluminum hydride.

25 Alternatively, the above-described carboxylic acid intermediate may be converted by conventional means to the corresponding methyl ketone of formula



and the methyl ketone may be halogenated on methyl and coupled with various amines, alcohols, or other halogenated compounds to give a compound of Formula I wherein Q is $\text{CH}(\text{R}^6)\text{C}(\text{O})$.

- 5 Alternatively, the above-described carboxylic acid intermediate or bromo- or iodo-intermediates may be converted by conventional means to the corresponding nitrile of formula



- and the nitrile condensed with an amine or alcohol under non-nucleophilic basic conditions (e.g., 1,8-diazaundecane) to give a compound of Formula I wherein Q is $\text{N}(\text{R}^6)\text{C}(\text{NR}^6)$ or $\text{OC}(\text{NR}^6)$, respectively.
- 10

Alternatively, compounds of Formula I wherein Q is a lactam diradical may be prepared by conventional means by cyclizing the corresponding gamma-amino acids.

- 15 The compounds of Formula I can be evaluated in standard assays for their ability to inhibit the catalytic activity of MMP enzymes. The assays used to evaluate the MMP biological activity of the invention compounds are well-known and routinely used by those skilled in the study of MMP inhibitors and their use to treat clinical conditions. For example, compounds of Formula I may be readily
- 20 identified by assaying a test compound for inhibition of MMP-13 according to Biological Methods 1 or 2, and further assaying the test compound for allosteric inhibition of MMP-13 according to Biological Methods 3 or 4, as described below.

- 25 The compounds of Formula I will be shown to be potent inhibitors of MMP-13 catalytic domain. Potencies, as measured by IC_{50} 's, with MMP-13

catalytic domain for the invention compounds will typically range from about 0.001 μM to about 30 μM .

5 Invention compounds can be further screened with full-length MMP-2, full-length MMP-7, full-length MMP-9, and MMP-14 catalytic domain to determine selectivity of the inhibitors with MMP-13 versus the other MMP enzymes also. Selectivities of the invention compounds for MMP-13 catalytic domain versus another MMP enzyme (full-length or catalytic domain), as determined by dividing the IC_{50} for the inhibitor with a comparator MMP enzyme by the IC_{50} of the inhibitor with MMP-13 catalytic domain, are expected to range
10 from 5 to 50,000 fold.

To determine the inhibitory profiles, a compound of Formula I, or a pharmaceutically acceptable salt thereof, may be evaluated in standard assays for their ability to inhibit the catalytic activity of various MMP enzymes. The assays used to evaluate the MMP biological activity of the invention compounds are
15 well-known and routinely used by those skilled in the study of MMP inhibitors and their use to treat clinical conditions. The compound of Formula I will be shown to be selective for inhibition of MMP-13CD versus MMP-1FL, MMP-2FL, MMP-3CD, MMP-7FL, MMP-9FL, MMP-12CD, and MMP-14CD with typical selectivity ranging between about 50 and about 500 fold, as measured by dividing
20 the IC_{50} of the compound of Formula I with MMP-1FL, MMP-2FL, MMP-3CD, MMP-7FL, MMP-9FL, MMP-12CD, or MMP-14CD by the IC_{50} of the compound of Formula I with MMP-13CD.

The assays measure the amount by which a test compound reduces the hydrolysis of a thiopeptolide substrate catalyzed by a matrix metalloproteinase
25 enzyme. Such assays are described in detail by Ye et al., in *Biochemistry*, 1992;31(45):11231-11235, which is incorporated herein by reference. One such assay is described below in Biological Method 1.

Some of the particular methods described below use the catalytic domain of the MMP-13 enzyme, namely matrix metalloproteinase-13 catalytic domain
30 ("MMP-13CD"), rather than the corresponding full-length enzyme, MMP-13. It has been shown previously by Ye Qi-Zhuang, Hupe D., and Johnson L. (*Current Medicinal Chemistry*, 1996;3:407-418) that inhibitor activity against a catalytic

domain of an MMP is predictive of the inhibitor activity against the respective full-length MMP enzyme.

BIOLOGICAL METHOD 1

5

Thiopeptolide substrates show virtually no decomposition or hydrolysis at or below neutral pH in the absence of a matrix metalloproteinase enzyme. A typical thiopeptolide substrate commonly utilized for assays is Ac-Pro-Leu-Gly-thioester-Leu-Leu-Gly-OEt. A 100 μ L assay mixture will contain 50 mM of N-2-hydroxyethylpiperazine-N'-2-ethanesulfonic acid buffer ("HEPES," pH 7.0), 10 mM CaCl_2 , 100 μ M thiopeptolide substrate, and 1 mM 5,5'-dithio-bis-(2-nitrobenzoic acid) (DTNB). The thiopeptolide substrate concentration may be varied, for example from 10 to 800 μ M to obtain K_m and K_{cat} values. The change in absorbance at 405 nm is monitored on a Thermo Max microplate reader (molecular Devices, Menlo Park, CA) at room temperature (22°C). The calculation of the amount of hydrolysis of the thiopeptolide substrate is based on $E_{412} = 13600 \text{ M}^{-1} \text{ cm}^{-1}$ for the DTNB-derived product 3-carboxy-4-nitrothiophenoxide. Assays are carried out with and without matrix metalloproteinase inhibitor compounds, and the amount of hydrolysis is compared for a determination of inhibitory activity of the test compounds.

20

Test compounds were evaluated at various concentrations in order to determine their respective IC_{50} values, the micromolar concentration of compound required to cause a 50% inhibition of catalytic activity of the respective enzyme.

25

It should be appreciated that the assay buffer used with MMP-3CD was 50 mM N-morpholinoethane sulfonate ("MES") at pH 6.0 rather than the HEPES buffer at pH 7.0 described above.

30

The test described above for the inhibition of MMP-13 may also be adapted and used to determine the ability of the compounds of Formula I to inhibit the matrix metalloproteases MMP-1, MMP-2, MMP-3, MMP-7, MMP-9, MMP-12 and MMP-14.

BIOLOGICAL METHOD 2

Some representative compounds of Formula I have been evaluated for their ability to inhibit MMP-13. Inhibitor activity versus other MMPs with the compounds may be determined using, for example, MMP-1FL, which refers to full length interstitial collagenase; MMP-2FL, which refers to full length Gelatinase A; MMP-3CD, which refers to the catalytic domain of stromelysin; MMP-7FL, which refers to full length matrilysin; MMP-9FL, which refers to full length Gelatinase B; MMP-13CD, which refers to the catalytic domain of collagenase 3; and MMP-14CD, which refers to the catalytic domain of MMP-14. Test compounds can be evaluated at various concentrations in order to determine their respective IC₅₀ values, the micromolar concentration of compound required to cause a 50% inhibition of the hydrolytic activity of the respective enzyme.

The results of the above assays with other MMPs will establish that the compounds of Formula I are potent inhibitors of MMP enzymes, and are especially useful due to their selective inhibition of MMP-13. Because of this potent and selective inhibitory activity, the compounds are especially useful to treat diseases mediated by the MMP enzymes.

Allosteric inhibitors of MMP-13 which are compounds of Formula I may be readily identified by assaying a test compound for inhibition of MMP-13 according to the methods described below in Biological Methods 3 and 4.

BIOLOGICAL METHOD 3

Fluorogenic peptide-1 substrate based assay for identifying compounds of Formula I as allosteric inhibitors of MMP-13:

Final assay conditions:

50 mM HEPES buffer (pH 7.0)

10 mM CaCl₂

10 μ M fluorogenic peptide-1 ("FP1") substrate

0 or 15 mM acetohydroxamic acid (AcNHOH) = 1 K_d

2% DMSO (with or without inhibitor test compound)

0.5 nM MMP-13CD enzyme

Stock solutions:

- 1) 10X assay buffer: 500 mM HEPES buffer (pH 7.0) plus 100 mM CaCl_2
- 2) 10 mM FP1 substrate: (Mca)-Pro-Leu-Gly-Leu-(Dnp)-Dpa-Ala-Arg- NH_2
(Bachem, M-1895; "A novel coumarin-labeled peptide for sensitive
5 continuous assays of the matrix metalloproteinases," Knight C.G.,
Willenbrock F., and Murphy, G., FEBS Lett., 1992;296:263-266). Is prepared
10 mM stock by dissolving 5 mg FP1 in 0.457 mL DMSO.
- 3) 3 M AcNHOH: Is prepared by adding 4 mL H_2O and 1 mL 10X assay buffer
to 2.25 g AcNHOH (Aldrich 15,903-4). Adjusting pH to 7.0 with NaOH.
10 Diluting volume to 10 mL with H_2O . Final solution will contain 3 M
AcNHOH, 50 mM HEPES buffer (pH 7.0), and 10 mM CaCl_2 .
- 4) AcNHOH dilution buffer: 50 mM HEPES buffer (pH 7.0) plus 10 mM CaCl_2
- 5) MMP-13CD enzyme: Stock concentration = 250 nM.
- 6) Enzyme dilution buffer: 50 mM HEPES buffer (pH 7.0), 10 mM CaCl_2 , and
15 0.005% BRIJ 35 detergent (Calbiochem 203728; Protein Grade, 10%)

Procedure (for one 96-well microplate):

A. Prepared assay mixture:

- 1100 μL 10X assay buffer
- 11 μL 10 mM FP1
- 20 55 μL 3 M AcNHOH or 55 μL AcNHOH dilution buffer
- 8500 μL H_2O

B. Diluted MMP-13CD to 5 nM working stock:

- 22 μL MMP-13CD (250 nM)
- 1078 μL enzyme dilution buffer

25 C. Ran kinetic assay:

1. Dispense 2 μL inhibitor test sample (in 100% DMSO) into well.
2. Add 88 μL assay mixture and mix well, avoiding bubbles.
3. Initiate reactions with 10 μL of 5 nM MMP-13CD; mix well, avoid bubbles.
4. Immediately measure the kinetics of the reactions at room temperature.

Fluorimeter: F_{max} Fluorescence Microplate Reader & SOFTMAX PRO
Version 1.1 software (Molecular Devices Corporation; Sunnyvale, CA
94089).

Protocol menu:

5 excitation: 320 nm emission: 405 nm
 run time: 15 min interval: 29 sec
 RFU min: -10 RFU max: 200
 V_{max} points: 32/32

D. Compared % of control activity and/or IC₅₀ with inhibitor test compound
10 ±AcNHOH.

Hydrolysis of the fluorogenic peptide-1 substrate, [(Mca)Pro-Leu-Gly-Leu-
Dpa-Ala-Arg-NH₂; Bachem, catalog number M-1895], wherein “Mca” is
(7-methoxy-coumarin-4-yl)acetyl and “Dpa” is (3-[2,4-dinitrophenyl]-L-
2,3-diaminopropionyl), is used to screen for MMP-13 catalytic domain (CD)
15 inhibitors. (Dpa may also be abbreviated as “Dnp”.) Reactions (100 µL) contain
0.05 M Hepes buffer (pH 7), 0.01 M calcium chloride, 0.005% polyoxyethylene
(23) lauryl ether (“Brij 35”), 0 or 15 mM acetohydroxamic acid, 10 µM FP1, and
0.1 mM to 0.5 nM inhibitor in DMSO (2% final).

After recombinant human MMP-13CD (0.5 nM final) is added to initiate
20 the reaction, the initial velocity of FP1 hydrolysis is determined by monitoring the
increase in fluorescence at 405 nm (upon excitation at 320 nm) continuously for
up to 30 minutes on a microplate reader at room temperature. Alternatively, an
endpoint read can also be used to determine reaction velocity provided the initial
fluorescence of the solution, as recorded before addition of enzyme, is subtracted
25 from the final fluorescence of the reaction mixture. The inhibitor is assayed at
different concentration values, such as, for example, 100 µM, 10 µM, 1 µM,
100 nM, 10 nM, and 1 nM. Then the inhibitor concentration is plotted on the
X-axis against the percentage of control activity observed for inhibited
experiments versus uninhibited experiments (i.e., (velocity with inhibitor) divided
30 by (velocity without inhibitor) × 100) on the Y-axis to determine IC₅₀ values.
This determination is done for experiments done in the presence, and experiments
done in the absence, of acetohydroxamic acid. Data are fit to the equation: percent

control activity = $100/[1+([I]/IC_{50})^{\text{slope}}]$, where [I] is the inhibitor concentration, IC_{50} is the concentration of inhibitor where the reaction rate is 50% inhibited relative to the control, and slope is the slope of the IC_{50} curve at the curve's inflection point, using nonlinear least-squares curve-fitting equation regression.

Results may be expressed as an IC_{50} Ratio (+/-) ratio, which means a ratio of the IC_{50} of the inhibitor with MMP-13 and an inhibitor to the catalytic zinc of MMP-13, divided by the IC_{50} of the inhibitor with MMP-13 without the inhibitor to the catalytic zinc of MMP-13. Compounds of Formula I which are allosteric inhibitors of MMP-13 are expected to have an IC_{50} Ratio (+/-) ratio of less than 1, and are expected to be synergistic with the inhibitor to the catalytic zinc of MMP-13 such as, for example, AcNHOH. Compounds of Formula I which are not allosteric inhibitors of MMP-13 will be inactive in the assay or will have an IC_{50} Ratio (+/-) of greater than 1, unless otherwise indicated. Results can be confirmed by kinetics experiments which are well known in the biochemical art.

BIOLOGICAL METHOD 4

Fluorogenic peptide-1 based assay for identifying allosteric inhibitors of matrix metalloproteinase-13 catalytic domain ("MMP-13CD"):

In a manner similar to Biological Method 3, an assay is run wherein 1,10-phenanthroline is substituted for acetohydroxamic acid to identify compounds of Formula ICD.

Animal models may be used to establish that the instant compounds of Formula I, or a pharmaceutically acceptable salt thereof, would be useful for preventing, treating, and inhibiting cartilage damage, and thus for treating osteoarthritis, for example. Examples of such animal models are described below in Biological Methods 5 and 6.

BIOLOGICAL METHOD 5

Monosodium Iodoacetate-induced Osteoarthritis in Rat Model of Cartilage Damage (“MIA Rat”):

One end result of the induction of osteoarthritis in this model, as determined by histologic analysis, is the development of an osteoarthritic condition within the affected joint, as characterized by the loss of Toluidine blue staining and formation of osteophytes. Associated with the histologic changes is a concentration-dependent degradation of joint cartilage, as evidenced by effects on hind-paw weight distribution of the limb containing the affected joint, the presence of increased amounts of proteoglycan or hydroxyproline in the joint upon biochemical analysis, or histopathological analysis of the osteoarthritic lesions.

Generally, In the MIA Rat model on Day 0, the hind-paw weight differential between the right arthritic joint and the left healthy joint of male Wistar rats (150 g) are determined with an incapacitance tester, model 2KG (Linton Instrumentation, Norfolk, United Kingdom). The incapacitance tester has a chamber on top with an outwardly sloping front wall that supports a rat's front limbs, and two weight sensing pads, one for each hind paw, that facilitates this determination. Then the rats are anesthetized with isofluorine, and the right, hind leg knee joint is injected with 1.0 mg of mono-iodoacetate (“MIA”) through the infrapatellar ligament. Injection of MIA into the joint results in the inhibition of glycolysis and eventual death of surrounding chondrocytes. The rats are further administered either an invention compound or vehicle (in the instant case, water) daily for 14 days or 28 days. The invention compound is typically administered at a dose of 30 mg per kilogram of rat per day (30 mg/kg/day), but the invention compound may be administered at other doses such as, for example, 10 mg/kg/day, 60 mg/kg/day, 90-mg/kg/day, or 100 mg/kg/day according to the requirements of the compound being studied. It is well within the level of ordinary skill in the pharmaceutical arts to determine a proper dosage of an invention compound in this model. Administration of the invention compound in this model is optionally by oral administration or intravenous administration via an osmotic pump. After 7 and 14 days for a two-week study, or 7, 14, and 28 days for a four-week study, the hind-paw weight distribution is again determined. Typically, the animals administered vehicle alone place greater weight on their unaffected left

hind paw than on their right hind paw, while animals administered an invention compound show a more normal (i.e., more like a healthy animal) weight distribution between their hind paws. This change in weight distribution was proportional to the degree of joint cartilage damage. Percent inhibition of a change in hind paw joint function is calculated as the percent change in hind-paw weight distribution for treated animals versus control animals. For example, for a two week study,

Percent inhibition of a change in hind paw weight distribution

$$= \left\{ 1 - \left[\frac{(\Delta W_G)}{(\Delta W_C)} \right] \right\} \times 100$$

wherein: ΔW_C is the hind-paw weight differential between the healthy left limb and the arthritic limb of the control animal administered vehicle alone, as measured on Day 14; and

ΔW_G is the hind-paw weight differential between the healthy left limb and the arthritic limb of the animal administered an invention compound, as measured on Day 14.

In order to measure biochemical or histopathological end points in the MIA Rat model, some of the animals in the above study may be sacrificed, and the amounts of free proteoglycan in both the osteoarthritic right knee joint and the contralateral left knee joint may be determined by biochemical analysis. The amount of free proteoglycan in the contralateral left knee joint provides a baseline value for the amount of free proteoglycan in a healthy joint. The amount of proteoglycan in the osteoarthritic right knee joint in animals administered an invention compound, and the amount of proteoglycan in the osteoarthritic right knee joint in animals administered vehicle alone, are independently compared to the amount of proteoglycan in the contralateral left knee joint. The amounts of proteoglycan lost in the osteoarthritic right knee joints are expressed as percent loss of proteoglycan compared to the contralateral left knee joint control. The percent inhibition of proteoglycan loss, may be calculated as $\{[(\text{proteoglycan loss from joint (\%)} \text{ with vehicle}) - (\text{proteoglycan loss from joint with an invention compound})] \div (\text{proteoglycan loss from joint (\%)} \text{ with vehicle})\} \times 100$.

The MIA Rat data that are expected from the analysis of proteoglycan loss would establish that an invention compound is effective for inhibiting cartilage damage and inflammation and/or alleviating pain in mammalian patients, including human.

5 The results of these studies with oral dosing may be presented in tabular format in the columns labelled “IJFL (%+/- SEM)”, wherein IJFL means Inhibition of Joint Function Limitation, “SDCES”, wherein SDCES means Significant Decrease In Cartilage Erosion Severity, and “SIJWHLE”, wherein SIJWHLE means Significant Increase in Joints Without Hind Limb Erosion.

10 The proportion of subjects without hind limb erosions may be analyzed via an *Exact Sequential Cochran-Armitage Trend* test (SAS® Institute, 1999). The Cochran-Armitage Trend test is employed when one wishes to determine whether the proportion of positive or “Yes” responders increases or decreases with increasing levels of treatment. For the particular study, it is expected that the
15 number of animals without joint erosions increased with increasing dose.

 The ridit analysis may be used to determine differences in overall erosion severity. This parameter takes into account both the erosion grade (0 = no erosion, I = erosion extending into the superficial or middle layers, or II = deep layer erosion), and area (small, medium and large, quantified by dividing the area
20 of the largest erosion in each score into thirds) simultaneously. The analysis recognizes that each unit of severity is different, but does not assume a mathematical relationship between units.

 Another animal model for measuring effects of an invention compound on cartilage damage and inflammation and/or pain is described below in Biological
25 Method 6.

BIOLOGICAL METHOD 6

Induction of Experimental Osteoarthritis in Rabbit (“EOA in Rabbit”):

 Normal rabbits are anaesthetized and anteromedial incisions of the right knees performed. The anterior cruciate ligaments are visualized and sectioned.
30 The wounds are closed and the animals are housed in individual cages, exercised, and fed ad libitum. Rabbits are given either vehicle (water) or an invention

compound dosed three times per day with 30-mg/kg/dose or 10-mg/kg/dose. The invention compound may be administered at other doses such as, for example, 3 times 20 mg/kg/day or 3 times 60 mg/kg/day according to the requirements of the invention compound being studied. The rabbits are euthanized 8 weeks after surgery and the proximal end of the tibia and the distal end of the femur are removed from each animal.

Macroscopic Grading

The cartilage changes on the femoral condyles and tibial plateaus are graded separately under a dissecting microscope (Stereozoom, Bausch & Lomb, Rochester, NY). The depth of erosion is graded on a scale of 0 to 4 as follows: grade 0 = normal surface; Grade 1 = minimal fibrillation or a slight yellowish discoloration of the surface; Grade 2 = erosion extending into superficial or middle layers only; Grade 3 = erosion extending into deep layers; Grade 4 = erosion extending to subchondral bone. The surface area changes are measured and expressed in mm². Representative specimens may also be used for histologic grading (see below).

Histologic Grading

Histologic evaluation is performed on sagittal sections of cartilage from the lesional areas of the femoral condyle and tibial plateau. Serial sections (5 µm) are prepared and stained with safranin-O. The severity of OA lesions is graded on a scale of 0 - 14 by two independent observers using the histologic-histochemical scale of Mankin *et al.* This scale evaluates the severity of OA lesions based on the loss of safranin-O staining (scale 0 - 4), cellular changes (scale 0 - 3), invasion of tidemark by blood vessels (scale 0 - 1) and structural changes (scale 0 - 6). On this latter scale, 0 indicates normal cartilage structure and 6 indicates erosion of the cartilage down to the subchondral bone. The scoring system is based on the most severe histologic changes in the multiple sections.

Representative specimens of synovial membrane from the medial and lateral knee compartments are dissected from underlying tissues. The specimens are fixed, embedded, and sectioned (5 µm) as above, and stained with hematoxylin-eosin. For each compartment, two synovial membrane specimens are

examined for scoring purposes and the highest score from each compartment is retained. The average score is calculated and considered as a unit for the whole knee. The severity of synovitis is graded on a scale of 0 to 10 by two independent observers, adding the scores of 3 histologic criteria: synovial lining cell
5 hyperplasia (scale 0 - 2); villous hyperplasia (scale 0 - 3); and degree of cellular infiltration by mononuclear and polymorphonuclear cells (scale 0 - 5): 0 indicates normal structure.

Statistical Analysis

Mean values and SEM is calculated and statistical analysis was done using
10 the Mann-Whitney U-test.

The results of these studies would be expected to show that an invention compound would reduce the size of the lesion on the tibial plateaus, and perhaps the damage in the tibia or on the femoral condyles. In conclusion, these results would show that an invention compound would have significant inhibition effects
15 on the damage to cartilage.

The foregoing studies would establish that an invention compound is effective for the inhibition of cartilage damage and inflammation and/or alleviating pain, and thus useful for the treatment of osteoarthritis or rheumatoid arthritis in human, and other mammalian disorders. Such a treatment offers a
20 distinct advantage over existing treatments that only modify pain or inflammation or and other secondary symptoms. The effectiveness of an invention compound in this model would indicate that the invention compound will have clinically useful effects in preventing and/or treating cartilage damage, pain and/or inflammation.

Administration according to the invention method of an invention
25 compound to a mammal to treat the diseases listed above is preferably, although not necessarily, accomplished by administering the compound, or a salt thereof, in a pharmaceutical dosage form.

The compounds of Formula I, or a pharmaceutically acceptable salt thereof, can be prepared and administered according to the invention method in a
30 wide variety of oral and parenteral pharmaceutical dosage forms. Thus, the compounds of Formula I, or a pharmaceutically acceptable salt thereof, can be administered by injection, that is, intravenously, intramuscularly,

intracutaneously, subcutaneously, intraduodenally, or intraperitoneally. Also, the compounds of Formula I, or a pharmaceutically acceptable salt thereof, can be administered by inhalation, for example, intranasally. Additionally, the compounds of Formula I, or a pharmaceutically acceptable salt thereof, can be administered transdermally. It will be obvious to those skilled in the art that the following dosage forms may comprise as the active component an invention compound. The invention compounds generally are present in a concentration of about 5% to about 95% by weight of the formulation.

For preparing pharmaceutical compositions from the compounds of Formula I, or a pharmaceutically acceptable salt thereof, (i.e., the active component) pharmaceutically acceptable carriers can be either solid or liquid. Solid form preparations are preferred. Solid form preparations include powders, tablets, pills, capsules, cachets, suppositories, and dispersible granules. A solid carrier can be one or more substances which may also act as diluents, flavoring agents, solubilizers, lubricants, suspending agents, binders, preservatives, tablet disintegrating agents, or an encapsulating material.

In powders, the carrier is a finely divided solid which is in a mixture with the finely divided active component. Powders suitable for intravenous administration or administration by injection may be lyophilized.

In tablets, the active component is mixed with the carrier having the necessary binding properties in suitable proportions and compacted in the shape and size desired.

The powders and tablets preferably contain from about 5% to about 70%, total, of the active component. Suitable carriers are magnesium carbonate, magnesium stearate, talc, sugar, lactose, pectin, dextrin, starch, gelatin, tragacanth, methylcellulose, sodium carboxymethylcellulose, a low melting wax, cocoa butter, and the like. The term "preparation" is intended to include the formulation of the active component with encapsulating material as a carrier providing a capsule in which the active component, with or without other carriers, is surrounded by a carrier, which is thus in association with it. Similarly, cachets and lozenges are included. Tablets, powders, capsules, pills, cachets, and lozenges can be used as solid dosage forms suitable for oral administration.

For preparing suppositories, a low melting wax, such as a mixture of fatty acid glycerides or cocoa butter, is first melted and the active component is dispersed homogeneously therein, as by stirring. The molten homogenous mixture is then poured into convenient sized molds, allowed to cool, and thereby to solidify.

Liquid form preparations include solutions, suspensions, and emulsions, for example, water or water propylene glycol solutions. For parenteral injection, liquid preparations can be formulated in solution in aqueous polyethylene glycol solution.

Aqueous solutions suitable for oral use can be prepared by dissolving the active component in water and adding suitable colorants, flavors, stabilizing, and thickening agents as desired.

Aqueous suspensions suitable for oral use can be made by dispersing the finely divided active component in water with viscous material, such as natural or synthetic gums, resins, methylcellulose, sodium carboxymethylcellulose, and other well-known suspending agents.

Also included are solid form preparations which are intended to be converted, shortly before use, to liquid form preparations for oral administration. Such liquid forms include solutions, suspensions, and emulsions. These preparations may contain, in addition to the active component, colorants, flavors, stabilizers, buffers, artificial and natural sweeteners, dispersants, thickeners, solubilizing agents, and the like.

The pharmaceutical preparation is preferably in unit dosage form. In such form, the preparation is subdivided into unit doses containing an appropriate quantity of the active component. The unit dosage form can be a packaged preparation, the package containing discrete quantities of preparation, such as packeted tablets, capsules, and powders in vials or ampoules. Also, the unit dosage form can be a capsule, tablet, cachet, or lozenge itself, or it can be the appropriate number of any of these in packaged form.

The quantity of active component in a unit dose preparation may be varied or adjusted from 0.01 to 1000 mg, preferably 1 to 500 mg according to the particular application and the potency of the active components. The composition can, if desired, also contain other compatible therapeutic agents.

In therapeutic use as agents to treat the above-listed diseases, the compounds of Formula I, or a pharmaceutically acceptable salt thereof, are administered at a dose that is effective for treating at least one symptom of the disease or disorder being treated. The initial dosage of about 1 mg/kg to about 100 mg/kg daily of the active component will be effective. A daily dose range of about 25 mg/kg to about 75 mg/kg of the active component is preferred. The dosages, however, may be varied depending upon the requirements of the patient, the severity of the condition being treated, and the particular invention compound being employed in the invention combination. Determination of the proper dosage for a particular situation is within the skill of the art as described above. Typical dosages will be from about 0.1 mg/kg to about 500 mg/kg, and ideally about 25 mg/kg to about 250 mg/kg, such that it will be an amount that is effective to treat the particular disease or disorder being treated.

A preferred composition for dogs comprises an ingestible liquid peroral dosage form selected from the group consisting of a solution, suspension, emulsion, inverse emulsion, elixir, extract, tincture and concentrate, optionally to be added to the drinking water of the dog being treated. Any of these liquid dosage forms, when formulated in accordance with methods well known in the art, can either be administered directly to the dog being treated, or may be added to the drinking water of the dog being treated. The concentrate liquid form, on the other hand, is formulated to be added first to a given amount of water, from which an aliquot amount may be withdrawn for administration directly to the dog or addition to the drinking water of the dog.

A preferred composition provides delayed-, sustained- and/or controlled-release of an invention compound. Such preferred compositions include all such dosage forms which produce $\geq 40\%$ inhibition of cartilage degradation, and result in a plasma concentration of the active component of at least 3 fold the active component's ED_{40} for at least 2 hours; preferably for at least 4 hours; preferably for at least 8 hours; more preferably for at least 12 hours; more preferably still for at least 16 hours; even more preferably still for at least 20 hours; and most preferably for at least 24 hours. Preferably, there is included within the above-described dosage forms those which produce $\geq 40\%$ inhibition of cartilage

degradation, and result in a plasma concentration of the active component of at least 5 fold the active component's ED₄₀ for at least 2 hours, preferably for at least 2 hours, preferably for at least 8 hours, more preferably for at least 12 hours, still more preferably for at least 20 hours and most preferably for at least 24 hours.

5 More preferably, there is included the above-described dosage forms which produce $\geq 50\%$ inhibition of cartilage degradation, and result in a plasma concentration of the active component of at least 5 fold the active component's ED₄₀ for at least 2 hours, preferably for at least 4 hours, preferably for at least 8 hours, more preferably for at least 12 hours, still more preferably for at least 20

10 hours and most preferably for at least 24 hours.

The following Formulation Examples 1 to 8 illustrate the invention pharmaceutical compositions. When the formulations comprise the invention compound and a pharmaceutically acceptable carrier, diluent, or excipient, they contain a cartilage damage treating effective amount or a therapeutically effective

15 amount such as, for example, an anti-osteoarthritic effective amount of the invention compound. The examples are representative only, and are not to be construed as limiting the invention in any respect.

FORMULATION EXAMPLE 1

Tablet Formulation:

Ingredient	Amount (mg)
An invention compound	25
Lactose	50
Cornstarch (for mix)	10
Cornstarch (paste)	10
Magnesium stearate (1%)	5
Total	100

20 The invention compound, lactose, and cornstarch (for mix) are blended to uniformity. The cornstarch (for paste) is suspended in 200 mL of water and heated with stirring to form a paste. The paste is used to granulate the mixed powders. The wet granules are passed through a No. 8 hand screen and dried at 80°C. The dry granules are lubricated with the 1% magnesium stearate and pressed into a

tablet. Such tablets can be administered to a human from one to four times a day for inhibiting cartilage damage or treating osteoarthritis.

FORMULATION EXAMPLE 2

Coated Tablets:

- 5 The tablets of Formulation Example 1 are coated in a customary manner with a coating of sucrose, potato starch, talc, tragacanth, and colorant.

FORMULATION EXAMPLE 3

Injection vials:

- 10 The pH of a solution of 500 g of an invention compound and 5 g of disodium hydrogen phosphate is adjusted to pH 6.5 in 3 L of double-distilled water using 2 M hydrochloric acid. The solution is sterile filtered, and the filtrate is filled into injection vials, lyophilized under sterile conditions, and aseptically sealed. Each injection vial contains 25 mg of the invention compound.

FORMULATION EXAMPLE 4

- 15 Suppositories:

A mixture of 25 g of an invention compound, 100 g of soya lecithin, and 1400 g of cocoa butter is fused, poured into molds, and allowed to cool. Each suppository contains 25 mg of the invention compound.

FORMULATION EXAMPLE 5

- 20 Solution:

- 25 A solution is prepared from 1 g of an invention compound, 9.38 g of $\text{NaH}_2\text{PO}_4 \cdot 12\text{H}_2\text{O}$, 28.48 g of $\text{Na}_2\text{HPO}_4 \cdot 12\text{H}_2\text{O}$, and 0.1 g benzalkonium chloride in 940 mL of double-distilled water. The pH of the solution is adjusted to pH 6.8 using 2 M hydrochloric acid. The solution is diluted to 1.0 L with double-distilled water, and sterilized by irradiation. A 25 mL volume of the solution contains 25 mg of the invention compound.

FORMULATION EXAMPLE 6

Ointment:

500 mg of an invention compound is mixed with 99.5 g of petroleum jelly under aseptic conditions. A 5 g portion of the ointment contains 25 mg of the invention compound.

FORMULATION EXAMPLE 7

Capsules:

2 kg of an invention compound are filled into hard gelatin capsules in a customary manner such that each capsule contains 25 mg of the invention compound.

FORMULATION EXAMPLE 8

Ampoules:

A solution of 2.5 kg of an invention compound is dissolved in 60 L of double-distilled water. The solution is sterile filtered, and the filtrate is filled into ampoules. The ampoules are lyophilized under sterile conditions and aseptically sealed. Each ampoule contains 25 mg of the invention compound.

The following Formulation Examples 9 to 16 illustrate the invention pharmaceutical compositions containing an invention combination in a single formulation with a pharmaceutically acceptable carrier, diluent, or excipient. The examples are representative only, and are not to be construed as limiting the invention in any respect.

FORMULATION EXAMPLE 9

Tablet Formulation:

Ingredient	Amount (mg)
An invention compound	25
A COX-2 inhibitor	20
Lactose	50
Cornstarch (for mix)	10
Cornstarch (paste)	10
Magnesium stearate (1%)	5
Total	120

The invention compound or COX-2 inhibitor, lactose, and cornstarch (for mix) are blended to uniformity. The cornstarch (for paste) is suspended in 200 mL of water and heated with stirring to form a paste. The paste is used to granulate the mixed powders. The wet granules are passed through a No. 8 hand screen and dried at 80°C. The dry granules are lubricated with the 1% magnesium stearate and pressed into a tablet. Such tablets can be administered to a human from one to four times a day for treatment of one of the above-listed diseases.

FORMULATION EXAMPLE 10

Coated Tablets:

The tablets of Formulation Example 9 are coated in a customary manner with a coating of sucrose, potato starch, talc, tragacanth, and colorant.

FORMULATION EXAMPLE 11

Injection vials:

The pH of a solution of 250 g of a COX-2 inhibitor, 500 g of an invention compound, and 5 g of disodium hydrogen phosphate is adjusted to pH 6.5 in 3 L of double-distilled water using 2 M hydrochloric acid. The solution is sterile filtered, and the filtrate is filled into injection vials, lyophilized under sterile conditions, and aseptically sealed. Each injection vial contains 12.5 mg of COX-2 inhibitor and 25 mg of the invention compound.

FORMULATION EXAMPLE 12

Suppositories:

A mixture of 50 g of a COX-2 inhibitor, 25 g of an invention compound, 100 g of soya lecithin, and 1400 g of cocoa butter is fused, poured into molds, and allowed to cool. Each suppository contains 50 mg of the COX-2 inhibitor and 25 mg of the invention compound.

FORMULATION EXAMPLE 13

Solution:

A solution is prepared from 0.5 g of a COX-2 inhibitor, 1 g of an invention compound, 9.38 g of $\text{NaH}_2\text{PO}_4 \cdot 12\text{H}_2\text{O}$, 28.48 g of $\text{Na}_2\text{HPO}_4 \cdot 12\text{H}_2\text{O}$, and 0.1 g benzalkonium chloride in 940 mL of double-distilled water. The pH of the solution is adjusted to pH 6.8 using 2 M hydrochloric acid. The solution is diluted to 1.0 L with double-distilled water, and sterilized by irradiation. A 25 mL volume of the solution contains 12.5 mg of the COX-2 inhibitor and 25 mg of the invention compound.

FORMULATION EXAMPLE 14

Ointment:

100 mg of a COX-2 inhibitor, 500 mg of an invention compound is mixed with 99.4 g of petroleum jelly under aseptic conditions. A 5 g portion of the ointment contains 5 mg of the COX-2 inhibitor and 25 mg of the invention compound.

FORMULATION EXAMPLE 15

Capsules:

2 kg of a COX-2 inhibitor and 20 kg of an invention compound are filled into hard gelatin capsules in a customary manner such that each capsule contains 25 mg of the COX-2 inhibitor and 250 mg of the invention compound.

FORMULATION EXAMPLE 16

Ampoules:

A solution of 2.5 kg of a COX-2 inhibitor and 2.5 kg of an invention compound is dissolved in 60 L of double-distilled water. The solution is sterile filtered, and the filtrate is filled into ampoules. The ampoules are lyophilized under sterile conditions and aseptically sealed. Each ampoule contains 25 mg each of the COX-2 inhibitor and the invention compound.

While it may be desirable to formulate a COX-2 inhibitor and an invention compound together in one capsule, tablet, ampoule, solution, and the like, for simultaneous administration, it is not necessary for the purposes of practicing the invention methods. A COX-2 inhibitor and an invention compound alternatively can each be formulated independently in any form such as, for example, those of any one Formulation Examples 1 to 16, and administered to a patient either simultaneously or at different times.

The following examples illustrate the invention pharmaceutical compositions containing discrete formulations of the active components of an invention combination and a pharmaceutically acceptable carrier, diluent, or excipient. The examples are representative only, and are not to be construed as limiting the invention in any respect.

FORMULATION EXAMPLE 17

Tablet Formulation of an invention compound:

Ingredient	Amount (mg)
An invention compound	25
Lactose	50
Cornstarch (for mix)	10
Cornstarch (paste)	10
Magnesium stearate (1%)	5
Total	100

An invention compound, lactose, and cornstarch (for mix) are blended to uniformity. The cornstarch (for paste) is suspended in 200 mL of water and heated

with stirring to form a paste. The paste is used to granulate the mixed powders. The wet granules are passed through a No. 8 hand screen and dried at 80°C. The dry granules are lubricated with the 1% magnesium stearate and pressed into a tablet.

5

Injection vial formulation of a COX-2 inhibitor:

The pH of a solution of 500 g of a COX-2 inhibitor and 5 g of disodium hydrogen phosphate is adjusted to pH 6.5 in 3 L of double-distilled water using 2 M hydrochloric acid. The solution is sterile filtered, and the filtrate is filled into
10 injection vials, lyophilized under sterile conditions, and aseptically sealed. Each injection vial contains 25 mg of the COX-2 inhibitor.

Such tablets containing the invention compound can be administered to a human from one to four times a day for treatment of the above-listed diseases, and the injection solutions containing the COX-2 inhibitor can be administered to a
15 human 1 or 2 times per day, wherein the administration by injection is optionally simultaneous with administration of the tablets or at different times, for the treatment of one of the above-listed diseases.

FORMULATION EXAMPLE 18

Coated Tablets containing an invention compound:

20 The tablets of Formulation Example 17 are coated in a customary manner with a coating of sucrose, potato starch, talc, tragacanth, and colorant.

Capsules containing valdecoxib or celecoxib:

2 kg of a COX-2 inhibitor are filled into hard gelatin capsules in a customary manner such that each capsule contains 25 mg of the COX-2 inhibitor.

25 Such coated tablets containing the invention compound can be administered to a human from one to four times a day for treatment of the above-listed diseases, and the capsules containing the COX-2 inhibitor can be administered to a human 1 or 2 times per day, wherein the administration of the capsules is optionally simultaneous with administration of the tablets or at
30 different times, for the treatment of one of the above-listed diseases.

Still further, it should be appreciated that the invention methods comprising administering an invention combination to a mammal to treat diseases

or disorders listed above may be used to treat different diseases simultaneously. For example, administration of a COX-2 inhibitor in accordance with the invention combination may be carried out as described above to treat inflammation, arthritic pain, pain associated with menstrual cramping, and
5 migraines, while an invention compound may be administered to treat OA or inhibit cartilage damage.

As shown above, the invention methods comprising administering an invention compound offer a distinct advantage over existing treatments for diseases such as OA that comprise cartilage damage, wherein the existing
10 treatments modify pain or secondary symptoms, but do not show a disease modifying effect.

While the invention has been described and illustrated with reference to certain particular embodiments thereof, those skilled in the art will appreciate that various adaptations, changes, modifications, substitutions, deletions, or additions
15 of procedures and protocols may be made without departing from the spirit and scope of the invention. It is intended, therefore, that the invention be defined by the scope of the claims that follow and that such claims be interpreted as broadly as is reasonable.

All references cited above are hereby incorporated by reference herein.

20 Having described the invention method, various embodiments of the invention are hereupon claimed.